- 15 Brimblecombe F S W, Cruikshank R, Masters P L, Reid D D, Stewart G T. Family studies of respiratory infections. *Br Med J* 1958; **i**: 119-28.
- 16 Tyrrell D A J. Clinical clues in virus infections. *Br Med J* 1963; **i**: 493-6.
- 17 Panush R S. Acute arthritis associated with febrile viral-like respiratory syndromes. *J Rheumatol* 1974; **1**: 299-307.
- 18 Rahal J J, Millian S J, Noriega E R. Coxsackie virus and adenovirus infection. Association with acute febrile and juvenile rheumatoid arthritis. *JAMA* 1976; **235**: 2496-501.
- 19 Isaacs D, Clarke J R, Tyrrell D A J, Webster A D B, Valman H B. Deficiency of production of leucocyte interferon (interferon- α) in vitro and in vivo in children with recurrent respiratory tract infections. *Lancet* 1981; **ii**: 950-2.
- 20 Bryson Y J, Winter H S, Gard S E, Fischer T J, Stiehm E R. Deficiency of immune interferon production by leukocytes of normal newborns. *Cell Immunol* 1980; **55**: 191-200.
- 21 Glynne L E. In: Dumonde D C, ed. *Infection and immunology in the rheumatic diseases*. Oxford: Blackwell, 1976: 75.
- 22 Frick O L, German D F, Mills J. Development of allergy in children. 1. Association with virus infections. *J Allergy Clin Immunol* 1979; **63**: 228-41.
- 23 Andersson J, Schreier M H, Melchers F. T-cell-dependent B-cell stimulation is H-2 restricted and antigen dependent only at the resting B-cell level. *Proc Natl Acad Sci USA* 1980; **77**: 1612-6.

Book review

Advances in Assessing Rheumatoid Arthritis: Symposium/Workshop at the London Hospital, 17-18 March 1983. British Journal of Rheumatology Supplement 1983. Available from Bailliere Tindall, 1 Vincent Square, London SW1, £8.00 including postage and packing.

I was fortunate to be invited to attend this symposium and commend its published proceedings for 2 reasons.

In the first place the design of the programme was imaginative and highly efficient, providing an excellent model for clinical workshops. Preconference questionnaires about the quality of clinical trial reports were distributed to the participants and results were analysed in detail at the meeting. At the opening session invited lecturers addressed 4 aspects of the problem—statistics, judgment analysis, radiology, and the assessment of disability. The meeting was then divided into a number of group discussions, each with its chairman, reporter, and introductory speakers, which thrashed out prescribed topics—definition and characterisation of rheumatoid arthritis, the nature of drugs to be tested, reporting clinical trials, multicentre trial management, and measurement of patient and other outcomes. In the final session the results of these group discussions were presented and debated. I can think of no better way for an active interchange of ideas.

Secondly, the ideas thrown up were interesting and sometimes provocative. For example, R. Peto, speaking from a wealth of experience, proposed that any controlled trial of disease-modifying drugs should involve several dozen rheumatologists studying 2000 patients over a 5-year period, but using very simple assessments of compliance, toxicity, and therapeutic benefit. C. R. B. Joyce and K. R. Hammond, using a mathematical model, showed how the different factors of knowledge and consistency can influence clinical judgment, a theme developed by other speakers. J. F. Fries discussed in depth modern techniques of

outcome assessment, in which disability is an important but by no means the only consideration. From the group discussion on the reporting of clinical trials emerged an important checklist of data which should be presented in such reports. The point was strongly made that most published clinical studies have been too seriously deficient in requirements to draw valid conclusions about therapeutic response. The interest generated by the symposium might perhaps have been enhanced by some consideration of assessing therapeutic agents other than drugs—physical treatment, surgery, or recent immunological *tour-de-force* such as lymphocyte depletion by one way or another, although the same principles of evaluation no doubt apply.

During the meeting I could not help reflecting on the comparison with another common rheumatic disorder—gout—in which I have taken an interest over the years. Here we have considerable knowledge about the causative agent, hyperuricaemia, which can be tidily measured in assessing the effect of 'disease-modifying' drugs such as allopurinol, with an almost certain knowledge of ultimate 'patient outcome'. It did not take 2000 patients to make the point: our original UK study on allopurinol dealt with 4 patients only, and was pretty conclusive. In rheumatoid arthritis we have no uric acid to estimate and certainly no allopurinol to deal with it. Time will change all this and one or two of our younger rheumatologists, chancing upon this dust-covered Supplement in years to come, will marvel at the skill and dedication at present being given to what is after all only the fringe of the problem. *Parturient montes, nascetur ridiculus mus*.

Nevertheless, we have to chase the wee mouse as he runs at the moment, and we should do it properly. The publication is well referenced and will undoubtedly be a definitive work on the subject for the next few years. It should be at the elbows of trialists and editors alike.

J. T. SCOTT