REPORT

Integrin alpha V beta 3 as a target for treatment of rheumatoid arthritis and related rheumatic diseases

R L Wilder

Ann Rheum Dis 2002;**61**(Suppl II):ii96–ii99

A substantial and persuasive body of data now exists that supports the view that integrin alpha V beta 3 plays a critical part in activated macrophage dependent inflammation, osteoclast development, migration, and bone resorption, and inflammatory angiogenesis. All of these processes play an important part in the pathogenesis of rheumatoid arthritis (RA) and related arthropathies. Animal arthritis model data further support these concepts and also suggest that therapeutic antagonism of integrin alpha V beta 3 is worthy of further investigation in RA and related arthropathies. To this end, Vitaxin, also known as MEDI-522, has been developed. Vitaxin is a humanised monoclonal IgG₁ antibody that specifically binds a conformational epitope formed by both the integrin alpha V and beta 3 subunits. It blocks the interaction of alpha V beta 3 with various ligands such as osteopontin and vitronectin. Clinical trials with Vitaxin in patients with RA are in progress.

Although medical treatment for rheumatic diseases such as rheumatoid arthritis (RA) has improved dramatically in recent years (most notably with the development of anti-tumour necrosis factor α agents used in combination with methotrexate), treatment for a substantial number of patients with RA remains suboptimal. Most patients fail to completely respond to these new treatments, and for many other patients, the beneficial effects of these treatments diminish with time. Many of patients with RA still will ultimately require costly joint arthroplasties as an additional effort to relieve pain and to improve or maintain functional abilities. In other words, the cost of the disease to the patients, to the patients' families, and to society remains extremely high. The continued development of new and more effective treatments is clearly warranted.

MEDI-522, also known as Vitaxin, is a new humanised monoclonal IgG₁ antibody that binds human integrin alpha V beta 3 that has now entered clinical development for RA and related diseases. MEDI-522 has a higher affinity for integrin alpha V beta 3 and is more stable than MEDI-523 (also known as Vitaxin 1), which has been phased out of clinical development. MEDI-522 was engineered from the murine monoclonal LM609, an antibody directed against a conformational epitope of the integrin alpha V beta 3 complex. In other words, it specifically recognises an epitope formed by both subunits of integrin alpha V and beta 3¹ and does not recognise the alpha V or beta 3 subunits alone. In this paper, the biology of integrin alpha V beta 3 and the rationale for its selection as a target for monoclonal antibody treatment of RA will be briefly reviewed.

INTEGRIN ALPHA V BETA 3: WHAT IS IT?

Integrins are heterodimeric transmembrane proteins formed by non-covalent association of alpha and beta subunits. Both

subunits are type 1 membrane proteins with large extracellular ectodomains and short cytoplasmic tails.² In mammals, the integrin family contains at least 18 alpha subunits that associate with at least nine beta subunits. The alpha and beta subunits assemble into at least 24 structurally distinct receptors.³ Ligands for these integrin receptors are diverse, but all use an acidic residue for recognition by the integrin. Specificity for an individual ligand is determined by additional contacts with the integrin. Ligand binding induces structural changes in the integrin and interactions with other molecules. In general, integrins serve important cellular biosensing roles and convey signals from both outside the cell to inside ("outside-in" signalling) and, vice versa, from inside the cell to outside ("inside-out" signalling).

The integrin alpha V (CD51) subunit can form heterodimers with at least five distinct beta subunits: beta 1, beta 3 (CD61), beta 5, beta 6, and beta 8. The integrin beta 3 subunit, on the other hand, can associate with not only alpha V but also alpha IIb subunits (CD41). This molecular association information is important from the perspective of designing and assessing potential therapeutic monoclonal antibodies. Antibodies that only target alpha V may affect the function of up to five distinct integrins, and antibodies that only target beta 3 may affect both integrins alpha V beta 3 and alpha IIb beta 3. Alpha IIb beta 3 is the major integrin on platelets. As noted above, Vitaxin recognises a specific conformational epitope formed by both the alpha V and beta 3 subunits. In other words, it does not bind integrins alpha V beta 5 or alpha IIb beta 3.

The integrin alpha V beta 3, also known as the vitronectin receptor, consists of a 125 kDa alpha V subunit and a 105 kDa beta 3 subunit. It has been the focus of intensive research because of its major role in several distinct processes, particularly osteoclast mediated bone resorption, angiogenesis and pathological neovascularisation, and tumour metastasis.^{4–12} Evidence of the intense interest in this integrin is provided by the recent determination of the crystal structure of the extracellular segment of integrin alpha V beta 3.¹³ This work shows that the heterodimeric molecule can exist in either an extended or "flexed" conformation. The available data indicate that the "flexed" or "bent" conformation represents the inactive form.¹⁴ In addition, the crystal structure of alpha V beta 3 binding an Arg-Gly-Asp ligand has also been determined.¹⁵

INTEGRIN ALPHA V BETA 3: WITH WHAT MOLECULES DOES IT ASSOCIATE?

Integrin alpha V beta 3 has distinct functional properties that are mediated through interactions with a variety of extracellular matrix (ECM) proteins in addition to vitronectin.⁶ These ECM proteins include osteopontin, fibronectin, fibrinogen, thrombospondin, proteolysed collagen, von Willebrand factor,

Abbreviations: RA, rheumatoid arthritis; ECM, extracellular matrix

and others. The specific nature of the extracellular interactions of integrin alpha V beta 3 with ECM has additional functional effects on the expression of other cell surface receptors. For example, culturing endothelial cells on vitronectin increases the presence of VEGFR1, as well as VEGFR2, FGFR1, and FGFR2 on the endothelial cell surface. Each of these receptors plays an important part mediating the effects of angiogenic growth factors such as VEGF, FGF1, and FGF2. The increased expression of VEGFR1 and the other receptors is blocked with if anti-alpha V beta 3 and alpha V beta 5 antibodies are added to the culture system. In contrast, plating endothelial cells on fibrin, instead of vitronectin, decreases the expression of these receptors.¹⁶ Thus, these integrins function to transmit signals into the cell, and the cell responds by changing its expression of surface receptors—that is, in this case, the expression of angiogenic growth factor receptors.

Integrin alpha V beta 3 not only modulates the levels of expression of selected cell surface molecules, it also physically associates with a number of important cell surface molecules including VEGFR2, IAP, MTI-MMP, MMP2, and CD47.^{17–21} The associations with these various cell surface molecules is modulated by alpha V beta 3 interactions with various ligands, and vice versa, implying that alpha V beta 3 plays a critical part in regulating the localisation or clustering of important cell surface molecules involved in cell adhesion, growth, survival, migration, and invasion through the ECM.

In addition to the ECM and cell surface membrane molecular associations, integrin alpha V beta 3 also interacts with a number of important intracellular signalling molecules. These include paxillin, focal adhesion kinase, caspase 8, and others.^{22–23} These interactions also play a part in regulating intracellular signalling, cell migration, cell proliferation, and cell survival.

INTEGRIN ALPHA V BETA 3: WHERE IS IT FOUND? WHAT TISSUES? WHAT CELLS?

Alpha V beta 3 is expressed at low levels in most normal tissues including intestinal, vascular, and smooth muscle cells, but, of particular interest, is that high level expression is limited to bone, the mid-menstrual cycle endometrium, placenta, inflammatory sites, and invasive tumours. The cell types that express high levels of this integrin include mature, bone resorbing osteoclasts and activated macrophages, a small fraction of neutrophils, angiogenic endothelial cells, and migrating smooth muscle cells.²⁴ These cell types clearly reflect the involvement of integrin alpha V beta 3 in bone resorption, neovascularisation, and inflammation. As pathological osteoclast mediated bone resorption, macrophage dependent inflammation and neovascularisation are characteristic features of diseases such as RA and related arthropathies, these observations provide the first line of supportive evidence that alpha V beta 3 may represent a rationale therapeutic target for these diseases.^{25–29} Pathological bone resorption and neovascularisation are also characteristic features of many malignant tumours. Moreover, expression of integrin alpha V beta 3 is also characteristic of several invasive tumour types including melanoma, glioma, ovarian, and breast cancer.^{7–30–33} These observations suggest that pharmacological antagonism of the integrin alpha V beta 3 may represent, in addition to rheumatic diseases such as RA, a rational approach to treatment of cancer.

INTEGRIN ALPHA V BETA 3: WHAT DOES IT DO? BIOLOGICAL FUNCTIONS?

From the preceding discussion of the integrin alpha V beta 3, it should be clear that this integrin, through interactions with ECM, various intramembranous and intracellular molecules, modulates the growth, survival, motility, and differentiation of a limited group of cells. These cells include osteoclasts,

activated macrophages, and angiogenic endothelial cells. Cellular functional activities are regulated by changes in the affinity and avidity of the integrin, through changes in its phosphorylation state, the nature of ECM ligands, and other factors.^{34–36}

The function of the alpha V and beta 3 subunits has also been assessed through studies of gene knockout mice. About 20% of alpha V integrin knockout mice survive to delivery. Placentas from these mice show defects, and the mice have abnormalities in central nervous system and gastrointestinal blood vessels. Cleft palate is also a frequent abnormality. The alpha V integrin subunit associates with beta 1, beta 3, beta 5, beta 6 and beta 8 integrin subunits.³⁷ Beta 3 integrin knockout mice are fertile and survive.³⁸ Their important problems are defective platelet aggregation, defective osteoclast resorptive capacity, and osteosclerosis.^{39–40} These also have to propensity to permit transplanted tumours to grow more rapidly.^{37–41} The report that transplanted tumours grow more rapidly in beta 3 knockout mice has been subject of extensive controversy because the conclusions contrast strikingly with studies with integrin antagonists on tumour growth.^{23–42} In other words, conclusions from gene knockout and studies with integrin antagonists on tumour growth have not been consistent with-out more in depth analysis.

The functional role of integrin alpha V beta 3 has been most extensively studied in the context of osteoclastogenesis and bone resorption, macrophage migration and activation, and angiogenesis. The integrin alpha V beta 3 is highly expressed on activated macrophages and osteoclasts.⁴³ These cell types are found in abundance at sites of bone destruction in RA patients—that is, activated macrophages are markedly increased in both subchondral bone and inflamed synovial tissues, and osteoclasts are markedly increased in subchondral bone at sites of bone erosion and resorption. In patients with RA, mature osteoclast numbers are highly associated with both periarticular and systemic bone loss.

Osteoclasts are highly specialised and differentiated multinucleated cells related to macrophages. Under normal physiological conditions, mature osteoclasts expressing alpha V beta 3 are continuously generated from immature bone marrow monocytic precursor cells. Stimulatory factors such as macrophage colony stimulating factor (M-CSF) and receptor activator of NF- κ B (RANK) ligand (RANKL) largely determine this differentiation process. During inflammatory diseases such as RA, tumour necrosis factor α (TNF α) and interleukin 1 (IL1) also significantly amplify osteoclastogenesis and generation of activated macrophages.^{43–44} Studies of macrophages and osteoclasts have shown that blocking alpha V beta 3 inhibits adhesion, migration, and, for osteoclasts, bone resorption. Resorbing osteoclasts develop a specialised membrane domain termed the tight sealing zone or clear zone. Alpha V beta 3 and one of its ligands, osteopontin, are enriched in the clear zone of the resorbing osteoclasts. The clear zone is believed to mediate attachment to the bone matrix and the formation of acidic resorption lacunae that are required for bone resorption.^{45–46} Other data suggest that alpha V beta 3 is involved in activated macrophage and osteoclast podosome formation. Podosomes are involved in transducing signals that regulate cell attachment, survival, and function internally into the macrophage and osteoclast.²² Disrupting alpha V beta 3 signalling molecules such as gelsolin impairs podosome formation, cell movements and bone resorption.⁴⁷ As osteoclast mediated bone resorption and macrophage dependent inflammation are such a central pathogenic feature of RA, these data provide a strong support for the concept that therapeutic inhibition of alpha V beta 3 is sensible.

Some of highest relative expression levels of alpha V beta 3 are observed on growth factor/cytokine activated endothelial cells, especially blood vessels in granulating wounds or tumours.⁷ It is also intensely expressed in inflamed synovial tissues of patients with RA and rabbits with experimental

RA-like arthritis.^{48–50} Alpha V beta 3 ligands, such as osteopontin and fibrinogen, are also abundant in rheumatoid synovial tissues. The interaction of these ligands with alpha V beta 3 seems to play an important part in development of the hyperplastic, tumour-like, invasive synovitis that contributes to destruction of bone and cartilage in RA.⁵¹ Endothelial cells in rheumatoid synovium are subject to continuous production of angiogenic stimuli (TNF α , VEGF, FGF1, FGF2), resulting in the expression of alpha V beta 3 on sprouting endothelial cell buds and new blood vessel development.²⁷ Alpha V beta 3 plays a very important part in this process by facilitating the attachment to and migration through the ECM by the newly developing blood vessels. These observations also support the view that inhibition of alpha V beta 3 in the synovium of RA patients may have therapeutic benefits.

BLOCKADE OF ALPHA V BETA 3 IN PRECLINICAL RA MODELS

A persuasive body of evidence indicates that alpha V beta 3 plays an important part in mediating the migration, differentiation, proliferation, and survival of a limited group of cells that express this integrin. It is particularly noteworthy that the cells that express the highest levels of alpha V beta 3 include activated macrophages, which are involved in producing proinflammatory cytokines, osteoclasts, which mediate inflammatory osteolysis, and endothelial cells, which are involved in pathological neovascularisation. Macrophage dependent processes, inflammatory osteolysis, and neovascularisation are clearly involved in the pathobiology of RA. These observations have stimulated the notion that blocking the function of integrin alpha V beta 3 may have therapeutic benefit in RA, and this hypothesis has been explored to a limited extent in preclinical animal models of inflammatory arthritis.

Intraarticular administration of a cyclic peptide alpha V beta 3 antagonist to rabbits with antigen induced arthritis, a model with features that resemble RA, inhibits synovial angiogenesis, inflammatory cell infiltration, and bone and cartilage destruction. The antagonist treatment also has a disease inhibitory effect when given to animals with chronic disease.⁵⁰ In addition, SB273005, a non-peptide antagonist of alpha V beta 3, was reported to significantly reduce symptoms and signs of adjuvant arthritis in the LEW rat. This arthritis model has a very good record of predicting therapeutic efficacy in patients with RA. SB273005 is also a well documented inhibitor of endothelial cell migration in vitro and a potent inhibitor in several animal models of bone resorption in vivo.⁵² Oral dosing with SB273005, started before disease onset or at the time of disease onset, significantly inhibits joint swelling and the destruction of both bone and cartilage (36%–52%, depending on schedule and dose).⁵³

More recently, the role of osteopontin has been examined in an experimental RA model. Osteopontin is an extracellular matrix protein containing Arg-Gly-Asp (RGD) sequence, which interacts with alpha V beta 3 integrins. Its expression in the joints of a murine RA model was associated with joint swelling, destruction of the surface structures of the joint based on scanning electron microscopy, and loss of toluidine blue positive proteoglycan content in the articular cartilage in wild type mice. In contrast, osteopontin deficiency prevented the mice from such surface destruction, loss of proteoglycan in the articular joint cartilage, and swelling of the joints. These preclinical results provide further support to the evolving view that alpha V beta 3 is an appropriate target for treatment of RA and related diseases.⁵⁴

Author's affiliations

R L Wilder, Clinical Development, Medimmune, Inc, 35 W Watkins Mill Road, Gaithersburg, MD 20878, USA

Correspondence to: Dr R L Wilder; wilderr@medimmune.com

REFERENCES

- 1 Wu H, Beuerlein G, Nie Y, Smith H, Lee BA, Hensler M, et al. Stepwise in vitro affinity maturation of Vitaxin, an alphav beta3- specific humanized mAb. *Proc Natl Acad Sci USA* 1998;95:6037–42.
- 2 Hynes RO. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 1992;69:11–25.
- 3 Rupp PA, Little CD. Integrins in vascular development. *Circ Res* 2001;89:566–72.
- 4 Brooks PC, Clark RA, Cheresh DA. Requirement of vascular integrin alpha v beta 3 for angiogenesis. *Science* 1994;264:569–71.
- 5 Teitelbaum SL. Osteoclasts, integrins, and osteoporosis. *J Bone Miner Metab* 2000;18:344–9.
- 6 Horton MA. The alpha v beta 3 integrin "vitronectin receptor". *Int J Biochem Cell Biol* 1997;29:721–5.
- 7 Eliceiri BP, Cheresh DA. The role of alphav integrins during angiogenesis: insights into potential mechanisms of action and clinical development. *J Clin Invest* 1999;103:1227–30.
- 8 Clover J, Dodds RA, Gowen M. Integrin subunit expression by human osteoblasts and osteoclasts in situ and in culture. *J Cell Sci* 1992;103 (Pt 1):267–71.
- 9 Helfrich MH, Nesbitt SA, Horton MA. Integrins on rat osteoclasts: characterization of two monoclonal antibodies (F4 and F11) to rat beta 3. *J Bone Miner Res* 1992;7:345–51.
- 10 Nesbitt S, Nesbitt A, Helfrich M, Horton M. Biochemical characterization of human osteoclast integrins. Osteoclasts express alpha v beta 3, alpha 2 beta 1, and alpha v beta 1 integrins. *J Biol Chem* 1993;268:16737–45.
- 11 Shinar DM, Schmidt A, Halperin D, Rodan GA, Weinreb M. Expression of alpha v and beta 3 integrin subunits in rat osteoclasts in situ. *J Bone Miner Res* 1993;8:403–14.
- 12 Felding-Habermann B, O'Toole TE, Smith JW, Fransvea E, Ruggeri ZM, Ginsberg MH, et al. Integrin activation controls metastasis in human breast cancer. *Proc Natl Acad Sci USA* 2001;98:1853–8.
- 13 Xiong JP, Stehle T, Diefenbach B, Zhang R, Dunker R, Scott DL, et al. Crystal structure of the extracellular segment of integrin {alpha}V{beta}3. *Science* 2001;294:339–45.
- 14 Beglova N, Blacklow SC, Takagi J, Springer T. Cysteine-rich module structure reveals a fulcrum for integrin rearrangement upon activation. *Nat Struct Biol* 2002;9:282–7.
- 15 Xiong JP, Stehle T, Zhang R, Joachimiak A, Frech M, Goodman SL, et al. Crystal structure of the extracellular segment of integrin {alpha}V{beta}3 in complex with an arg-gly-asp ligand. *Science* 2002;296:151–5.
- 16 Tsou R, Isik FF. Integrin activation is required for VEGF and FGF receptor protein presence on human microvascular endothelial cells. *Mol Cell Biochem* 2001;224:81–9.
- 17 Soldi R, Mitola S, Strasyly M, Defilippi P, Tarone G, Bussolino F. Role of alphavbeta3 integrin in the activation of vascular endothelial growth factor receptor-2. *EMBO J* 1999; 18:882–92.
- 18 Weerasinghe D, McHugh KP, Ross FP, Brown EJ, Gislis RH, Imhof BA. A role for the alphavbeta3 integrin in the transmigration of monocytes. *J Cell Biol* 1998;142:595–607.
- 19 Deryugina EI, Bourdon MA, Jungwirth K, Smith JW, Strongin AY. Functional activation of integrin alpha V beta 3 in tumor cells expressing membrane-type 1 matrix metalloproteinase. *Int J Cancer* 2000;86:15–23.
- 20 Deryugina EI, Ratnikov B, Monosov E, Postnova TI, DiScipio R, Smith JW, et al. MT1-MMP initiates activation of pro-MMP-2 and integrin alphavbeta3 promotes maturation of MMP-2 in breast carcinoma cells. *Exp Cell Res* 2001;263:209–23.
- 21 Hermant P, Armand M, Brown E, Rubio M, Ishihara H, Ulrich D, et al. The vitronectin receptor and its associated CD47 molecule mediates proinflammatory cytokine synthesis in human monocytes by interaction with soluble CD23. *J Cell Biol* 1999;144:767–75.
- 22 Pfaff M, Jurdic P. Podosomes in osteoclast-like cells: structural analysis and cooperative roles of paxillin, proline-rich tyrosine kinase 2 (Pyk2) and integrin alphavbeta3. *J Cell Sci* 2001;114 (pt 15):2775–86.
- 23 Stupack DG, Puente XS, Boutsabouloy S, Storgard CM, Cheresh DA. Apoptosis of adherent cells by recruitment of caspase-8 to unligated integrins. *J Cell Biol* 2001;155:459–70.
- 24 Eliceiri BP, Cheresh DA. The role of alphav integrins during angiogenesis. *Mol Med* 1998;4:741–50.
- 25 Jasin HE. Mechanisms of tissue damage in rheumatoid arthritis. In: Koopman WJ, ed. *Arthritis and allied conditions. A textbook of rheumatology*. Philadelphia: Lippincott Williams and Wilkins, 2001:1128–52.
- 26 Pap T, Gay R, Gay S. Mechanisms of joint destruction. In: Firestein GS, Panayi GS, Wollheim FA, eds. *Rheumatoid arthritis. New frontiers in pathogenesis and treatment*. Oxford: Oxford University Press, 2000:189–99.
- 27 Koch AE. The role of angiogenesis in rheumatoid arthritis: recent developments. *Ann Rheum Dis* 2000;59 (suppl 1):i65–71.
- 28 Goldring SR, Gravalles EM. Pathogenesis of bone erosions in rheumatoid arthritis. *Curr Opin Rheumatol* 2000;12:195–9.
- 29 Gravalles EM, Goldring SR. Cellular mechanisms and the role of cytokines in bone erosions in rheumatoid arthritis. *Arthritis Rheum* 2000;43:2143–51.
- 30 Albelda SM, Mette SA, Elder DE, Stewart R, Damjanovich L, Herlyn M, et al. Integrin distribution in malignant melanoma: association of the beta 3 subunit with tumor progression. *Cancer Res* 1990;50:6757–64.
- 31 Brooks PC, Montgomery AM, Rosenfeld M, Reisfeld RA, Hu T, Klier G, et al. Integrin alpha v beta 3 antagonists promote tumor regression by inducing apoptosis of angiogenic blood vessels. *Cell* 1994;79:1157–64.

- 32 **Brooks PC**, Stromblad S, Klemke R, Visscher D, Sarkar FH, Cheresh DA. Antiintegrin alpha v beta 3 blocks human breast cancer growth and angiogenesis in human skin. *J Clin Invest* 1995;96:1815–22.
- 33 **Carron CP**, Meyer DM, Pegg JA, Engleman VW, Nickols MA, Settle SL, *et al.* A peptidomimetic antagonist of the integrin alpha(v)beta3 inhibits Leydig cell tumor growth and the development of hypercalcemia of malignancy. *Cancer Res* 1998;58:1930–5.
- 34 **Boettiger D**, Huber F, Lynch L, Blystone S. Activation of alpha(v)beta3-vitronectin binding is a multistage process in which increases in bond strength are dependent on Y747 and Y759 in the cytoplasmic domain of beta3. *Mol Biol Cell* 2001;12:1227–37.
- 35 **Boettiger D**, Lynch L, Blystone S, Huber F. Distinct ligand-binding modes for integrin avb3-mediated adhesion to fibronectin vs vitronectin. *J Biol Chem* 2001;276:31684–90.
- 36 **Gottschalk KE**, Gunther R, Kessler H. A three-state mechanism of integrin activation and signal transduction for Integrin alphavbeta3. *Chembiochem* 2002;3:470–3.
- 37 **Carmeliet P**. Integrin indecision. *Nat Med* 2002;8:14–16.
- 38 **Bouvard D**, Brakebusch C, Gustafsson E, Aszodi A, Bengtsson T, Berna A, *et al.* Functional consequences of integrin gene mutations in mice. *Circ Res* 2001;89:211–23.
- 39 **Hodivala-Dilke KM**, McHugh KP, Tsakiris DA, Rayburn H, Crowley D, Ullman-Cullere M, *et al.* Beta3-integrin-deficient mice are a model for Glanzmann thrombasthenia showing placental defects and reduced survival. *J Clin Invest* 1999;103:229–38.
- 40 **McHugh KP**, Hodivala-Dilke K, Zheng MH, Namba N, Lam J, Novack D, *et al.* Mice lacking beta3 integrins are osteosclerotic because of dysfunctional osteoclasts. *J Clin Invest* 2000;105:433–40.
- 41 **Reynolds LE**, Wyder L, Lively JC, Taverna D, Robinson SD, Huang X, *et al.* Enhanced pathological angiogenesis in mice lacking beta3 integrin or beta5 integrins. *Nat Med* 2002;8:27–34.
- 42 **Cheresh DA**, Stupack DG. Integrin-mediated death: an explanation of the integrin-knockout phenotype? *Nat Med* 2002;8:193–4.
- 43 **Teitelbaum SL**. Bone resorption by osteoclasts. *Science* 2000;289:1504–8.
- 44 **Inoue M**, Ross FP, Erdmann JM, Abu-Amer Y, Wei S, Teitelbaum SL. Tumor necrosis factor alpha regulates alpha(v)beta5 integrin expression by osteoclast precursors in vitro and in vivo. *Endocrinology* 2000;141:284–90.
- 45 **Reinholt FP**, Hulthén K, Oldberg A, Heinegård D. Osteopontin—a possible anchor of osteoclasts to bone. *Proc Natl Acad Sci USA* 1990;87:4473–5.
- 46 **Nakamura I**, Pilkington MF, Lakkakorpi PT, Lipfert L, Sims SM, Dixon SJ, *et al.* Role of alpha(v)beta(3) integrin in osteoclast migration and formation of the sealing zone. *J Cell Sci* 1999;112 (pt 22):3985–93.
- 47 **Chellaiyah M**, Kizer N, Silva M, Alvarez U, Kwiatkowski D, Hruska KA. Gelsolin deficiency blocks podosome assembly and produces increased bone mass and strength. *J Cell Biol* 2000;148:665–78.
- 48 **Johnson BA**, Haines GK, Harlow LA, Koch AE. Adhesion molecule expression in human synovial tissue. *Arthritis Rheum* 1993;36:137–46.
- 49 **Walsh DA**, Wade M, Mapp PI, Blake DR. Focally regulated endothelial proliferation and cell death in human synovium. *Am J Pathol* 1998;52:691–702.
- 50 **Storgard CM**, Stupack DG, Jonczyk A, Goodman SL, Fox RI, Cheresh DA. Decreased angiogenesis and arthritic disease in rabbits treated with an alphavbeta3 antagonist. *J Clin Invest* 1999;103:47–54.
- 51 **Petrov PK**, Hummel KM, Schedel J, Franz JK, Klein CL, Muller-Ladner U, *et al.* Expression of osteopontin messenger RNA and protein in rheumatoid arthritis: effects of osteopontin on the release of collagenase 1 from articular chondrocytes and synovial fibroblasts. *Arthritis Rheum* 2000;43:1597–605.
- 52 **Miller WH**, Keenan RM, Willette RN, Lark MW. Identification and in vivo efficacy of small-molecule antagonists of integrin alphavbeta3 (the vitronectin receptor). *Drug Discov Today* 2000;5:397–408.
- 53 **Badger AM**, Blake S, Kapadia R, Sarkar S, Levin J, Swift BA, *et al.* Disease-modifying activity of SB 273005, an orally active, nonpeptide alphavbeta3 (vitronectin receptor) antagonist, in rat adjuvant-induced arthritis. *Arthritis Rheum* 2001;44:128–37.
- 54 **Yumoto K**, Ishijima M, Rittling SR, Tsuji K, Tsuchiya Y, Kon S, *et al.* Osteopontin deficiency protects joints against destruction in anti-type II collagen antibody-induced arthritis in mice. *Proc Natl Acad Sci USA* 2002;99:4556–61.