



OPEN ACCESS

# Clinical relevance of HEp-2 indirect immunofluorescent patterns: the International Consensus on ANA patterns (ICAP) perspective

Jan Damoiseaux,<sup>1</sup> Luis Eduardo Coelho Andrade,<sup>2</sup> Orlando Gabriel Carballo,<sup>3,4</sup> Karsten Conrad,<sup>5</sup> Paulo Luiz Carvalho Francescantonio,<sup>6</sup> Marvin J Fritzler,<sup>7</sup> Ignacio Garcia de la Torre,<sup>8</sup> Manfred Herold,<sup>9</sup> Werner Klotz,<sup>10</sup> Wilson de Melo Cruvinel,<sup>11</sup> Tsuneyo Mimori,<sup>12</sup> Carlos von Muhlen,<sup>13</sup> Minoru Satoh,<sup>14</sup> Edward K Chan<sup>15</sup>

**Handling editor** Josef S Smolen

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2018-214436>).

For numbered affiliations see end of article.

## Correspondence to

Dr Jan Damoiseaux, Central Diagnostic Laboratory, Maastricht University Medical Center, Maastricht 6229 HX, The Netherlands; [jan.damoiseaux@mumc.nl](mailto:jan.damoiseaux@mumc.nl)

Received 13 September 2018  
Accepted 23 January 2019  
Published Online First 12 March 2019

## ABSTRACT

The indirect immunofluorescence assay (IIFA) on HEp-2 cells is widely used for detection of antinuclear antibodies (ANA). The dichotomous outcome, negative or positive, is integrated in diagnostic and classification criteria for several systemic autoimmune diseases. However, the HEp-2 IIFA test has much more to offer: besides the titre or fluorescence intensity, it also provides fluorescence pattern(s). The latter include the nucleus and the cytoplasm of interphase cells as well as patterns associated with mitotic cells. The International Consensus on ANA Patterns (ICAP) initiative has previously reached consensus on the nomenclature and definitions of HEp-2 IIFA patterns. In the current paper, the ICAP consensus is presented on the clinical relevance of the 29 distinct HEp-2 IIFA patterns. This clinical relevance is primarily defined within the context of the suspected disease and includes recommendations for follow-up testing. The discussion includes how this information may benefit the clinicians in daily practice and how the knowledge can be used to further improve diagnostic and classification criteria.

## INTRODUCTION

Autoantibodies, as detected by the indirect immunofluorescence assay (IIFA) on HEp-2 cells (IIFA HEp-2), are recognised as important diagnostic markers in a plethora of autoimmune diseases, in particular the systemic autoimmune rheumatic diseases (SARD).<sup>1</sup> Although somewhat dated by today's standards, members of the American College of Rheumatology (ACR) prepared an evidence-based guideline for the usefulness of the HEp-2 IIFA results for diagnostic and prognostic purposes and also for meeting diagnostic criteria.<sup>2</sup> That guideline was based on reactivity with nuclear antigens as detected by IIFA on rodent tissue or HEp-2 cells. More recently, the IIFA on HEp-2 cells was reinforced as the gold standard for autoantibody screening in SARD.<sup>3</sup>

Interestingly, the HEp-2 IIFA test reveals much more information than the mere absence or presence of autoantibodies, that is, the level of antibody as well as the HEp-2 IIFA pattern. Based on titration or appropriate evaluation of the fluorescence intensity, the antibody level can be determined and this information has general concordance with the

clinical relevance of the test result. Indeed, higher antibody levels are better associated with SARD and have an increased likelihood to identify the auto-antigen in follow-up testing.<sup>4–6</sup> The importance of the level of autoantibodies is also recognised in the ACR guideline as well as by the recommendations issued by the European Autoimmunity Standardization Initiative (EASI) and the International Union of Immunologic Societies (IUIS) Autoantibody Standardization Subcommittee.<sup>2,7</sup>

The HEp-2 IIFA pattern may also reveal clinically relevant information. This information is not restricted to giving direction to follow-up testing for antigen-specificity, but, for instance, the centromere pattern is included in the classification criteria for systemic sclerosis,<sup>8</sup> while the nuclear dense fine speckled pattern is reported to be more prevalent in apparently healthy individuals as compared with patients with SARD.<sup>9</sup> To harmonise the names and descriptions of the distinct HEp-2 IIFA patterns, an ordered classification taxonomy was proposed.<sup>10</sup> This proposal was subsequently elaborated on by the International Consensus on ANA Patterns (ICAP), initiated in parallel to the 12th International Workshop on Autoantibodies and Autoimmunity (2014) held in Sao Paulo, Brazil. During this workshop, a consensus was reached on the nomenclature and definitions of 28 HEp-2 IIFA patterns. Each HEp-2 IIFA pattern was ascribed an alphanumeric code from AC-1 to AC-28.<sup>11</sup> The consensus nomenclature for each pattern and representative images were also made available online at the ICAP website (<http://www.ANAPatterns.org>).

In addition to the nuclear patterns, important cytoplasmic and mitotic patterns may also be observed in HEp-2 IIFA analysis. Although reporting non-nuclear patterns is considered clinically relevant,<sup>7</sup> for various jurisdictional reasons there is no clear-cut consensus viewpoint on reporting non-nuclear patterns as a negative or positive test.<sup>12</sup> With the understanding that the term 'Antinuclear antibody (ANA) test' may be inappropriate to designate a test that also addresses autoantibodies to antigens in the cytoplasm and mitotic apparatus, an alternative name, anticellular antibodies, was suggested in the EASI/IUIS recommendations.<sup>7</sup> Recent publications from ICAP have preferred the term HEp-2 IIFA as it covers the



© Author(s) (or their employer(s)) 2019. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ.

**To cite:** Damoiseaux J, Andrade LEC, Carballo OG, et al. *Ann Rheum Dis* 2019;**78**:879–889.

## Recommendation

whole spectrum of patterns that can be observed when using the HEp-2 cells as substrate.<sup>13 14</sup>

Originally, the HEp-2 IIFA patterns were associated with diseases, but it was anticipated that many of these associations are only valid if the antigen-specificity was confirmed by follow-up testing. In subsequent ICAP workshops, it was agreed that the disease associations should be replaced by clinical relevance. In this current paper, we present the consensus on the clinical relevance of the distinct HEp-2 IIFA patterns as achieved by consecutive workshops and discussions among the executive ICAP members.

## MATERIALS AND METHODS

For discussion about the structure of clinical relevance templates were prepared for AC-2 (LECA), AC-3 (JD) and AC-5 (MS). This formed the basis of a guideline for description of each AC pattern (EC). Of highest importance, it was agreed that the information should be objective and helpful for the clinician, the pattern-antigen associations should be put in the right clinical context and information should be evidence-based.

In preparation for the third ICAP workshop in Kyoto (2016), composition of the clinical relevance documents was started for the nuclear patterns (JD, LECA, MS), cytoplasmic patterns (CAvM, EKLC) and mitotic patterns (MH, TM). As far as already available, the documents were commented on by the ICAP executive board and, after appropriate adjustment, discussed with the workshop participants. The feedback from participants mainly focused on the structure of the information provided, on the required level of detail and the format of recommended follow-up testing.

In anticipation of the fourth ICAP workshop in Dresden (2017), the set of clinical relevance documents was completed for all patterns. Further comments from the ICAP executive board were included. The resulting documents were individually discussed with the workshop participants for nuclear (JD), cytoplasmic (CAvM) and mitotic (MH) patterns. Besides several substantive comments, there was general agreement that the information should be provided in tabular format at two distinct levels. The first level should contain information on relevant follow-up testing in the respective clinical context, the recommended follow-up tests should be commercially available and detailed test characteristics should not be given because of potential geographic and jurisdictional differences. Information based on case reports or small patient cohorts, as well as information on possible follow-up testing that is only available in specialised research laboratories, should only be provided in the second level information.

Tables for nuclear, cytoplasmic and mitotic patterns were prepared for first and second level information (JD). These tables were commented by the ICAP executive board and finalised by JD. Of note, since the starting point of the tables on clinical relevance is the HEp-2 IIFA pattern and not the clinically suspected disease, the tables do not list all autoantibodies related to the respective disease.

## RESULTS

### Nuclear HEp-2 IIFA patterns

To date, a total of 15 nuclear HEp-2 IIFA patterns have been described, that is, AC-1–AC-14 and AC-29. Table 1 summarises the clinical relevance of these patterns.<sup>8 9 14–79</sup> Since AC-29 was only recently described,<sup>14</sup> the advice for follow-up testing for autoantibodies to topoisomerase I (Scl-70) in case of clinical suspicion of systemic sclerosis is also added as a note to the

clinical relevance of AC-1. In particular, disease-specific immunoassays, like autoimmune liver disease profile, inflammatory myopathy profile, systemic sclerosis profile, are often only available in specialty clinical laboratories.

For six nuclear HEp-2 IIFA patterns (AC-3, 5, 7, 8, 12 and 13), additional information about clinical relevance is summarised in online supplementary table S1. Although some assays for anti-CENP-A antibodies are commercially available, these antibodies are included in online supplementary table S1 because the majority of sera revealing the AC-3 pattern are also reactive with CENP-B. In contrast to CENP-A, CENP-B is included in many routine extractable nuclear antigens profiles.

### Cytoplasmic HEp-2 IIFA patterns

Table 2 summarises the clinical relevance of the nine cytoplasmic HEp-2 IIFA patterns, that is, AC-15–AC-23.<sup>26 33 80–101</sup> It is recognised that the distinction between AC-19 (dense fine speckled) and AC-20 (fine speckled) can be challenging. Moreover, within the spectrum of anti-tRNA synthetase antibodies, not all produce an HEp-2 IIFA pattern and only some anti-Jo-1 antibodies are considered to give the AC-20 pattern, while the other anti-tRNA synthetase antibodies (EJ, KS, OJ, PL-7 and PL-12) are more likely to reveal the AC-19 pattern. Solid information on the pattern of two additional anti-tRNA synthetase antibodies (Ha and Zo) is lacking. Overall, the relation between these two cytoplasmic HEp-2 IIFA patterns and the distinct anti-tRNA synthetase antibodies is subject to further discussion. In clinical practice, the complete spectrum of the anti-tRNA synthetase antibodies should be determined irrespective of the subtype of cytoplasmic speckled pattern, that is, AC-19 or AC-20.

For seven cytoplasmic HEp-2 IIFA patterns (AC-15–AC-19, AC-22 and AC-23), more detailed information is provided in online supplementary table S2. In particular, for AC-16–AC-18, the clinical associations are quite diverse, depending on the antigen recognised. Overall, the clinical associations provided are primarily based on antigen-specific immunoassays and not on the HEp-2 IIFA pattern as such.

### Mitotic HEp-2 IIFA patterns

The clinical relevance of the five mitotic patterns is summarised in table 3,<sup>102–122</sup> with more detailed information in online supplementary table S3. As for the cytoplasmic patterns, clinical associations for the mitotic patterns are primarily based on antigen-specific immunoassays and not on the HEp-2 IIFA pattern as such.

## DISCUSSION

In the current paper, we present the ICAP consensus on the clinical relevance of 29 HEp-2 IIFA patterns defined by ICAP.<sup>11 14</sup> The consensus on clinical relevance is defined in the clinical context of the patient, that is, suspected disease, and includes recommended follow-up testing within the spectrum of antigen-specificities that are commercially available. Obviously, if follow-up testing identifies the antigen, the clinical relevance can be further refined.<sup>123</sup>

Defining the clinical relevance of HEp-2 IIFA patterns in the context of disease manifestations is meant to be an important tool for the clinician in the diagnostic work-up of patients suspected of SARD. Unfortunately, good data on the association between HEp-2 IIFA patterns and the distinct diseases are lacking, probably due to reasons summarised below. There are several reasons for not finding a perfect association between HEp-2 IIFA patterns and diseases. First, pattern assignment in

**Table 1** Nuclear HEp-2 IIFA patterns

Code	AC pattern—clinical relevance	Refs
AC-1	<p><b>HOMOGENEOUS</b></p> <ul style="list-style-type: none"> <li>Found in patients with SLE, chronic autoimmune hepatitis or juvenile idiopathic arthritis</li> <li>If SLE is clinically suspected, it is recommended to perform a follow-up test for anti-dsDNA antibodies, alone or in combination with dsDNA/histone complexes (nucleosomes/chromatin); anti-dsDNA antibodies are included in the classification criteria for SLE</li> <li>If chronic autoimmune hepatitis or juvenile idiopathic arthritis is suspected, follow-up testing is not recommended because the respective autoantigens revealing the AC-1 pattern are not completely defined</li> </ul> <p>Notes: Although autoantibodies to Topoisomerase I (formerly Scl-70) may be reported as nuclear homogeneous, they typically reveal a composite AC-29 HEp-2 IIFA pattern; as such, clinical suspicion of SSc may warrant follow-up testing for reactivity to this antigen.</p> <p>Although AC-1 is the most prevalent pattern in chronic autoimmune hepatitis, other HEp-2 IIFA patterns may occur, but also for these patterns the autoantigens are not completely defined.</p>	<p>15, 16</p> <p>17</p> <p>14, 18</p> <p>19</p>
AC-2	<p><b>DENSE FINE SPECKLED</b></p> <ul style="list-style-type: none"> <li>Commonly found as high titer HEp-2 IIFA-positive in apparently healthy individuals or in patients who do not have a systemic autoimmune rheumatic disease (SARD)</li> <li>The negative association with SARD is only valid if the autoreactivity is confirmed as being directed to DFS70 (also known as LEDGF/p75) and if no other common ENA is recognized</li> <li>Both in apparently healthy individuals as well as patients who do not have a SARD the AC-2 pattern may be caused by autoantibodies to other antigens than DFS70</li> </ul> <p>Note: Confirmatory assays for anti-DFS70 antibodies may be available only in specialty clinical laboratories.</p>	<p>9</p> <p>20, 21</p> <p>22</p>
AC-3	<p><b>CENTROMERE</b> (see online supplementary table S1 for further details)</p> <ul style="list-style-type: none"> <li>Commonly found in patients with limited cutaneous SSc, and as such included in the classification criteria for SSc</li> <li>In combination with Raynaud phenomenon, the AC-3 pattern is prognostic for onset of limited cutaneous SSc</li> <li>Strongly associated with antibodies to CENP-B; especially in case of low titers, confirmation by an antigen-specific immunoassay is recommended to support the association with limited cutaneous SSc; the CENP-B antigen is included in many routine ENA profiles</li> <li>The AC-3 pattern is also apparent in a subset of patients with PBC; these patients often have both SSc as well as PBC</li> </ul>	<p>8, 15, 23</p> <p>15, 23</p> <p>15</p> <p>15</p>
AC-4	<p><b>FINE SPECKLED</b></p> <ul style="list-style-type: none"> <li>Present to a varying degree in distinct SARD, in particular SjS, SLE, subacute cutaneous lupus erythematosus, neonatal lupus erythematosus, congenital heart block, DM, SSc, and SSc-AIM overlap syndrome</li> <li>If SjS, SLE, subacute cutaneous lupus erythematosus, neonatal lupus erythematosus, or congenital heart block is clinically suspected, it is recommended to perform follow-up tests for anti-SS-A/Ro (Ro60) and anti-SS-B/La antibodies; in most laboratories these antigens are included in the routine ENA profile</li> <li>Autoantibodies to SS-A/Ro are part of the classification criteria for SjS (the criteria do not distinguish between Ro60 and Ro52/TRIM21)</li> <li>If SSc, AIM, or to a lesser extent SLE, is clinically suspected, it is recommended to perform follow-up tests for detecting autoantibodies to Mi-2, TIF1γ, and Ku; these antigens are typically included in disease specific immunoassays (i.e., inflammatory myopathy profile*)</li> <li>Autoantibodies to Mi-2 and TIF1γ are associated with DM; autoantibodies to TIF1γ in patients with DM, although rare in the overall AC-4 pattern, is strongly associated with malignancy in old patients</li> <li>Autoantibodies to Ku are associated with SSc-AIM and SLE-SSc-AIM overlap syndromes</li> </ul> <p>Notes: Anti-SS-A/Ro (Ro60) and AIM-specific autoantibodies may be undetected in HEp-2 IIFA-screening.</p>	<p>15</p> <p>15</p> <p>25</p> <p>26</p> <p>26, 27</p> <p>26</p> <p>28</p>
AC-5	<p><b>LARGE/COARSE SPECKLED</b> (see online supplementary table S1 for further details)</p> <ul style="list-style-type: none"> <li>Present to a varying degree in distinct SARD, in particular SLE, SSc, MCTD, SSc-AIM overlap syndrome, and UCTD (i.e, patients with rheumatic symptoms without a definite SARD diagnosis)</li> <li>If SLE is clinically suspected, it is recommended to perform follow-up tests for anti-Sm and anti-U1RNP antibodies; these antigens are commonly included in the routine ENA profile; anti-Sm antibodies are included in the classification criteria for SLE</li> <li>If SSc is clinically suspected, it is recommended to perform a follow-up test for anti-RNApol III antibodies (e.g., SSc profile*); the anti-RNApol III antibodies are included in the classification criteria for SSc</li> <li>If MCTD is clinically suspected, it is recommended to perform a follow-up test for anti-U1RNP antibodies; the antigen is commonly included in the routine ENA profile; anti-U1RNP antibodies are included in the diagnostic criteria for MCTD</li> <li>If the SSc-AIM overlap syndrome is clinically suspected, it is recommended to perform follow-up tests for anti-U1RNP and anti-Ku antibodies; these antigens are included in the routine ENA profile (U1RNP), or in disease specific immunoassays (Ku, i.e., inflammatory myopathy profile* and SSc profile*)</li> <li>In non-SARD individuals in the general population, the presence of the AC-5 pattern is not associated with the autoantigens mentioned above and most often concerns low antibody titers</li> </ul>	<p>29</p> <p>16, 30, 31</p> <p>8</p> <p>32</p> <p>26, 33</p>

Continued

## Recommendation

Table 1 Continued

Code	AC pattern—clinical relevance	Refs
AC-6	<p><b>MULTIPLE NUCLEAR DOTS</b></p> <ul style="list-style-type: none"> <li>Found in a broad spectrum of autoimmune diseases, including PBC, AIM (DM), as well as other inflammatory conditions <a href="#">34</a></li> <li>If PBC is clinically suspected, it is recommended to perform follow-up tests for anti-Sp100 (and PML/Sp140) antibodies; in particular anti-Sp100 antibodies have the best clinical association with PBC and have added value, especially when associated with AMA; the Sp100 (and PML-Sp140) antigen is included in disease specific immunoassays (ie, liver profile*) <a href="#">17, 35, 36</a></li> <li>If DM is clinically suspected, it is recommended to perform a follow-up test for anti-MJ/NXP-2 antibodies; these anti-MJ/NXP-2 antibodies are highly specific for AIM, are found in up to one third of patients with juvenile DM, and have been reported to be associated with malignancies in adult AIM patients; the antigen is included in disease specific immunoassays (i.e., inflammatory myopathy profile*) <a href="#">37–39</a></li> </ul>	
AC-7	<p><b>FEW NUCLEAR DOTS</b> (see online supplementary table S1 for further details)</p> <ul style="list-style-type: none"> <li>The AC-7 pattern has low positive predictive value for any disease <a href="#">40, 41</a></li> <li>Antigens primarily localized in the dots include p80-coilin and SMN complex; specific immunoassays for these autoantibodies are currently not commercially available <a href="#">42, 43</a></li> </ul>	
AC-8	<p><b>HOMOGENEOUS NUCLEOLAR</b> (see online supplementary table S1 for further details)</p> <ul style="list-style-type: none"> <li>Found in patients with SSc, SSc-AIM overlap syndrome, and patients with clinical manifestations of other SARD <a href="#">44–46</a></li> <li>If limited cutaneous SSc is clinically suspected, it is recommended to perform a follow-up test for anti-Th/To antibodies; the antigen is included in disease specific immunoassays (ie, SSc profile*) <a href="#">44, 45</a></li> <li>If SSc-AIM overlap syndrome is clinically suspected, it is recommended to perform a follow-up test for anti-PM/Scl antibody reactivity; the antigen may be included in the routine ENA profile and is included in disease specific immunoassays (i.e., inflammatory myopathy profile* and the SSc profile*); in general, anti-PM/Scl antibodies yield a diffuse nuclear fine speckled staining in addition to the AC-8 pattern <a href="#">46</a></li> <li>Other antigens recognized include B23/nucleophosmin, No55/SC65, and C23/nucleolin, but the clinical significance of these autoantibodies is not well established; specific immunoassays for these autoantibodies are currently not commercially available</li> </ul> <p>Notes: Although some anti-Th/To antibody immunoassays are commercially available, technical issues relating to the limited sensitivity of these immunoassays should be taken in to consideration. <a href="#">44, 47</a></p>	
AC-9	<p><b>CLUMPY NUCLEOLAR</b></p> <ul style="list-style-type: none"> <li>Found in patients with SSc <a href="#">48</a></li> <li>If SSc is clinically suspected, it is recommended to perform a follow-up test for anti-U3RNP/fibrillarin antibodies; the antigen is included in disease specific immunoassays (i.e, SSc profile*) <a href="#">48</a></li> <li>If confirmed as anti-U3RNP/fibrillarin reactivity by immunoassay, the clinical association is with diffuse SSc, increased incidence of pulmonary arterial hypertension, skeletal muscle disease, severe cardiac involvement, and gastrointestinal dysmotility <a href="#">23, 48–50</a></li> <li>Among SSc patients, anti-U3RNP/fibrillarin antibodies are most commonly found in African American and Latin American patients <a href="#">48, 49, 51</a></li> </ul> <p>Notes: Although some anti-U3RNP/fibrillarin immunoassays are commercially available, technical issues relating to the limited sensitivity of these immunoassays should be taken into consideration. <a href="#">24</a></p>	
AC-10	<p><b>PUNCTATE NUCLEOLAR</b></p> <ul style="list-style-type: none"> <li>The AC-10 pattern can be seen in various conditions, including SSc, Raynaud's phenomenon, SjS, and cancer <a href="#">52–56</a></li> <li>If the AC-10 pattern is observed in the serum of patients with conditions mentioned above, follow-up testing for anti-NOR90(hUBF) antibodies is to be considered; the antigen is included in disease specific immunoassays (i.e. SSc profile*) <a href="#">54, 55</a></li> <li>While AC-10 is associated with anti-RNAPol I antibodies, these antibodies almost always coexist with anti-RNAPol III antibodies which reveal the AC-5 pattern; therefore, if SSc is clinically suspected, it is recommended to perform a follow-up test for anti-RNAPol III antibodies (See also AC-5); specific immunoassays for anti-RNAPol I antibodies are currently not commercially available <a href="#">52, 53, 57</a></li> </ul>	
AC-11	<p><b>SMOOTH NUCLEAR ENVELOPE</b></p> <ul style="list-style-type: none"> <li>The AC-11 pattern is infrequently found in routine autoantibody testing and has been described in autoimmune-cytopenias, autoimmune liver diseases, linear scleroderma, APS, and SARD; current information on clinical associations is based mainly on case reports and small cohorts <a href="#">58–60</a></li> <li>Antigens recognized include lamins (A, B, C) and LAP-2; specific immunoassays for these autoantibodies are currently not commercially available <a href="#">58–60</a></li> </ul>	
AC-12	<p><b>PUNCTATE NUCLEAR ENVELOPE</b> (see online supplementary table S1 for further details)</p> <ul style="list-style-type: none"> <li>Found in patients with PBC, as well as patients with other autoimmune liver diseases and SARD <a href="#">61</a></li> <li>If PBC is clinically suspected, it is recommended to perform a follow-up test for anti-gp210 antibodies; the antigen is included in disease specific immunoassays (ie, extended liver profile*) <a href="#">62–64</a></li> <li>Other antigens recognized include p62 nucleoporin, LBR, and Tpr; specific immunoassays for these autoantibodies are currently not commercially available <a href="#">65–68</a></li> </ul>	

Continued



Table 1 Continued

Code	AC pattern—clinical relevance	Refs
AC-13	PCNA-like (see online supplementary table S1 for further details) <ul style="list-style-type: none"><li>▶ The AC-13 pattern has formerly been considered highly specific for SLE, but this specificity is debated</li><li>▶ If SLE is clinically suspected, it is recommended to perform a follow-up test for anti-PCNA antibodies; the antigen is included in several routine ENA profiles</li><li>▶ Recent studies with antigen-specific immunoassays show clinical associations also with SSc, AIM, RA, HCV, and other conditions</li></ul>	69, 70 69 70–73
AC-14	CENP-F-like <ul style="list-style-type: none"><li>▶ The majority of sera exhibiting the AC-14 pattern are from patients with a diversity of neoplastic conditions (breast, lung, colon, lymphoma, ovary, brain); paradoxically, the frequency of the AC-14 pattern in patient cohorts with these malignancies is low</li><li>▶ The AC-14 pattern is also seen in inflammatory conditions (Crohn’s disease, autoimmune liver disease, SjS, graft-versus-host disease); current information on clinical associations is based mainly on case reports and series of cases</li><li>▶ Possible associations only hold if the reactivity to CENP-F is confirmed in an antigen-specific immunoassay; current information on clinical associations is based mainly on case reports and series of cases; specific immunoassays for this autoantibody are currently not commercially available</li></ul>	74–78
AC-29	TOPOI-like <ul style="list-style-type: none"><li>▶ The AC-29 pattern is highly specific for SSc, in particular with diffuse cutaneous SSc and more aggressive forms of SSc</li><li>▶ If SSc is clinically suspected, it is recommended to perform a follow-up test for anti-Topoisomerase I (formerly Scl-70) antibodies; the anti-Topoisomerase I antibodies are included in the classification criteria for SSc and the antigen is included in routine ENA profiles</li></ul>	14, 18, 23 8, 23, 79

\*Availability of the inflammatory myopathy profile, the SSc profile and the (extended) liver profile may be limited to specialty clinical laboratories. AIM, autoimmune myopathy; AMA, antimitochondrial antibodies; APS, antiphospholipid syndrome; CENP, centromere-associated protein; DFS, dense fine speckled; DM, dermatomyositis; ENA, extractable nuclear antigens; HCV, hepatitis C virus; IIFA, indirect immunofluorescence assay; LAP, lamin-associated polypeptide; LBR, lamin B receptor; LEDGF, lens epithelial derived growth factor; NOR, nucleolus organiser region; NXP, nuclear matrix protein; PBC, primary biliary cholangitis; PCNA, proliferating cell nuclear antigen; PML, promyelocytic leukaemia; PM/Scl, polymyositis-scleroderma; RA, rheumatoid arthritis; RNAPol, RNA polymerase; RNP, ribonucleoprotein; SARD, systemic autoimmune rheumatic diseases; SLE, systemic lupus erythematosus; SMN, survival of motor neuron; SSc, systemic sclerosis; SjS, Sjögren’s syndrome; TIF, transcription intermediary factor; TRIM, tripartite motif; Tpr, translocated promoter region; UCTD, undifferentiated connective tissue disease; dsDNA, double stranded DNA; hUBF, human upstream binding factor.

clinical laboratories is rather inconsistent as shown by external quality assessments.<sup>14 124 125</sup> This is exactly the reason why ICAP was initiated: the consensus on nomenclature and definitions of HEp-2 IIFA patterns allows to align pattern description across laboratories. Also, the integration of computer-aided immunofluorescence microscopy (CAIFM) may further improve the consistency in pattern assignments.<sup>126–131</sup> As such, it is promising that several companies involved in CAIFM have declared their intention to accommodate to the ICAP classification. Second, even apparently healthy individuals may have autoantibodies as detected by the HEp-2 IIFA. Such autoantibodies, being either innocent bystander antibodies or predictive antibodies, may still be present on development of SARD and interfere with the SARD-related pattern. Interestingly, the pattern best associated with apparently healthy individuals is the nuclear dense fine speckled pattern (AC-2), but this association only holds if the specificity is confirmed as monospecific for DFS70.<sup>20 21 132</sup> Third, the HEp-2 IIFA patterns may slightly differ depending on the cellular substrate used. For this reason, the ICAP website contains for each pattern multiple pictures taken from different brands of HEp-2 slides. Fourth, diseases like systemic lupus erythematosus and autoimmune inflammatory myopathies may be associated with distinct autoantibodies, each associated with a distinct HEp-2 IIFA pattern. If the autoantigens are ill defined, as is the case, for instance, in autoimmune hepatitis, only the most prevalent patterns are included. Altogether, it is evident that, with the exception of the centromere pattern (AC-3), all patterns are to be confirmed by antigen-specific immunoassay for a solid association with the respective autoimmune diseases. While consensus statements have been generated for all 29 HEp-2 IIFA patterns, and it is highly recommended to report

patterns,<sup>7 11</sup> it is anticipated that laboratories may restrict their reports to the so-called ‘competent level’ patterns (<http://www.ANAPatterns.org>).<sup>133</sup> Although, for instance, the nucleolar patterns may not be reported as distinct entities (AC-8, AC-9 and AC-10), all three subtypes represent autoantibodies reactive with antigens associated with systemic sclerosis, either alone or in combination with autoimmune inflammatory myopathies. Follow-up testing, therefore, anyhow involves the systemic sclerosis multiparameter assay including all the relevant autoantibodies. Traditionally, only nuclear HEp-2 IIFA patterns have been considered as a true positive HEp-2 IIFA test, and this is most likely related to the time-honoured terminology ‘Antinuclear Antibody Test’,<sup>12</sup> but it is evident from this report that even for nuclear HEp-2 IIFA patterns, the clinical associations are quite diverse. In particular, the nuclear dense fine speckled pattern (AC-2) seems to have an inverse association with SARD.<sup>9 134</sup> On the other hand, the cytoplasmic HEp-2 IIFA patterns, and to a lesser extent the mitotic patterns, are also clinically relevant and may demand dedicated follow-up testing in daily clinical practice. Therefore, the ICAP executive board advocates that information on HEp-2 IIFA patterns should be reported to the clinician and should also be incorporated in diagnostic and classification criteria instead of the simple assignment ‘ANA-positive’.<sup>135</sup> Although the HEp-2 IIFA has been considered the gold standard for autoantibody detection in SARD,<sup>3</sup> the limitations of this assay are understood.<sup>136–138</sup> Indeed, up to 35% of healthy controls may be positive if a screening dilution of 1/40 is used.<sup>139</sup> Therefore, in the EASI/IUIS recommendations, it is advocated that each laboratory verifies that the screening dilution is defined by a cut-off set at the 95th percentile.<sup>7</sup> However, by

## Recommendation

Table 2 Cytoplasmic HEp-2 IIFA patterns

Code	AC pattern—clinical relevance	Refs
AC-15	<p>FIBRILLAR LINEAR (see online supplementary table S2 for further details)</p> <ul style="list-style-type: none"> <li>► Found in patients with AIH type 1, chronic HCV infection, and celiac disease (IgA isotype); rare in SARD <a href="#">17</a></li> <li>► If AIH type 1 is clinically suspected, it is recommended to confirm reactivity with smooth muscle antibodies (IgG isotype), typically detected by IIFA on rodent tissue (liver, stomach, kidney); anti-smooth muscle antibodies are included in the international criteria for AIH type 1 <a href="#">17,80</a></li> <li>► F-actin is the main target antigen of anti-smooth muscle antibodies in AIH type 1; autoantibodies to F-actin are of more clinical importance than antibodies to G-actin <a href="#">81–83</a></li> </ul> <p>Notes: Although anti-F-actin immunoassays are commercially available, technical issues relating to the sensitivity of these immunoassays should be taken into consideration.</p>	
AC-16	<p>FIBRILLAR FILAMENTOUS (see online supplementary table S2 for further details)</p> <ul style="list-style-type: none"> <li>► Found in various diseases, but AC-16 is not typically found in SARD</li> <li>► Antigens recognized include cytokeratins 8, 18, &amp; 19, tubulin, and vimentin; specific immunoassays for these autoantibodies are currently not commercially available</li> </ul>	
AC-17	<p>FIBRILLAR SEGMENTAL (see online supplementary table S2 for further details)</p> <ul style="list-style-type: none"> <li>► Found very infrequently in a routine serology diagnostic setting</li> <li>► Antigens recognized include <math>\alpha</math>-Actinin and Vinculin; specific immunoassays for these autoantibodies are currently not commercially available</li> </ul>	
AC-18	<p>DISCRETE DOTS (see online supplementary table S2 for further details)</p> <ul style="list-style-type: none"> <li>► Autoantibodies revealing the AC-18 pattern have been reported in distinct SARD and in a variety of other diseases; their prevalence in unselected or specified disease cohorts has not been thoroughly studied <a href="#">84</a></li> <li>► Antigens recognized include GW-body (Processing or P body) antigens (Ge-1/Hedls, GW182, and Su/Ago2) and endosomal antigens (EEA1, CLIP-170, GRASP-1, and LBPA); specific immunoassays for these autoantibodies are currently not commercially available</li> </ul> <p>Notes: Autoantibodies to GW-bodies and endosomes may yield slightly different HEp-2 IIFA patterns. <a href="#">84, 85</a></p>	
AC-19	<p>DENSE FINE SPECKLED (see online supplementary table S2 for further details)</p> <ul style="list-style-type: none"> <li>► Found in patients with SLE and the anti-synthetase syndrome (a subset of AIM), interstitial lung disease, polyarthritis, Raynaud's phenomenon, and mechanic's hands; these features may occur in various combinations or as an isolated manifestation, especially interstitial lung disease <a href="#">33, 86, 87</a></li> <li>► If SLE is clinically suspected, follow-up tests for antibodies to ribosomal P phosphoproteins (P0, P1, P2, C22 RibP peptide) are recommended; these antigens may be included in the routine ENA profile</li> <li>► Anti-RibP antibodies have been associated in some studies with neuropsychiatric lupus, and in childhood-onset SLE with autoimmune hemolytic anemia <a href="#">86, 88, 89</a></li> <li>► If AIM, in particular the anti-synthetase syndrome, is suspected, it is recommended to perform follow-up tests for antibodies to tRNA synthetases; antigens are included in disease specific immunoassays (ie, inflammatory myopathy profile*) <a href="#">26, 33</a></li> <li>► If AIM, in particular necrotizing myopathy, is suspected, it is recommended to perform follow-up tests for anti-SRP antibodies; the antigen is included in disease specific immunoassays (ie, inflammatory myopathy profile*) <a href="#">26</a></li> </ul> <p>Notes: The fine distinction between AC-19 and -20 may depend on HEp-2 substrates and/or antibody concentration; antibodies to both RibP as well as tRNA synthetases may be undetected in HEp-2 IIFA-screening.</p>	
AC-20	<p>FINE SPECKLED</p> <ul style="list-style-type: none"> <li>► Found in patients with the anti-synthetase syndrome (a subset of AIM), interstitial lung disease, polyarthritis, Raynaud's phenomenon, and mechanic's hands; these features may occur in various combinations or as an isolated manifestation, especially interstitial lung disease <a href="#">33, 90</a></li> <li>► Autoantibodies associated with the AC-20 pattern are primarily reported for the anti-Jo-1 antibody, which recognizes histidyl-tRNA synthetase; since AC-20 is not specific for Jo-1, it is recommended to perform a follow-up test for anti-Jo-1 antibodies; the antigen is included in the routine ENA profile, as well as in disease specific immunoassays (ie, inflammatory myopathy profile*); the anti-Jo-1 antibodies are included in the classification criteria for AIM <a href="#">91, 92</a></li> </ul> <p>Notes: The fine distinction between AC-19 and -20 may depend on HEp-2 substrates and/or antibody concentration; antibodies to Jo-1 may be undetected in HEp-2 IIFA-screening.</p>	

Continued

Table 2 Continued

Code	AC pattern—clinical relevance	Refs
AC-21	<p>RETICULAR/AMA</p> <ul style="list-style-type: none"><li>Commonly found in PBC, but also detected in SSs, including PBC-SSs overlap syndrome and PBC-SjS overlap syndrome</li><li>If PBC is clinically suspected it is recommended to perform a follow-up test for AMA, historically detected by IIFA on rodent tissue (liver, stomach, kidney); these autoantibodies are primarily directed to the PDH complex, and in particular the E2-subunit (PDH-E2); the antigen is included in disease specific immunoassays (i.e., liver profile*) as well as in some routine ENA profiles</li><li>Additional antigens recognized include the E1α and E1β subunits of PDH, the E3-binding protein of PDH, and the 2-OGDC; these antigens are only included in extended disease specific immunoassays (i.e., extended liver profile*)</li></ul>	<p>93–97</p> <p>93, 94</p> <p>93, 94</p>
AC-22	<p>POLAR/GOLGI-like (see online supplementary table S2 for further details)</p> <ul style="list-style-type: none"><li>Found in small numbers of patients with a variety of conditions</li><li>Antigens recognized include giantin/macrogolgin and distinct golgin molecules; specific immunoassays to detect autoantibodies directed to specific Golgi antigens are currently not commercially available</li></ul>	<p>85</p>
AC-23	<p>RODS and RINGS (see (online supplementary table S2 for further details)</p> <ul style="list-style-type: none"><li>Most commonly found in HCV patients who have been treated with pegylated interferon-α/ribavirin combination therapy, but autoantibodies revealing the AC-23 patterns were undetected prior to treatment; as the use of interferon-α/ribavirin in HCV treatment is decreasing, the frequency and clinical associations of the AC-23 pattern may change</li><li>Specific immunoassays to detect autoantibodies directed to specific Rods and Rings antigens, for instance IMPDH2, are not commercially available</li></ul> <p>Note: Presence of the AC-23 pattern depends on the Hep-2 cell substrate.</p>	<p>98–101</p>

\*Availability of the inflammatory myopathy profile, the SSs profile and the (extended) liver profile may be limited to specialty clinical laboratories. AIH, autoimmune hepatitis; AIM, autoimmune myopathy; AMA, anti-mitochondrial antibodies; APS, antiphospholipid syndrome; Ago, argonaute protein; CENP, centromere-associated protein; CLIP, class II-associated invariant chain peptide; DFS, dense fine speckled; DM, dermatomyositis; EEA, early endosome antigen; ENA, extractable nuclear antigens; HCV, hepatitis C virus; IFA, indirect immunofluorescence assay; LAP, lamin-associated polypeptide; LBR, lamin B receptor; LEDGF, lens epithelial derived growth factor; NOR, nucleolus organizer region; NXP, nuclear matrix protein; PBC, primary biliary cholangitis; PCNA, proliferating cell nuclear antigen; PML, promyelocytic leukaemia; PM/ScI, polymyositis-scleroderma; RA, rheumatoid arthritis; RNAPol, ribonucleic acid polymerase; RNP, ribonucleoprotein; SARD, systemic autoimmune rheumatic diseases; SLE, systemic lupus erythematosus; SMN, survival of motor neuron; SRP, signal recognition protein; SSs, systemic sclerosis; SjS, Sjögren's syndrome; TIF, transcription intermediary factor; TRIM, tripartite motif; Tpr, translocated promoter region; dsDNA, double stranded deoxyribonucleic acid; HUBF, human upstream binding factor; tRNA, transfer ribonucleic acid.

taking into account that the Hep-2 IIFA nowadays is ordered by a wide spectrum of clinical disciplines,<sup>1</sup> the number of clinically unexpected positive results, that is, positive test results with no clinical evidence of an associated autoimmune disease, is ever increasing and may even equal the likelihood of a clinically true-positive result.<sup>140 141</sup> A study performed in a community setting concluded that many patients with a positive ANA test are incorrectly given a diagnosis of systemic lupus erythematosus and sometimes even treated with toxic medications.<sup>142</sup> These arguments are used to introduce a gating strategy in order to restrict test-ordering to those cases that have a sufficiently high pretest probability for having a SARD. However, it can also be argued that patients with a low pretest probability should be tested using the Hep-2 IIFA in order to prevent true cases, especially those with very early disease manifestations, from being missed. This is a paradigm shift to disease prediction and prevention.<sup>143</sup> In this strategy, the Hep-2 IIFA could be integrated in multianalyte ‘omic’ profiles for case finding and establishing an early diagnosis and preventing severe complications.<sup>143 144</sup> Obviously, it is anticipated that the added value of the Hep-2 IIFA in this approach can be increased by incorporating information on both patterns as well as titres in combination with well-directed advices on follow-up testing.

Although the current consensus on the clinical relevance of Hep-2 IIFA patterns has come across after extensive discussion and debate within the ICAP executive board as well as with the workshop participants, the information provided is not based on a systematic review or meta-analysis of the existing literature. Because of the short history of ICAP, being founded in 2014, inclusion of older literature might have been hampered

by potential differences in pattern nomenclature and definitions. For instance, the nuclear dense fine speckled (AC-2) and topo I-like (AC-29) patterns were previously often considered homogeneous, speckled or even mixed patterns. The centromere pattern (AC-3) or the cytoplasmic reticular/AMA (AC-21) patterns, on the other hand, are examples that probably have been less prone to change in pattern definition over time. The universal use of the ICAP nomenclature and pattern definitions, both in daily clinical practice as well as in the scientific literature, may enable systematic reviews in the future, and may well fine-tune current consensus based on expert opinions only.

In conclusion, the consensus statements on clinical relevance should be readily available to clinicians and this will enable further harmonisation of test-result interpretation with respect to Hep-2 IIFA patterns. Obviously, clinicians should be aware of the clinical suspicion for the respective patient, and therefore should order specific tests accordingly, also taking into account the anticipation of prevalence of Hep-2 IIFA negative (AC-0)<sup>13</sup> results in SARD. The information on clinical relevance of Hep-2 IIFA patterns is intended to support the decision strategy of the clinician. Information presented in the online supplementary tables 1–3 is primarily intended to be used for complex cases in the consultation of the laboratory specialist by the clinician. Depending on various jurisdictional regulations, follow-up testing can be automated in predefined algorithms which eventually will shorten the diagnostic delay. Eventually, appropriate integration of Hep-2 IIFA pattern information may help to better define disease criteria and even enable a paradigm shift in the pretest probability paradox.

## Recommendation

Table 3 Mitotic HEp-2 IIFA patterns

Code	AC pattern—clinical relevance	Refs
AC-24	<p>CENTROSOME (see online supplementary table 3 for further details)</p> <ul style="list-style-type: none"> <li>▶ The AC-24 pattern has low positive predictive value for any disease</li> <li>▶ Within the spectrum of the SARD, the AC-24 pattern is found in patients with Raynaud's phenomenon, localized scleroderma, SSc, SLE and RA, either alone or in combination with other SSc-associated antibodies;</li> <li>▶ Antigens recognized include <math>\alpha</math>-enolase, <math>\gamma</math>-enolase, ninein, Cep-250, Mob1, PCM-1/2, pericentrin; specific immunoassays for these autoantibodies are currently not commercially available</li> </ul>	102–105 104, 106–108
AC-25	<p>SPINDLE FIBERS (see online supplementary table 3 for further details)</p> <ul style="list-style-type: none"> <li>▶ The AC-25 pattern has low positive predictive value for any disease</li> <li>▶ Found very infrequently in a routine serology diagnostic setting</li> <li>▶ Antigen recognized includes HsEg5; specific immunoassays for this autoantibody, or other spindle fiber targets, are currently not commercially available</li> </ul>	109 109 110, 111
AC-26	<p>NuMA-like</p> <ul style="list-style-type: none"> <li>▶ Approximately one-half of the patients with the AC-26 pattern have clinical features of a SARD (SJS, SLE, UCTD, limited SSc, or RA); the AC-26 pattern is also observed in patients with organ-specific autoimmune diseases and less frequently in non-autoimmune conditions, especially when in low titer</li> <li>▶ Found very infrequently in a routine serology diagnostic setting</li> <li>▶ Antigens recognized include NuMA, centrophilin, SP-H antigen and NMP-22; specific immunoassays for these autoantibodies are currently not commercially available</li> </ul>	109, 111–114 109 115
AC-27	<p>INTERCELLULAR BRIDGE (see online supplementary table 3 for further details)</p> <ul style="list-style-type: none"> <li>▶ The AC-27 pattern has low positive predictive value for any disease</li> <li>▶ Found very infrequently in a routine serology diagnostic setting</li> <li>▶ Antigens recognized include, among other, CENP-E, CENP-F, TD60, MSA36, KIF-14, MKLP-1, MPP1/KIF20B, and INCENP; specific immunoassays for these autoantibodies are currently not commercially available</li> </ul>	116 117 116, 118, 119
AC-28	<p>MITOTIC CHROMOSOMAL (see online supplementary table 3 for further details)</p> <ul style="list-style-type: none"> <li>▶ The AC-28 pattern has low positive predictive value for any disease</li> <li>▶ Found very infrequently in a routine serology diagnostic setting</li> <li>▶ Antigens recognized include DCA, MCA1, and MCA5; specific immunoassays for these autoantibodies are currently not commercially available</li> </ul>	120 120–122

CENP, centromere-associated protein; Cep, centrosomal protein; DCA, dividing cell antigen; IIFA, indirect immunofluorescence assay; INCENP, inner centromere protein; KIF, kinesin family; MCA, mitotic chromosomal antigen; MKLP, mitotic kinesin-like protein; MPP, M-phase phosphoprotein; MSA, mitotic spindle apparatus; NMP, nuclear matrix protein; NuMA, nuclear mitotic apparatus; PCM, pericentriolar material; RA, rheumatoid arthritis; SARD, systemic autoimmune rheumatic diseases; SLE, systemic lupus erythematosus; SSc, systemic sclerosis; SJS, Sjögren's syndrome; UCTD, undifferentiated connective tissue disease.

## Author affiliations

<sup>1</sup>Central Diagnostic Laboratory, Maastricht University Medical Center, Maastricht, The Netherlands

<sup>2</sup>Rheumatology Division, Universidade Federal de São Paulo, São Paulo, Brazil

<sup>3</sup>Department of Immunology, Instituto Universitario del Hospital Italiano de Buenos Aires, Buenos Aires, Argentina

<sup>4</sup>Laboratory of Immunology, Hospital General de Agudos Carlos G Durand, Buenos Aires, Argentina

<sup>5</sup>Immunology, Medical Faculty TU Dresden, Dresden, Germany

<sup>6</sup>Pontificia Universidade Católica de Goiás, Goiás, Brazil

<sup>7</sup>Medicine, Health Sciences Centre, Calgary, Alberta, Canada

<sup>8</sup>Department of Immunology and Rheumatology, Hospital General de Occidente, Guadalajara, Mexico

<sup>9</sup>Rheumatology Unit, Clinical Department of General Internal Medicine, Innsbruck Medical University, Innsbruck, Austria

<sup>10</sup>Department of Internal Medicine II, Medical University of Innsbruck, Innsbruck, Austria

<sup>11</sup>Pontificia Universidade Católica de Goiás, Goiânia, Brazil

<sup>12</sup>Department of Rheumatology and Clinical Immunology, Kyoto University Graduate School of Medicine, Kyoto, Japan

<sup>13</sup>Brazilian Society of Autoimmunity, Porto Alegre, Brazil

<sup>14</sup>Department of Clinical Nursing, University of Occupational and Environmental Health, Kitakyushu, Japan

<sup>15</sup>Department of Oral Biology, University of Florida, Gainesville, Florida, USA

**Acknowledgements** We thank all the workshop participants for their constructive comments and fruitful discussions.

**Contributors** All authors actively participated in the respective workshops in Kyoto and Dresden. They also participated in the discussions of the executive ICAP committee. The draft of the manuscript was made by JD and was commented on by all authors. Final discussions have taken place at the international autoimmunity meeting in Lisbon. Required amendments were made by JD and approved by all authors.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** The ICAP committee is funded by unrestricted educational grants by several in vitro diagnostics companies (for details see [www.anapatterns.org/sponsors.php](http://www.anapatterns.org/sponsors.php)). JD has received lecture fees from Euroimmun and Thermo Fisher. MJF is a consultant to Inova Diagnostics and Werfen International; none of the other authors declare any competing interest.

**Patient consent for publication** Not required.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data sharing statement** No additional data are available.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## REFERENCES

- Mahler M, Meroni P-L, Bossuyt X, *et al*. Current concepts and future directions for the assessment of autoantibodies to cellular antigens referred to as anti-nuclear antibodies. *J Immunol Res* 2014;2014:1–18.
- Solomon DH, Kavanaugh AJ, Schur PH, *et al*. Evidence-based guidelines for the use of immunologic tests: antinuclear antibody testing. *Arthritis Rheum* 2002;47:434–44.
- Meroni PL, Schur PH. ANA screening: an old test with new recommendations. *Ann Rheum Dis* 2010;69:1420–2.
- Damoiseaux JGMC, Cohen Tervaert JW. From ANA to ena: how to proceed? *Autoimmun Rev* 2006;5:10–17.
- Op De Beeck K, Vermeersch P, Verschueren P, *et al*. Detection of antinuclear antibodies by indirect immunofluorescence and by solid phase assay. *Autoimmun Rev* 2011;10:801–8.
- Oyaert M, Bossuyt X, Ravelingien I, *et al*. Added value of indirect immunofluorescence intensity of automated antinuclear antibody testing in a secondary hospital setting. *Clin Chem Lab Med* 2016;54:e63–6.
- Agmon-Levin N, Damoiseaux J, Kallenberg C, *et al*. International recommendations for the assessment of autoantibodies to cellular antigens referred to as anti-nuclear antibodies. *Ann Rheum Dis* 2014;73:17–23.
- van den Hoogen F, Khanna D, Fransen J, *et al*. 2013 classification criteria for systemic sclerosis: an American College of rheumatology/European League against rheumatism collaborative initiative. *Ann Rheum Dis* 2013;72:1747–55.
- Mariz HA, Sato EI, Barbosa SH, *et al*. Pattern on the antinuclear antibody-HEp-2 test is a critical parameter for discriminating antinuclear antibody-positive healthy individuals and patients with autoimmune rheumatic diseases. *Arthritis & Rheumatism* 2011;63:191–200.
- Wiik AS, Høier-Madsen M, Forslid J, *et al*. Antinuclear antibodies: a contemporary nomenclature using HEp-2 cells. *J Autoimmun* 2010;35:276–90.
- Chan EKL, Damoiseaux J, Carballo OG, *et al*. Report of the first international consensus on standardized Nomenclature of antinuclear antibody HEp-2 cell patterns 2014–2015. *Front Immunol* 2015;6.
- Damoiseaux J, von Mühlén CA, García-De La Torre I, Torre G-de la I, *et al*. International consensus on ANA patterns (ICAP): the bumpy road towards a consensus on reporting ANA results. *Autoimmun Highlights* 2016;7.



- 13 Herold M, Klotz W, Andrade LEC, *et al.* International consensus on antinuclear antibody patterns: defining negative results and reporting unidentified patterns. *Clin Chem Lab Med* 2018;56:1799–802.
- 14 Andrade LEC, Klotz W, Herold M, *et al.* International consensus on antinuclear antibody patterns: definition of the ac-29 pattern associated with antibodies to DNA topoisomerase I. *Clin Chem Lab Med* 2018;56:1783–8.
- 15 Conrad K, Schössler W, Hiepe F. *Autoantibodies in systemic autoimmune diseases: a diagnostic reference*. 2. 3th edn. Autoantigens autoantibodies autoimmunity, 2015.
- 16 Petri M, Orbai AM, Alarcón GS, *et al.* Derivation and validation of the systemic lupus international collaborating clinics classification criteria for systemic lupus erythematosus. *Arthritis Rheum* 2012;64:2677–86.
- 17 Conrad K, Schössler W, Hiepe F. *Autoantibodies in organ specific autoimmune diseases. a diagnostic reference*. 8. 2th edn. Autoantigens autoantibodies autoimmunity, 2017.
- 18 Dellavance A, Gallindo C, Soares MG, *et al.* Redefining the Scl-70 indirect immunofluorescence pattern: autoantibodies to DNA topoisomerase I yield a specific compound immunofluorescence pattern. *Rheumatology* 2009;48:632–7.
- 19 European Association for the Study of the Liver. EASL clinical practice guidelines: autoimmune hepatitis. *J Hepatol* 2015;62:971–1004.
- 20 Watanabe A, Koda M, Sugiura K, *et al.* Anti-DFS70 antibodies in 597 healthy hospital workers. *Arthritis Rheum* 2004;50:892–900.
- 21 Mahler M, Parker T, Peebles CL, *et al.* Anti-DFS70/LEDGF antibodies are more prevalent in healthy individuals compared to patients with systemic autoimmune rheumatic diseases. *J Rheumatol* 2012;39:2104–10.
- 22 Ochs RL, Mahler M, Basu A, *et al.* The significance of autoantibodies to DFS70/LEDGFp75 in health and disease: integrating basic science with clinical understanding. *Clin Exp Med* 2016;16:273–93.
- 23 Johnson SR, Fransen J, Khanna D, *et al.* Validation of potential classification criteria for systemic sclerosis. *Arthritis Care Res* 2012;64:358–67.
- 24 Mehra S, Walker J, Patterson K, *et al.* Autoantibodies in systemic sclerosis. *Autoimmun Rev* 2013;12:340–54.
- 25 Shiboski CH, Shiboski SC, Seror R, *et al.* American College of Rheumatology/ European League against rheumatism classification criteria for primary Sjögren's syndrome: a consensus and data-driven methodology involving three international patient cohorts. *Ann Rheum Dis* 2016;2017:9–16.
- 26 Betteridge Z, McHugh N. Myositis-specific autoantibodies: an important tool to support diagnosis of myositis. *J Intern Med* 2016;280:8–23.
- 27 Trallero-Araguás E, Rodrigo-Pendás JA, Selva-O'Callaghan A, *et al.* Usefulness of anti-p155 autoantibody for diagnosing cancer-associated dermatomyositis: a systematic review and meta-analysis. *Arthritis Rheum* 2012;64:523–32.
- 28 Bossuyt X, Frans J, Hendrickx A, *et al.* Detection of anti-SSA antibodies by indirect immunofluorescence. *Clin Chem* 2004;50:2361–9.
- 29 Northway JD, Tan EM. Differentiation of antinuclear antibodies giving Speckled staining patterns in immunofluorescence. *Clin Immunol Immunopathol* 1972;1:140–54.
- 30 Satoh M, Fritzler MJ, Chan EKL. Antihistone and antisplinesomal antibodies. In: Lahita RG, Tsokos G, Buyon JP, *et al.* eds. *Systemic lupus erythematosus*. San Diego: Academic Press, 2011: 275–92.
- 31 Satoh M, Vázquez-Del Mercado M, Chan EK. Clinical interpretation of antinuclear antibody tests in systemic rheumatic diseases. *Mod Rheumatol* 2009;19:219–28.
- 32 Sharp GC, Irwin WS, Tan EM, *et al.* Mixed connective tissue disease—an apparently distinct rheumatic disease syndrome associated with a specific antibody to an extractable nuclear antigen (ENA). *Am J Med* 1972;52:148–59.
- 33 Satoh M, Tanaka S, Ceribelli A, *et al.* A comprehensive overview on myositis-specific antibodies: new and old biomarkers in idiopathic inflammatory myopathy. *Clin Rev Allergy Immunol* 2017;52:1–19.
- 34 Cozzani E, Drosera M, Riva S, *et al.* Analysis of a multiple nuclear dots pattern in a large cohort of dermatological patients. *Clin Lab* 2012;58:329–32.
- 35 Hu SL, Zhao FR, Hu Q, *et al.* Meta-analysis assessment of gp210 and Sp100 for the diagnosis of primary biliary cirrhosis. *PLoS One* 2014;9:e101916.
- 36 Granito A, Yang WH, Muratori L, *et al.* PML nuclear body component SP140 is a novel autoantigen in primary biliary cirrhosis. *Am J Gastroenterol* 2010;105:125–31.
- 37 Ichimura Y, Matsushita T, Hamaguchi Y, *et al.* Anti-NXP2 autoantibodies in adult patients with idiopathic inflammatory myopathies: possible association with malignancy. *Ann Rheum Dis* 2012;71:710–3.
- 38 Benveniste O, Stenzel W, Allenbach Y. Advances in serological diagnostics of inflammatory myopathies. *Curr Opin Neurol* 2016;29:662–73.
- 39 Ceribelli A, Fredi M, Taraborelli M, *et al.* Anti-MJ/NXP-2 autoantibody specificity in a cohort of adult Italian patients with polymyositis/dermatomyositis. *Arthritis Res Ther* 2012;14.
- 40 Onouchi H, Muro Y, Tomita Y. Clinical features and IgG subclass distribution of anti-p80 coilin antibodies. *J Autoimmun* 1999;13:225–32.
- 41 Fujimoto M, Kikuchi K, Tamaki T, *et al.* Distribution of anti-p80-coilin autoantibody in collagen diseases and various skin diseases. *Br J Dermatol* 1997;137:916–20.
- 42 Andrade LE, Chan EK, Raska I, *et al.* Human autoantibody to a novel protein of the nuclear coiled body: immunological characterization and cDNA cloning of p80-coilin. *J Exp Med* 1991;173:1407–19.
- 43 Satoh M, Chan JY, Ross SJ, *et al.* Autoantibodies to survival of motor neuron complex in patients with polymyositis: immunoprecipitation of D, E, F, and G proteins without other components of small nuclear ribonucleoproteins. *Arthritis Rheum* 2011;63:1972–8.
- 44 Mahler M, Fritzler MJ, Satoh M. Autoantibodies to the mitochondrial RNA processing (MRP) complex also known as Th/To autoantigen. *Autoimmun Rev* 2015;14:254–7.
- 45 Ceribelli A, Cavazzana I, Franceschini F, *et al.* Anti-Th/To are common antinuclear autoantibodies in Italian patients with scleroderma. *J Rheumatol* 2010;37:2071–5.
- 46 Mahler M, Rajmakers R. Novel aspects of autoantibodies to the PM/ScI complex: clinical, genetic and diagnostic insights. *Autoimmun Rev* 2007;6:432–7.
- 47 Mahler M, Gascon C, Patel S, *et al.* Rpp25 is a major target of autoantibodies to the Th/To complex as measured by a novel chemiluminescent assay. *Arthritis Res Ther* 2013;15.
- 48 Okano Y, Steen VD, Medsger TA. Autoantibody to U3 nucleolar ribonucleoprotein (fibrillarin) in patients with systemic sclerosis. *Arthritis Rheum* 1992;35:95–100.
- 49 Arnett FC, Reveille JD, Goldstein R, *et al.* Autoantibodies to fibrillarin in systemic sclerosis (scleroderma). An immunogenetic, serologic, and clinical analysis. *Arthritis Rheum* 1996;39:1151–60.
- 50 Tormey VJ, Bunn CC, Denton CP, *et al.* Anti-fibrillarin antibodies in systemic sclerosis. *Rheumatology* 2001;40:1157–62.
- 51 Nandiwada SL, Peterson LK, Mayes MD, *et al.* Ethnic differences in autoantibody diversity and hierarchy: More clues from a US cohort of patients with systemic sclerosis. *J Rheumatol* 2016;43:1816–24.
- 52 Reimer G, Rose KM, Scheer U, *et al.* Autoantibody to RNA polymerase I in scleroderma sera. *J Clin Invest* 1987;79:65–72.
- 53 Kuwana M, Kaburaki J, Mimori T, *et al.* Autoantibody reactive with three classes of RNA polymerases in sera from patients with systemic sclerosis. *J Clin Invest* 1993;91:1399–404.
- 54 Fritzler MJ, von Muhlen CA, Toffoli SM, *et al.* Autoantibodies to the nucleolar organizer antigen NOR-90 in children with systemic rheumatic diseases. *J Rheumatol* 1995;22:521–4.
- 55 Fujii T, Mimori T, Akizuki M. Detection of autoantibodies to nucleolar transcription factor nor 90/hUBF in sera of patients with rheumatic diseases, by recombinant autoantigen-based assays. *Arthritis Rheum* 1996;39:1313–8.
- 56 Imai H, Ochs RL, Kiyosawa K, *et al.* Nucleolar antigens and autoantibodies in hepatocellular carcinoma and other malignancies. *Am J Pathol* 1992;140:859–70.
- 57 Yamasaki Y, Honkanen-Scott M, Hernandez L, *et al.* Nucleolar staining cannot be used as a screening test for the scleroderma marker anti-RNA polymerase I/III antibodies. *Arthritis Rheum* 2006;54:3051–6.
- 58 Coppo P, Clauvel JP, Bengoufa D, *et al.* Autoimmune cytopenias associated with autoantibodies to nuclear envelope polypeptides. *Am J Hematol* 2004;77:241–9.
- 59 Konstantinov K, Foisner R, Byrd D, *et al.* Integral membrane proteins associated with the nuclear lamina are novel autoimmune antigens of the nuclear envelope. *Clin Immunol Immunopathol* 1995;74:89–99.
- 60 Reeves WH, Chaudhary N, Salerno A, *et al.* Lamin B autoantibodies in sera of certain patients with systemic lupus erythematosus. *J Exp Med* 1987;165:750–62.
- 61 Miyachi K, Hankins RW, Matsushima H, *et al.* Profile and clinical significance of anti-nuclear envelope antibodies found in patients with primary biliary cirrhosis: a multicenter study. *J Autoimmun* 2003;20:247–54.
- 62 Courvalin JC, Lassoued K, Bartnik E, *et al.* The 210-kD nuclear envelope polypeptide recognized by human autoantibodies in primary biliary cirrhosis is the major glycoprotein of the nuclear pore. *J Clin Invest* 1990;86:279–85.
- 63 Nickowitz RE, Worman HJ. Autoantibodies from patients with primary biliary cirrhosis recognize a restricted region within the cytoplasmic tail of nuclear pore membrane glycoprotein gp210. *J Exp Med* 1993;178:2237–42.
- 64 Bandin O, Courvalin JC, Poupon R, *et al.* Specificity and sensitivity of gp210 autoantibodies detected using an enzyme-linked immunosorbent assay and a synthetic polypeptide in the diagnosis of primary biliary cirrhosis. *Hepatology* 1996;23:1020–4.
- 65 Wesierska-Gadek J, Klima A, Komina O, *et al.* Characterization of autoantibodies against components of the nuclear pore complexes: high frequency of anti-p62 nucleoporin antibodies. *Ann N Y Acad Sci* 2007;1109:519–30.
- 66 Kraemer DM, Tony HP. Nuclear pore protein p62 autoantibodies in systemic lupus erythematosus. *Open Rheumatol J* 2010;4:24–7.
- 67 Courvalin JC, Lassoued K, Worman HJ, *et al.* Identification and characterization of autoantibodies against the nuclear envelope lamin B receptor from patients with primary biliary cirrhosis. *J Exp Med* 1990;172:961–7.
- 68 Ou Y, Enarson P, Rattner JB, *et al.* The nuclear pore complex protein TPR is a common autoantigen in sera that demonstrate nuclear envelope staining by indirect immunofluorescence. *Clin Exp Immunol* 2004;136:379–87.
- 69 Miyachi K, Fritzler MJ, Tan EM. Autoantibody to a nuclear antigen in proliferating cells. *J Immunol* 1978;121:2228–34.
- 70 Vermeersch P, De Beeck KO, Lauwerys BR, *et al.* Antinuclear antibodies directed against proliferating cell nuclear antigen are not specifically associated with systemic lupus erythematosus. *Ann Rheum Dis* 2009;68:1791–3.
- 71 Mahler M, Burlingame RW, Wu J, *et al.* Serological and clinical associations of anti-PCNA antibodies in systemic lupus erythematosus detected by a novel chemiluminescence assay. *Arthritis Rheum* 2011;63.

- 72 Mahler M, Miyachi K, Peebles C, *et al.* The clinical significance of autoantibodies to the proliferating cell nuclear antigen (PCNA). *Autoimmun Rev* 2012;11:771–5.
- 73 Hsu TC, Tsay GJ, Chen TY, *et al.* Anti-PCNA autoantibodies preferentially recognize C-terminal of PCNA in patients with chronic hepatitis B virus infection. *Clin Exp Immunol* 2006;144:110–6.
- 74 Casiano CA, Landberg G, Ochs RL, *et al.* Autoantibodies to a novel cell cycle-regulated protein that accumulates in the nuclear matrix during S phase and is localized in the kinetochores and spindle midzone during mitosis. *J Cell Sci* 1993;106:1045–56.
- 75 Casiano CA, Humbel RL, Peebles C, *et al.* Autoimmunity to the cell cycle-dependent centromere protein p330d/CENP-F in disorders associated with cell proliferation. *J Autoimmun* 1995;8:575–86.
- 76 Rattner JB, Rees J, Whitehead CM, *et al.* High frequency of neoplasia in patients with autoantibodies to centromere protein CENP-F. *Clin Invest Med* 1997;20:308–19.
- 77 Bencimon C, Salles G, Moreira A, *et al.* Prevalence of anticentromere F protein autoantibodies in 347 patients with non-Hodgkin's lymphoma. *Ann N Y Acad Sci* 2005;1050:319–26.
- 78 Welner S, Trier NH, Frisch M, *et al.* Correlation between centromere protein-F autoantibodies and cancer analyzed by enzyme-linked immunosorbent assay. *Mol Cancer* 2013;12.
- 79 Basu D, Reveille JD. Anti-scl-70. *Autoimmunity* 2005;38:65–72.
- 80 Liberal R, Grant CR, Longhi MS, *et al.* Diagnostic criteria of autoimmune hepatitis. *Autoimmun Rev* 2014;13:435–40.
- 81 Lidman K, Biberfeld G, Fagraeus A, *et al.* Anti-actin specificity of human smooth muscle antibodies in chronic active hepatitis. *Clin Exp Immunol* 1976;24:266–72.
- 82 Fusconi M, Cassani F, Zauli D, *et al.* Anti-actin antibodies: a new test for an old problem. *J Immunol Methods* 1990;130:1–8.
- 83 Chretien-Leprince P, Ballot E, Andre C, *et al.* Diagnostic value of anti-F-actin antibodies in a French multicenter study. *Ann N Y Acad Sci* 2005;1050:266–73.
- 84 Bhanji RA, Eystathiou T, Chan EK, *et al.* Clinical and serological features of patients with autoantibodies to GW/P bodies. *Clin Immunol* 2007;125:247–56.
- 85 Stinton LM, Eystathiou T, Selak S, *et al.* Autoantibodies to protein transport and messenger RNA processing pathways: endosomes, lysosomes, Golgi complex, proteasomes, assemblyosomes, exosomes, and GW bodies. *Clin Immunol* 2004;110:30–44.
- 86 Sciascia S, Bertolaccini ML, Roccatello D, *et al.* Autoantibodies involved in neuropsychiatric manifestations associated with systemic lupus erythematosus: a systematic review. *J Neurol* 2014;261:1706–14.
- 87 Yura H, Sakamoto N, Satoh M, *et al.* Clinical characteristics of patients with anti-aminoacyl-tRNA synthetase antibody positive idiopathic interstitial pneumonia. *Respir Med* 2017;132:189–94.
- 88 Valões CC, Molinari BC, Pitta AC, *et al.* Anti-ribosomal P antibody: a multicenter study in childhood-onset systemic lupus erythematosus patients. *Lupus* 2017;26:484–9.
- 89 Mahler M, Kessenbrock K, Szymrka M, *et al.* International multicenter evaluation of autoantibodies to ribosomal P proteins. *Clin Vaccine Immunol* 2006;13:77–83.
- 90 Marie I, Hatron PY, Cherin P, *et al.* Functional outcome and prognostic factors in anti-Jo1 patients with antisynthetase syndrome. *Arthritis Res Ther* 2013;15.
- 91 Fritzler MJ, Choi MY, Mahler M. The antinuclear antibody test in the diagnosis of antisynthetase syndrome and other autoimmune myopathies. *J Rheumatol* 2018;45:444.1–5.
- 92 Lundberg IE, Tjærnlund A, Bottai M, *et al.* 2017 European league against rheumatism/American college of rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies and their major subgroups. *Ann Rheum Dis* 2017;76:1955–64.
- 93 European Association for the Study of the Liver. EASL clinical practice guidelines: the diagnosis and management of patients with primary biliary cholangitis. *J Hepatol* 2017;67:145–72.
- 94 Shuai Z, Wang J, Badamagunta M, *et al.* The fingerprint of antimitochondrial antibodies and the etiology of primary biliary cholangitis. *Hepatology* 2017;65:1670–82.
- 95 Skare TL, Nishihara RM, Haider O, *et al.* Liver autoantibodies in patients with scleroderma. *Clin Rheumatol* 2011;30:129–32.
- 96 Zheng B, Vincent C, Fritzler MJ, *et al.* Prevalence of systemic sclerosis in primary biliary cholangitis using the new ACR/EULAR classification criteria. *J Rheumatol* 2017;44:33–9.
- 97 Nardi N, Brito-Zerón P, Ramos-Casals M, *et al.* Circulating auto-antibodies against nuclear and non-nuclear antigens in primary Sjögren's syndrome: prevalence and clinical significance in 335 patients. *Clin Rheumatol* 2006;25:341–6.
- 98 Covini G, Carcamo WC, Bredi E, *et al.* Cytoplasmic rods and rings autoantibodies developed during pegylated interferon and ribavirin therapy in patients with chronic hepatitis C. *Antivir Ther* 2012;17:805–11.
- 99 Calise SJ, Keppeke GD, Andrade LE, *et al.* Anti-rods/rings: a human model of drug-induced autoantibody generation. *Front Immunol* 2015;6.
- 100 Novembrino C, Aghemo A, Ferraris Fusarini C, *et al.* Interferon-ribavirin therapy induces serum antibodies determining 'rods and rings' pattern in hepatitis C patients. *J Viral Hepat* 2014;21:944–9.
- 101 Keppeke GD, Calise SJ, Chan EK, *et al.* Anti-rods/rings autoantibody generation in hepatitis C patients during interferon- $\alpha$ /ribavirin therapy. *World J Gastroenterol* 2016;22:1966–74.
- 102 Takahashi T, Asano Y, Hirakawa M, *et al.* Linear scleroderma with prominent multiple lymphadenopathy followed by the development of polymyositis: a case report and review of published work. *J Dermatol* 2016;43:1224–7.
- 103 Howng SL, Chou AK, Lin CC, *et al.* Autoimmunity against hNinein, a human centrosomal protein, in patients with rheumatoid arthritis and systemic lupus erythematosus. *Mol Med Rep* 2011;4:825–30.
- 104 Hamaguchi Y, Matsushita T, Hasegawa M, *et al.* High incidence of pulmonary arterial hypertension in systemic sclerosis patients with anti-centriole autoantibodies. *Mod Rheumatol* 2015;25:798–801.
- 105 Gavanescu I, Vazquez-Abad D, McCauley J, *et al.* Centrosome proteins: a major class of autoantigens in scleroderma. *J Clin Immunol* 1999;19:166–71.
- 106 Fritzler MJ, Zhang M, Stinton LM, *et al.* Spectrum of centrosome autoantibodies in childhood varicella and post-varicella acute cerebellar ataxia. *BMC Pediatr* 2003;3.
- 107 Rattner JB, Martin L, Waisman DM, *et al.* Autoantibodies to the centrosome (centriole) react with determinants present in the glycolytic enzyme enolase. *J Immunol* 1991;146:2341–4.
- 108 Mack GJ, Rees J, Sandblom O, *et al.* Autoantibodies to a group of centrosomal proteins in human autoimmune sera reactive with the centrosome. *Arthritis Rheum* 1998;41:551–8.
- 109 Mozo L, Gutiérrez C, Gómez J, Gámez Jes<sup>o</sup>s. Antibodies to mitotic spindle apparatus: clinical significance of NuMA and HsEg5 autoantibodies. *J Clin Immunol* 2008;28:285–90.
- 110 Whitehead CM, Winkfein RJ, Fritzler MJ, *et al.* The spindle kinesin-like protein HsEg5 is an autoantigen in systemic lupus erythematosus. *Arthritis Rheum* 1996;39:1635–42.
- 111 Szalat R, Ghillani-Dalbin P, Jallouli M, *et al.* Anti-NuMA1 and anti-NuMA2 (anti-HsEg5) antibodies: clinical and immunological features: a propos of 40 new cases and review of the literature. *Autoimmun Rev* 2010;9:652–6.
- 112 Grypiotis P, Ruffatti A, Tonello M, *et al.* [Clinical significance of fluoroscopic patterns specific for the mitotic spindle in patients with rheumatic diseases]. *Reumatismo* 2002;54:232–7.
- 113 Price CM, McCarty GA, Pettijohn DE. NuMA protein is a human autoantigen. *Arthritis Rheum* 1984;27:774–9.
- 114 Bonaci-Nikolic B, Andrejevic S, Bukilica M, *et al.* Autoantibodies to mitotic apparatus: association with other autoantibodies and their clinical significance. *J Clin Immunol* 2006;26:438–46.
- 115 Compton DA, Szilak I, Cleveland DW. Primary structure of NuMA, an intranuclear protein that defines a novel pathway for segregation of proteins at mitosis. *J Cell Biol* 1992;116:1395–408.
- 116 Rattner JB, Mack GJ, Fritzler MJ. Autoantibodies to components of the mitotic apparatus. *Mol Biol Rep* 1998;25:143–55.
- 117 Vermeersch P, Bossuyt X. Prevalence and clinical significance of rare antinuclear antibody patterns. *Autoimmun Rev* 2013;12:998–1003.
- 118 Humbel RL, Conrad K, ed. *Autoantibodies to mitotic cells. In Dresden autoantibody symposium.* Dresden Germany: Pabst Science, 2013: 327–39.
- 119 Rodríguez-Bayona B, Ruchaud S, Rodríguez C, *et al.* Autoantibodies against the chromosomal passenger protein INCENP found in a patient with Graham Little-Piccardi-Lassueur syndrome. *J Autoimmune Dis* 2007;4.
- 120 Gitlits VM, Macaulay SL, Toh BH, *et al.* Novel human autoantibodies to phosphopeptides on mitotic chromosomal autoantigens (MCAs). *J Invest Med* 2000;48:172–82.
- 121 Blaschek M, Muller S, Youinou P. Anti-"dividing cell antigen" autoantibody: a novel antinuclear antibody pattern related to histones in systemic lupus erythematosus. *J Clin Immunol* 1993;13:329–38.
- 122 Rayzman VM, SENTRY JW. MCA1 detection of histone H3 serine 10 phosphorylation, a novel biomarker for determination of mitotic indices. *Hum Antibodies* 2006;15:71–80.
- 123 Didier K, Bolko L, Giusti D, *et al.* Autoantibodies associated with connective tissue diseases: what meaning for clinicians? *Front Immunol* 2018;9.
- 124 Pham BN, Albareda S, Guyard A, *et al.* Impact of external quality assessment on antinuclear antibody detection performance. *Lupus* 2005;14:113–9.
- 125 Van Blerk M, Van Campenhout C, Bossuyt X, *et al.* Current practices in antinuclear antibody testing: results from the Belgian external quality assessment scheme. *Clin Chem Lab Med* 2009;47:102–8.
- 126 Foggia P, Percannella G, Soda P, *et al.* Benchmarking HEp-2 cells classification methods. *IEEE Trans Med Imaging* 2013;32:1878–89.
- 127 Bizzaro N, Antico A, Platzgummer S, *et al.* Automated antinuclear immunofluorescence antibody screening: a comparative study of six computer-aided diagnostic systems. *Autoimmun Rev* 2014;13:292–8.
- 128 Hobson P, Lovell BC, Percannella G, *et al.* Benchmarking human epithelial type 2 interphase cells classification methods on a very large dataset. *Artif Intell Med* 2015;65:239–50.
- 129 Hobson P, Lovell BC, Percannella G, *et al.* Computer aided diagnosis for anti-nuclear antibodies HEp-2 images: progress and challenges. *Pattern Recognit Lett* 2016;82:3–11.

- 130 Hobson P, Lovell BC, Percannella G, *et al.* HEp-2 staining pattern recognition at cell and specimen levels: datasets, algorithms and results. *Pattern Recognit Lett* 2016;82:12–22.
- 131 van Beers JJBC, Hahn M, Fraune J, *et al.* Performance analysis of automated evaluation of antinuclear antibody indirect immunofluorescent tests in a routine setting. *Autoimmun Highlights* 2018;9.
- 132 Infantino M, Shovman O, Pérez D, PÁ©rez D, *et al.* A better definition of the anti-DFS70 antibody screening by IIF methods. *J Immunol Methods* 2018;461:110–6.
- 133 Chan EK, Damoiseaux J, de Melo Cruvinel W, *et al.* Report on the second international consensus on ANA pattern (ICAP) workshop in Dresden 2015. *Lupus* 2016;25:797–804.
- 134 Mahler M, Meroni PL, Andrade LE, *et al.* Towards a better understanding of the clinical association of anti-DFS70 autoantibodies. *Autoimmun Rev* 2016;15:198–201.
- 135 Meroni PL, Chan EK, Damoiseaux J, *et al.* Unending story of the indirect immunofluorescence assay on HEp-2 cells: old problems and new solutions? *Ann Rheum Dis* 2019;78:e46.
- 136 Fritzler MJ. The antinuclear antibody test: last or lasting GASP? *Arthritis Rheum* 2011;63:19–22.
- 137 Pisetsky DS. Antinuclear antibody testing - misunderstood or misbegotten? *Nat Rev Rheumatol* 2017;13:495–502.
- 138 Pérez D, Gilburd B, Azoulay D, *et al.* Antinuclear antibodies: is the indirect immunofluorescence still the gold standard or should be replaced by solid phase assays? *Autoimmun Rev* 2018;17:548–52.
- 139 Tan EM, Feltkamp TE, Smolen JS, *et al.* Range of antinuclear antibodies in "healthy" individuals. *Arthritis Rheum* 1997;40:1601–11.
- 140 Abeles AM, Abeles M. The clinical utility of a positive antinuclear antibody test result. *Am J Med* 2013;126:342–8.
- 141 Avery TY, van de Cruys M, Austen J, *et al.* Anti-nuclear antibodies in daily clinical practice: prevalence in primary, secondary, and tertiary care. *J Immunol Res* 2014;2014:1–8.
- 142 Narain S, Richards HB, Satoh M, *et al.* Diagnostic accuracy for lupus and other systemic autoimmune diseases in the community setting. *Arch Intern Med* 2004;164:2435–41.
- 143 Fritzler MJ, Martinez-Prat L, Choi MY, *et al.* The utilization of autoantibodies in approaches to precision health. *Front Immunol* 2018;9.
- 144 Fritzler MJ. Choosing wisely: review and commentary on anti-nuclear antibody (ANA) testing. *Autoimmun Rev* 2016;15:272–80.