DeCORTISYL

PREDNISONE

and

PreCORTISYL

PREDNISOLONE

Prescribable on Form E.C.10

ADVANTAGES

Superior therapeutic effect.

Little or no sodium and water retention

INDICATED ESPECIALLY

when cortisone has failed or when electrolyte disturbance and water retention have occurred

DOSAGE: One quarter that of cortisone.

e.g. in adults: initially 20 to 40 mg. daily

maintenance 5 to 20 mg. daily

in children: initially $\frac{1}{2}$ to 1 mg./kg. daily

maintenance $\frac{1}{8}$ to $\frac{1}{4}$ mg./kg. daily

Transfer to DeCortisyl or PreCortisyl requires no special precautions, other than maintaining continuity of treatment in equivalent dosage.

Scored tablets of 5 mg.
Bottles of 30 and 100
Basic N.H.S. Price
(cost to chemist)
37/6 and 120/-



ECONOMY PRESCRIBING

without detriment to the treatment of patients

CREMALGIN is a long acting rubefacient of high quality supplied at realistic cost on E.C.10. Every CREMALGIN prescription issued for the treatment of Rheumatism, Fibrositis, Sciatica, Lumbago, Muscular Pain and associated conditions represents up to 50% saving to the National Health Service. Doctors are writing more than 150,000 prescriptions for rubefacient balms each month and have readily accepted this valuable means of sound N.H.S. Economy.

*Methyl Nicotinate 1.0% Glycol Salicylate 10.0% Histamine Dihydrochloride 0.1% Capsicin 0.1% Excipient q.s.

Basic Price to N.H.S.: 1 oz. dispensing tube 1/9d. 16 oz. dispensing jar, per oz.—1/2 d.

Note that these prices cover a full ounce prescribed

CREMALGIN

*** Long-acting Rubefacient**

WEST PHARMACEUTICAL COMPANY LTD.

82 VICTORIA STREET, LONDON, S.W.1

TEL: TAT 2580

No mere decoration



The tree-like Monstera is one of the most intriguing of all house plants and undoubtedly owes much of its popularity to the ease with which long-lasting decorative effects can be achieved under a variety of conditions. In antibiotic medicine, however, it is practical effectiveness that counts, for no doctor has time for a product which proves merely decorative. The popularity of the now improved injectable forms of Penidural is due to the dramatic, long-lasting effects that can be achieved, with intriguing simplicity, in treatment and prophylaxis, often using a single dose of these unique preparations.

PENIDURAL BY INJECTION



LONG-ACTING


200,000 units of benzathine penicillin per c.c. Packs of single 4 c.c. and 10 c.c. vials.

ALL-PURPOSE

600.000 units of benzathine penicillin fortified with 300.000 units of each of potassium and procaine penicillin G. Packs of single 1.2 megaunit vials.


The word 'Penidural' is a registered trade mark of


l·d.2




LEWIS LABORATORIES LTD., LEEDS 9

5073/E







 $\mathbf{I}^{\scriptscriptstyle\mathsf{T}}$ is generally agreed that cases of soft-tissue rheumatism and of many types of arthritis benefit by hyperaemia of the affected tissues. Transvasin, which contains two quick-acting esters of nicotinic acid, induces such a deep and prolonged hyperaemia at the same time as it produces superficial erythema.

In addition to the esters of nicotinic acid, Transvasin contains the fat- and water-soluble esters of salicylic and p-aminobenzoic acids. Both these well-tried analgesic drugs are readily conveyed through the skin in therapeutic quantities and enable an effective concentration to be built up where they are needed.

Transvasin is now being widely prescribed, with successful clinical results. Since a very small quantity is sufficient for each application, the cost of treatment is extremely low.

COMPOSITION


Salicylic acid tetrahydrofurfuryl-ester		14%
Nicotinic acid ethyl-ester		2%
Nicotinic acid n-hexyl-ester	••	2%
p-Aminobenzoic acid ethyl-ester		2%
Water-miscible base ad	••	100%

Transvasin is available in 1 oz. tubes, basic N.H.S. price of 2/6+Purchase Tax, and is not advertised to the public. Samples and literature will be gladly sent on application.



LLOYD-HAMOL LTD., 11 WATERLOO PLACE, LONDON, S.W.1. WHItehall 8654/5/6

'Transvasin' is a registered trade mark of Lloyd-Hamol Limited



"The common association of an anaemia with rheumatoid disease has been recognised for many years . . ."

Iron?—"Blood transfusions have been given with good results but improvement has been transient".

Oral iron?—"The prolonged administration of iron by mouth has not proved very effective".

Intravenous iron?—"... the difficulty of giving such injections to patients with joint deformity is a drawback as well as the liability to cause unpleasant side reactions".

Intramuscular iron?—"Clearly, an iron preparation for intramuscular injection free from side effects and fairly rapidly absorbed would be a great advantage It is concluded that . . . in the treatment of the anaemia of rheumatoid arthritis . . . intramuscular iron is as beneficial as intravenous iron and easier to administer." Ann. Rheum. Dis., 1956, 15, 51.

Conclusion:

A fall in haemoglobin is a common feature in arthritis and its restoration to normal is an essential part of the treatment. Restoration of a normal haemoglobin level can be quickly and surely achieved with Imferon.





Relief of Rheumatic Pain in General Practice

1. Formulation of succinate overcomes toxicity of massive dosage of aspirin

Pain caused by all forms of rheumatism can now be relieved promptly by massive aspirin dosage without fear of toxicity. The Berex formulation of aspirin (3.7 gr.) and calcium succinate (2.8 gr.) has overcome this old problem so effectively that your patients can safely administer the drug themselves. Repeated visits to the surgery for supervision of administration are unnecessary.

2. Side-effects fewer and milder

Even after prolonged dosage, Berex produces no toxic effects. The prothrombin level is maintained and there is no hæmorrhagic tendency.* Prolonged periods of relief can be enjoyed in safety. The patient is simply instructed to reduce the initial high dosage as soon as the pain diminishes. Side effects, if present at all, are fewer and milder.

3. Encourages tissue respiration

Experimentally, by Warburg test, it has been shown that the inhibitory action of salicylate on tissue respiration is completely offset when it is combined with succinate. Succinate encourages cellular respiration and stimulates the respiratory enzyme systems. It is to this stimulating action that the beneficial effect of succinate is attributed.

* "No abnormal prolongation of prothrombin time even after 68 days of succinate-salicylate." "The results also show that this succinate-salicylate formulation combines safety and efficacy, permitting wide use both for treatment and maintenance without the excessive supervision required in many other forms of therapy." Delaware State Med. J., 1954, 26, 22.



FOR PROMPT RELIEF OF PAIN

ASSOCIATED WITH

ALL FORMS OF RHEUMATISM



PROMPT RELIEF or all rheumatic pains by massive aspirin dosage is now made safe by the Berex succinate-aspirin formulation. Prolonged dosage does not lower the prothrombin level, and does not therefore produce hamorrhagic tendency.

FORMULA:

Calcium succinate 2.8 gr., acetylsalicylic acid 3.7 gr.


IN TABLET FORM:

basic N.H.S. price, 4 8½d.—100 tablets; 24 3d.—600 tablets.

BEREX HAS NEVER BEEN ADVERTISED TO THE PUBLIC

A professional sample will be sent on request to:

MEDICAL DEPARTMENT: CLINOD PHARMACEUTICALS LID BELVUE ROAD NORTHOLT, GREENFORD, MIDDX



Butazolidin's outstanding reputation in the treatment of rheumatic disorders should not overshadow its value in rapidly relieving pain and swelling and in restoring normal temperature in superficial thrombophlebitis.

It is rarely necessary for treatment to last more than one week.




Geigy

Detailed literature is available on request.

Geigy Pharmaceutical Company Ltd., Wythenshawe, Manchester, 23





"You mean Algesal?"

"Yes - local salicylate by inunction and all that"

"Very well. Matter of fact, I was meaning to tell you about it — it is extraordinarily effective for soft tissue rheumatism and it works without irritating the skin."

"Seems to bear out what that fellow said the other day at the Medico-Chi meeting, that the site of action of the salicylates in the rheumatic syndromes may be peripheral, at tissue level."

Algesal

10% diethylamine salicylate in a soothing vanishing cream

non-irritant, skin penetrating salicylate for the relief of rheumatic pain by inunction

Full literature and samples gladly sent on request to the manufacturers: E. G. H. LABORATORIES LTD, PERU STREET, SALFORD 3, LANCS.

Aspirin intolerance in rheumatoid arthritis

BUFFERIN—the new, better-tolerated analgesic

Most successful answer yet discovered The latest report by the Joint Council Committee of the Medical Research Council and the Nuffield Foundation (on clinical trials of cortisone, ACTH, and other therapeutic measures in chronic rheumatic diseases) confirms the findings of the previous year. "At the end of two years, therefore, as previously reported at the end of one year, it appears that for practical purposes there is remarkably little to choose between cortisone and aspirin in the management of this group of patients." (British Medical Journal, 1955, 2, 695.)

As aspirin is a far cheaper and less dangerous therapeutic agent it is obvious why practitioners prefer it to cortisone. But there is still the problem of aspirin intolerance which affects as many as 42% of arthritic patients. And Bufferin, the new analgesic which reduces gastric upset to a negligible minimum, is certainly the most successful answer yet discovered. (J.A.M.A., June 4, 1955, p. 387.)

Far better tolerated — even by arthritics Bufferin combines acetylsalicylic acid with the antacids aluminium glycinate and magnesium carbonate (gentle antacid agents which do not produce "acid rebound"), and is well tolerated — even by arthritics. In blind trials amongst arthritics with proven intolerance to ordinary aspirin, 70% had no gastric symptoms after taking large doses of Bufferin over periods of 4 to 16 months. (J.A.M.A., June 4, 1955, p. 387.)

Acts twice as quickly as aspirin Not only is Bufferin better tolerated by the stomach than ordinary aspirin, but its pain relieving ingredient is absorbed twice as quickly into the blood-stream. Trials show that on an average the 10-minute salicylate level after taking Bufferin was more than 20% higher than the 20-minute level after taking ordinary aspirin (J. Amer. Pharm. Assoc. 1950, 39:21.)

BUFFERIN HAS NO EQUIVALENT IN THE B.P. OR NATIONAL FORMULARY.
THIS FULLY JUSTIFIES THE PRESCRIPTION OF BUFFERIN ON E.C.10.

BUFFERING ANALGESIC

A detailed, fully documented leaflet and sample are available on request to the makers, Bristol-Myers Co. Ltd., Dept. B, 211 Blackfriars Road, London, S.E.1.

Formula: Acetylsalicylic Acid 5 gr.: Aluminium Glycinate \(\frac{3}{4} \) gr.: Magnesium Carbonate I\(\frac{1}{2} \) gr.

For OVER-ACIDITY prescribe

VICHYCELESTINS

WORLD-FAMOUS FRENCH SPA WATER

Bottled as it flows from the Spring

Holding an undisputed place in the therapeutics of rheumatism and arthritis, as well as in disorders of the digestive and urinary tract, Vichy-Celestins is invaluable to sufferers from these ailments.

Sole Agents in the United Kingdom:

INGRAM & ROYLE LIMITED

50 Manchester Street, London, W.1







CAIRN HYDRO HOTEL

HARROGATE

The only Hotel in Harrogate with a full suite of medical baths under expert attention. A comfortable friendly place with good food and good service and personal attention. Spacious bedrooms, with gas fires or radiators, some with private bathrooms. Pleasant gardens.

A few minutes' walk from the Royal Baths and Pump Room.

TELEPHONE: HARROGATE 4005/8

TRUST HOUSES LIMITED

