

21

**GENERATION AND CHARACTERISATION OF MONOCLONAL ANTIBODIES FROM SINGLE RA SYNOVIAL B CELLS**

Khaled Amara,<sup>1</sup> Eric Meffre,<sup>2</sup> Lena Israelsson,<sup>1</sup> Monika Hansson,<sup>1</sup> Omri Snir,<sup>1</sup> Johanna Steen,<sup>1</sup> Fiona Murray,<sup>1</sup> Hedda Wardemann,<sup>3</sup> Lars Klareskog,<sup>1</sup> Vivianne Malmström<sup>1</sup> <sup>1</sup>Rheumatology Unit, Karolinska Institutet, Center for Molecular Medicine, Stockholm, Sweden; <sup>2</sup>Department of Immunobiology, Yale University School of Medicine, New Haven, Connecticut, USA; <sup>3</sup>Max Planck Research Group, Molecular Immunology, Max-Planck-Institute for Infection Biology, Berlin, Germany

10.1136/annrheumdis-2011-201234.21

**Background and objectives** Anticitrulline protein antibodies (ACPA) are a hallmark of HLA-associated rheumatoid arthritis (RA), but it is still not established whether these antibodies represent a cause or a consequence of arthritis. It is however clear that such antibodies can enhance murine arthritis, but a corresponding effect in patients is difficult to prove. In this study the authors aimed to assess the specificity

and the immunoglobulin gene characteristics of B cells derived from RA synovial fluid of RA by using a method that allows in vitro production of monoclonal antibodies derived from single human memory B cells.

**Materials and methods** The authors have cloned and expressed antibodies from single flow cytometry purified CD19+IgG+ B cells from synovial fluid of two ACPA+ and two ACPA- RA patients and tested, by ELISA, the generated recombinant monoclonal antibodies for reactivity to different known citrullinated antigens.

**Results** On the molecular level, striking differences were found between ACPA+ and ACPA- patients taking into consideration the mutational pattern of the Ig genes. Indeed, based on DNA sequences, the authors could demonstrate that B cells from ACPA+ patients did not have more total mutations than ACPA- patients but when focusing on the CDR1, two and three regions of the Ig variable region, the ACPA+ clones displayed more replacement mutations than did the ACPA- clones.

Regarding the specificity of the generated monoclonal antibodies, the authors have analysed clones from one ACPA+ and one ACPA negative patient. Our results so far show that 10 out of 23 antibodies generated from an ACPA+ RA patient, proved to bind, to variable degrees, to cyclic citrullinated peptide and different citrullinated antigens including citrullinated a-enolase, fibrinogen, vimentin and type II collagen. In contrast, no ACPA positive clones were identified amongst the antibodies derived from the ACPA- RA patient.

**Conclusions** Our data suggest that citrulline-reactive B cells are common in synovial fluid and the Ig molecules bear clear signs of antigen-driven maturation.