



OPEN ACCESS

EXTENDED REPORT

Safety, tolerability and potential efficacy of injection of autologous adipose-derived stromal vascular fraction in the fingers of patients with systemic sclerosis: an open-label phase I trial

Brigitte Granel,^{1,2} Aurélie Dumas,¹ Elisabeth Jouve,³ Jean-Robert Harlé,¹ Pierre-Sébastien Nguyen,⁴ Christian Chabannon,⁵ Nathalie Colavolpe,⁶ Jean-Charles Reynier,⁷ Romain Truillet,³ Stéphanie Mallet,⁸ Antoine Baiada,⁴ Dominique Casanova,⁴ Laurent Giraudo,⁹ Laurent Arnaud,¹⁰ Julie Veran,⁹ Florence Sabatier,^{2,9,10} Guy Magalon^{4,9}

Handling editor Tore K Kvien

► Additional material is published online only. To view please visit the journal online (<http://dx.doi.org/10.1136/annrheumdis-2014-205681>).

For numbered affiliations see end of article.

Correspondence to

Professor Brigitte Granel, North Hospital, Assistance Publique Hôpitaux de Marseille (AP-HM), Marseille 139015, France; bgranel@ap-hm.fr

Received 4 April 2014

Revised 15 July 2014

Accepted 20 July 2014

Published Online First

11 August 2014



Open Access
Scan to access more
free content



CrossMark

To cite: Granel B, Dumas A, Jouve E, et al. *Ann Rheum Dis* 2015;**74**:2175–2182.

ABSTRACT

Background In patients with systemic sclerosis (scleroderma, SSs), impaired hand function greatly contributes to disability and reduced quality of life, and is insufficiently relieved by currently available therapies. Adipose tissue-derived stromal vascular fraction (SVF) is increasingly recognised as an easily accessible source of regenerative cells with therapeutic potential in ischaemic or autoimmune diseases. We aimed to measure for the first time the safety, tolerability and potential efficacy of autologous SVF cells local injections in patients with SSs with hand disability.

Methods We did an open-label, single arm, at one study site with 6-month follow-up among 12 female SSs patients with Cochin Hand Function Scale score >20/90. Autologous SVF was obtained from lipoaspirates, using an automated processing system, and subsequently injected into the subcutaneous tissue of each finger in contact with neurovascular pedicles. Primary outcome was the number and the severity of adverse events related to SVF-based therapy. Secondary endpoints were changes in hand disability and fibrosis, vascular manifestations, pain and quality of life from baseline to 2 and 6 months after cell therapy.

Findings All enrolled patients had surgery, and there were no dropouts or patients lost to follow-up. No severe adverse events occurred during the procedure and follow-up. Four minor adverse events were reported and resolved spontaneously. A significant improvement in hand disability and pain, Raynaud's phenomenon, finger oedema and quality of life was observed.

Interpretation This study outlines the safety of the autologous SVF cells injection in the hands of patients with SSs. Preliminary assessments at 6 months suggest potential efficacy needing confirmation in a randomised placebo-controlled trial on a larger population.

Funding GFRS (Groupe Francophone de Recherche sur la Sclérodémie).

Clinical Trials number NCT01813279.

Systemic sclerosis (scleroderma, SSs) is a rare systemic autoimmune disease characterised by microvascular lesions and progressive fibrosis of skin and internal organs. Hands involvement is the most

classical and visible manifestation of the disease and is a large burden on global disability and social relationship. Hand disability has multifactorial origins with microvascular lesions, paroxysmal vasospasm or permanent ischaemia and subsequently digital ulcers (DU), skin sclerosis, tendons retractions, bone and articular involvement, and subcutaneous calcinosis.^{1–3} In patients with SSs, radiological abnormalities included erosion (21%), joint space narrowing (28%), arthritis (18%), acro-osteolysis (22%), flexion contracture (27%) and calcinosis (23%).⁴ All these lesions cause pain, functional impairment, aesthetic and psychological distress.¹ To date, therapeutic interventions for SSs hands have mainly focused on vascular manifestations, including Raynaud's phenomenon and DUs.^{5–9} These treatments are often life-long therapies with considerable costs and potentially significant side effects. No antifibrotic drug is effective in SSs. Conventional immunosuppressive drugs effectively treat joint synovitis and sometimes even improve skin fibrosis but do not improve contractures of small joints.¹⁰

Most research priorities are aimed to prevent or treat life-threatening complications that arise from affection of internal organs; however, if asked, patients indicate that hand disability interfering with daily life and work matters most.¹¹ Thus, healthcare professionals should consider improving disability, patients' satisfaction and social comfort as clinically relevant objectives of therapy.

The delivery of autologous stem/progenitor cells into ischaemic tissue of patients is emerging as a novel therapeutic option for vascular diseases. To date, bone marrow-derived cells have mainly been used in such context^{12–14} but are limited by the invasive nature of bone marrow harvest, the delay and quality management issues associated with cells culture.¹⁵ In 2002, Zuk and colleagues¹⁶ characterised a multipotent population of cells within the stromal vascular fraction (SVF) of adipose tissue, referred to adipose-derived stromal/stem cells (ADSC). In addition to multipotency, this population exerts paracrine proangiogenic effects through the increased expression of vascular endothelial

growth factor, basic fibroblast growth factor and interleukin-6.¹⁷ Moreover, these cells have anti-inflammatory effects that appear to be mediated through secretion of anti-inflammatory soluble factors such as interleukin-10, interferon- γ , transforming growth factor- β or prostaglandin E2, with dendritic cells immunomodulation, stimulation of regulatory T cells and inhibition of inflammatory cytokines synthesis.¹⁸

Results from the recent ACellDREAM¹⁹ trial demonstrated the safety of autologous cultured ADSC in patients with critical limb ischaemia, with favourable secondary endpoints on efficacy. Recently, six patients with localised scleroderma (linear and plaque scleroderma, generalised morphea) have been locally treated with in vitro expanded autologous ADSC with a favourable safety profile and a significant improvement in the skin, at 1-year follow-up.²⁰

Thus, adipose tissue-derived SVF may have potential efficacy against ischaemia and skin fibrosis, two major manifestations of SSc. We tested in this early phase clinical trial the safety and tolerability of local injections of autologous SVF cells in fingers and assessed preliminary efficacy outcomes in patients with SSc with diminished hand function.

METHODS

Study design and eligibility criteria

Twelve patients with SSc were enrolled from December 2012 through May 2013 in an open-label, single-arm and single study site. All subjects fulfilled the 1980 American College of Rheumatology criteria²¹ or the 1988 LeRoy et al criteria.²² All enrolled patients had a score of ≥ 9 according to the 2013 Classification criteria for SSc.²³ Subjects had previously received optimal treatment for digital vasculopathy, according to EULAR.⁵

Patients were eligible if they were aged >18 and had a Cochin Hand Function Scale (CHFS)²⁴ $>20/90$. Exclusion criteria were new vasodilators or immunosuppressive therapy for SSc in the 3 months prior to enrolment or during the 6-month follow-up, surgical contraindication, clinical or radiological signs of digital infection, positive status for HIV, hepatitis B or C, HTLV1-2 or syphilis, pregnancy and body mass index <17 kg/m².

Patients were asked not to change their regular medications and physiotherapy of the hands during the trial. The study was approved by the local Ethics Committee and the French Drug Agency. The study was done in accordance with the Declaration of Helsinki and current Good Clinical Practices. An independent data monitoring committee ensured the integrity of the trial and safety of participants. All patients gave written informed consent before participation. This study is registered with ClinicalTrials.gov number NCT01813279.

Procedures

Tissue collection and SVF preparation

Adipose tissue collection and SVF injection were conducted under conscious sedation; harvesting areas were anaesthetised. Harvesting was performed with a 10 mL syringe in a closed circuit using a 3 mm Coleman cannula with 2 terminal 4 mm \times 2 mm openings with a two-way non-return AT valve, sterile tubing and a 250 mL collection bag. Once harvesting was complete, the bag was immediately transported to the registered Cell Therapy Unit. SVF was obtained within 2 h after lipoaspiration using the automated processing Celution800/CRS system (Cytori Therapeutics, San Diego, California, USA). Collected lipoaspirate was washed and enzymatically digested using GMP-grade reagents. Cells were concentrated, washed,

aseptically recovered and resuspended in 5–11 mL Lactate Ringer's solution. The final cell suspension was transferred into 10 syringes (1 mL) for injection. The remaining volume was used for sterility testing and biological characterisation. Total viable nucleated cell recovery and viability percentage were determined using the Nucleocounter NC100 (ChemoMetec, Denmark). Cellular components were identified by flow cytometry analysis (Beckman Navios instrument) with a panel of cell surface makers in agreement with International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT) recommendations.²⁵ The following markers (CD45, CD34, CD90, CD146, CD14) were used in combination with DRAQ5 and DAPI to exclude debris, red blood cells and dead cells. The frequency of adipose-derived mesenchymal-like stem cells was estimated using the Colony-Forming-Unit-Fibroblastic clonogenic assay.²⁵

SVF delivery

Entry points were preformed using a 25-gauge needle (0.5 mm \times 16 mm). SVF cells were then injected using a 25-gauge (0.5 mm \times 40 mm) reinforced cannula placed into the subcutaneous tissue in contact with the neurovascular pedicles: 0.5 mL SVF was injected into each lateral side of each digit, using a retro-tracing technique, from distal to proximal. Entry points were positioned at the metacarpophalangeal joint for the thumb, and the proximal interphalangeal joint where the palmar and dorsal skin joins for long fingers. Both hands were treated over a period of approximately 20 min.

Assessment of safety, tolerability and efficacy

See online supplementary table S1.^{26–30}

Statistical analysis

See online supplementary text.

RESULTS

Characteristics of the patients and the SVF

All enrolled patients had surgery, and there were no dropouts or patients lost to follow-up. The population mainly included patients with limited cutaneous form of the disease and without severe organ damage (table 1). No patient was treated with >10 mg steroids daily, cyclophosphamide or mycophenolate mofetil. Hand radiology revealed abnormalities in 10/12 patients (see online supplementary table S2). No patient had radial or ulnar occlusion on arterial Doppler ultrasound.

Patients received a mean of $3.76 \pm 1.85 \times 10^6$ viable cells into each finger. The infused cells contained a mean of $3.7 \pm 1.9\%$ CFU-F. Flow cytometry characterisation of cell subsets in SVF is shown in table 1.

Safety and tolerance profile of autologous adipose-derived SVF injection

All patients were discharged from hospital 24 h after surgery. No serious AEs occurred during follow-up. Four minor AEs reported by four patients were potentially related to the procedure: two abdominal bruises induced by the lipoaspiration of respectively 7 and 15 days duration, one transient paresthesia on the lateral side of the left fifth finger persisting for 11 days postinjection and one pain located on the lateral side of the left thumb persisting for 13 days postinjection. These events spontaneously resolved. Abdominal lipoaspiration sample points healed in less than 7 days postsurgery and the points of entry for SVF injection healed as soon as D+1. Abdominal pain remained moderate and transient: visual analogue scale (VAS) of 32 ± 13 on D+1, 47 ± 18

Table 1 Characteristics of the 12 patients with systemic sclerosis (SSc) and adipose tissue harvest

Demographic	
Number of patients, sex	12, all female
Age (years)	54.5±10.3 (34.0–68.0)
Body mass index (kg/m ²)	22.0±2.1 (18.6–26.2)
Dominant hand	
Right-handed/left-handed	10 (83.3%)/2 (16.7%)
Main causes of hand disability	
1. Vascular aspects (Raynaud's phenomenon)	12 patients (100%)
2. Skin fibrosis (sclerodactyly)	11 patients (92%)
3. Musculoskeletal aspects (joint synovitis)	5 patients (42%)
Cardiovascular risk factors	
Tobacco	0 (0.0%)
Diabetes	0 (0.0%)
Arterial hypertension	0 (0.0%)
Disease history	
Disease duration from diagnosis (years)	9.9±7.0 (2.0–24.0)
Early SSc disease <4 years	3 (25%)
History of Raynaud's phenomenon (years)	14.3±7.7 (5.0–34.0)
Systemic sclerosis cutaneous subclassification: limited/diffuse	7 (58.3%)/5 (41.7%)
Pulmonary arterial hypertension or previous renal crisis	0 (0.0%)
Rodnan	
Global modified Rodnan skin score	13.9±9.8 (3–32)
Gastrointestinal	
Gastrointestinal reflux	9 (75%)
Hypotonia of the inferior oesophagus sphincter (manometry)	9/9 tested
Lung	
Total lung capacity <80%	0 (0%)
Lung capacity for carbon monoxide/alveolar ventilation <75%	5 (41.7%)
Fibrosis on high-resolution chest tomography (bronchiectasia and/or honeycombing)	2 (16.7%)
Serology	
Anti-nuclear antibodies positive (indirect immunofluorescence on HEp2)	12 (100%)
Anti-topoisomerase 1 (Scl70) antibodies positive (ELIA)	6 (50.0%)
Anti-centromere antibodies positive (ELIA)	4 (33.3%)
Anti-SSA antibodies positive (ELIA)	2 (16.7%)
Ongoing and previous systemic sclerosis medications	
Calcium-channel blockers	6 ongoing (50.0%), 11 previously (91.7%)
Bosentan	2 ongoing (16.6%), 4 previously (33.3%)
Prednisone <10 mg/day	3 ongoing (25.0%)
Methotrexate	1 ongoing (8.3%)
Iloprost infusion	0 ongoing, 8 previously (66.7%)
Adipose tissue harvest and cell characterisation	
Amount of adipose tissue harvest (mL)	181.3±50.8 (120–270)
Number of total viable nucleated cells (×10 ⁶)	50.5±23.8 (16.7–92.6)
% of cell viability	88.0±4.8 (81.0–94.6)
Number of viable nucleated cells injected into each finger (×10 ⁶)	3.76±1.85 (1.19–7.07)
% of CFU-F	3.7±1.9 (1.6–8.1)
% of blood leucocytes (CD45 CD34– cells)	49.1±18.0
% of endothelial progenitor cells (CD34brightCD146dimCD45–)	3.4±3.2
% of circulating endothelial cells (CD34dimCD146brightCD45–)	6.4±6.8
% of stromal cells (CD45-CD34brightCD146-CD90+)	36.0±14.5
% of resident macrophages (CD45+CD14+CD34dim)	5.1±2.3


Data are mean±SD (minimum and maximum) or n (%) of patients.

on D+7 and 13±17 on D+21, and resolved completely at M2. Hand pain did not worsen. SVF injection led to a rapid decrease of the Raynaud's Condition Score (RCS) from 7.2±0.9 at baseline to 5.3±1.7 at D+7 and to 4.6±3.0 at D+21. The number of patients with normal finger colour increased from 5 at baseline to 8 at D+7 and D+21, without colour score worsening.

Efficacy profile of autologous adipose-derived SVF injection Effects on quality of life

Patients perceived their general health status to be improved as shown by a 45.2% and 42.4% decrease of the Scleroderma Health Assessment Questionnaire (SHAQ) at M2 (p=0.001) and at M6 (p=0.001), respectively (figure 1).

Figure 1 Six-month follow-up for patients with systemic sclerosis receiving stromal vascular fraction injection ($p < 0.05$ at M2 and M6).



Effects on hand disability and mobility

A 47.4% and 56.0% decrease of the CHFS at M2 and M6 in comparison to baseline was observed ($p < 0.001$ for both). Grip strength increased at M6 with a mean improvement of $+48 \pm 64$ kg for the dominant hand ($p = 0.033$) and $+40 \pm 3.5$ kg for the non-dominant hand ($p = 0.002$) (table 2, figure 1). Similarly, an increase in pinch strength at M6 was noted with a mean improvement of $+1.0 \pm 1.1$ kg for the dominant hand ($p = 0.009$) and $+0.8 \pm 1.2$ kg for the non-dominant hand ($p = 0.050$). The Kapandji score did not change. A mean increase in first corner distance of 13.3 ± 11.6 mm in dominant hand and 17.0 ± 14.7 mm in non-dominant hand was noted at M6 ($p = 0.002$ for both). Similarly, an increase of the sum of 2nd, 3rd and 4th corner distances was noted at M6 with a mean improvement of 8.3 ± 5.1 mm in non-dominant hand ($p < 0.001$) and of 9.8 ± 15.3 mm in dominant hand, which however did not reach statistical significance ($p = 0.073$). Finger flexion also improved but the difference did not reach statistical significance.

Effect on fingers' oedema, skin fibrosis and digital ulcer

SVF injection induced a significant reduction in the circumference of the fingers at M2 and M6 in both hands. Although four patients presented at least 25% reduction of MRSS focused on hands at M6, change in the whole population did not reach statistical significance (table 3). Global MRSS decreased from baseline to M2 ($p = 0.010$) and to M6 ($p = 0.013$). The number of patients with normal finger colour increased from 5 at baseline to 9 at M2 and to 10 at M6; no patient worsened. Among subjects having at least one DU at inclusion, total number of DU decreased, from 15 DUs at baseline, 10 at M2 and 7 at M6 (see online supplementary table S3 for the details).

Effect on Raynaud's severity, hand pain and capillaroscopic changes

Average reduction of the RCS from baseline was 53.7% at M2 ($p < 0.001$) and 67.5% at M6 ($p < 0.001$) (table 4, figure 1). Hand pain showed a significant decrease of 37.8% at M2 ($p = 0.001$) and 41.7% at M6 ($p < 0.001$). Also, 9 of the 12 patients with SSc had at least 25% improvement in all the following four parameters: CHFS, RCS, VAS pain and SHAQ.

Capillaroscopy evaluation showed no significant change in the number of nailfold capillary loops from baseline to M6. A slight decrease in the number of giant capillaries was noted at M6 for the dominant hand ($p = 0.035$), while the decrease did not reach statistical significance for the non-dominant hand ($p = 0.074$). A significant decrease in dystrophic capillaries was observed from

baseline to M6 in both hands ($p = 0.047$ for both). Finally, a significant decrease in the vascular suppression score was observed from baseline to M6 ($p = 0.01$ for the dominant hand and $p = 0.003$ for the non-dominant hand).

DISCUSSION

This study demonstrates that injection of autologous SVF from adipose tissue into the fingers of patients with SSc can be performed safely and is well tolerated. The only safety-related events were minor bruising, consistent with what is normally seen following lipoaspiration, and transient finger discomfort. Importantly, a priori efficacy outcomes variables revealed a decrease of ~50% in CHFS, Raynaud's severity and hand pain at 2 months with persisting benefits at 6 months. To our knowledge, SVF is the first treatment improving such hand manifestations and with favourable impact on quality of life in patients with SSc.

Adipose-derived SVF yields a heterogeneous population of cells including stem and progenitor cells with multipotent differentiation potential. SVF cells transcribe many genes that are implicated inflammation, angiogenesis and tissue repair.³¹ It is suggested that ADSC can have antifibrotic properties by the reduction of local infiltration of inflammatory cells into tissue by the secretion of antifibrotic factors such as interferon- γ and matrix metalloproteinases,³¹ and by the decrease of pro-fibrotic factors such as transforming growth factor- β .³² Advantages of SVF include (1) ease of obtaining cells from lipoaspirates, (2) larger pool of ADSC compared with the pool of bone marrow-derived mesenchymal stem cells (BM-MSC) and (3) stronger angiogenic and regenerative potential of ADSC compared with BM-MSC.³³ As a result, the use of SVF cells for tissue regeneration or autoimmune diseases is increasing as shown by numerous ongoing clinical trials (see Clinicaltrial.gov).

The present study mainly enrolled patients with SSc free of severe visceral manifestations but with long disease duration. Some of them had hand flexion contracture, joint space narrowing, severe vascular disease, calcinosis and acro osteolysis. These characteristics may explain the higher frequency of joint involvement and radiographic abnormalities compared with that previously reported by Avouac *et al.*⁴ Major hand disability in our cohort was also consistent with a CHFS score much higher than that reported by Rannou *et al.*²⁴ Despite such difference, we similarly noticed a strong correlation between CHFS and SHAQ, with the CHFS scores contributing to 71.3% of the SHAQ variance at M6. This result highlights the crucial contribution of hand disability in global patient disability. Importantly, treatments and physiotherapy schedule were not changed for

Table 2 Effect of autologous adipose-derived stromal vascular fraction injection on hand disability

	Baseline	2 months	Change (2 months–baseline)	p Value*	6 months	Change (6 months–baseline)	p Value†
CHFS total							
Mean±SD	48.5±10.8	25.8±17.0	–22.7±16.4	<0.001	21.2±15.4	–27.3±17.2	<0.001
Median (range)	48.5 (30–69)	25.5 (2–49)	–22.5 (–51–10)		20.0 (0–48)	–30.0 (–58–10)	
Jamar score (kg)							
Dominant hand							
Mean±SD	16.0±5.8	17.9±7.3	3.0±5.0	0.072	19.4±7.4	4.8±6.4	0.033
Median (range)	15.0 (9.0–26.5)	18.5 (5.5–28.0)	2.0 (–3.0–15.0)		20.0 (5.0–30.0)	3.0 (–6.0–17.0)	
Non-dominant hand							
Mean±SD	14.9±6.1	17.5±7.8	3.7±3.6	0.006	17.6±8.0	4.0±3.5	0.002
Median (range)	14.0 (6.0–26.0)	18.5 (4.0–28.0)	3.0 (–2.0–9.0)		20.0 (3.5–29.0)	3.0 (0.0–13.0)	
Pinch score (kg)							
Dominant hand							
Mean±SD	1.3±1.1	2.0±0.8	0.6±1.3	0.199	2.3±1.3	1.0±1.1	0.009
Median (range)	0.9 (0.2–4.1)	2.3 (0.5–3.2)	0.8 (–2.3–2.7)		2.0 (0.9–5.4)	0.9 (–1.1–3.4)	
Non-dominant hand							
Mean±SD	1.3±0.9	2.2±0.7	0.7±1.2	0.071	2.1±1.0	0.8±1.2	0.050
Median (range)	0.9 (0.2–3.2)	2.3 (1.4–3.2)	0.5 (–0.9–2.5)		2.0 (0.7–3.6)	0.5 (–1.1–3.4)	
Kapandji score/10							
Dominant hand							
Mean±SD	8.0±1.4	8.3±1.6	0.3±1.0	0.380	8.4±1.7	0.5±0.9	0.111
Median (range)	8.0 (5.0–10.0)	9.0 (5.0–10.0)	0.0 (–1.0–2.5)		9.3 (5.0–10.0)	0.0 (–1.0–2.5)	
Non-dominant hand							
Mean±SD	8.5±1.2	8.4±1.6	–0.1±0.7	1.000	8.8±1.3	0.4±0.7	0.188
Median (range)	9.0 (6.0–10.0)	9.0 (5.0–10.0)	0.0 (–1.0–1.5)		9.0 (6.0–10.0)	0.0 (–0.5–2.0)	
1st corner distance (mm)							
Dominant hand							
Mean±SD	105.6±24.7	112.9±29.2	7.3±8.2	0.010	118.9±31.2	13.3±11.6	0.002
Median (range)	112.0 (57–142)	118.5 (57–154)	5.5 (–6–23)		124.0 (58–178)	10.0 (–3–36)	
Non-dominant hand							
Mean±SD	115.8±24.5	122.3±20.9	6.5±9.8	0.042	132.8±28.0	17.0±14.7	0.002
Median (range)	118.5 (65–152)	121.5 (88–155)	4.5 (–8–23)		137.0 (95–190)	15.5 (–4–38)	
Sum of corners distances (mm)							
Dominant hand							
Mean±SD	133.9±18.5	131.2±20.7	1.0±13.7	0.822	140.2±26.6	9.8±15.3	0.073
Median (range)	130.5 (110–168)	131.0 (94–169)	2.0 (–18–28)		131.0 (105–188)	5.5 (–7–36)	
Non-dominant hand							
Mean±SD	132.1±24.6	133.7±29.4	4.5±6.8	0.052	137.9±27.0	8.3±5.1	<0.001
Median (range)	139.0 (73–158)	139.5 (64–166)	6.0 (–9–13)		147.0 (77–177)	10.0 (1–19)	
Sum of Pad/DPL distance (mm)							
Dominant hand							
Mean±SD	52.0±46.5	47.3±43.8	–4.8±18.6	0.395	43.9±42.7	–8.1±16.2	0.111
Median (range)	49.5 (0–160)	45.0 (0–115)	–2.5 (–45–23)		36.5 (0–125)	–4.0 (–35–19)	
Non-dominant hand							
Mean±SD	48.1±54.5	46.8±52.0	–1.3±19.6	0.830	37.8±43.0	–10.3±20.6	0.109
Median (range)	32.0 (0–144)	38.5 (0–160)	–0.5 (–43–28)		15.0 (0–111)	–2.0 (–41–22)	

Bold values correspond to the p value columns. p≤0.05 (2-sided) was considered significant.

Data are mean±SD, median (minimum–maximum).

*p Value from baseline to M2.

†p Value from baseline to M6.

CHFS, Cochin Hand Functional Scale; DPL, Distal Palmar Line.

3 months before entry and all during the trial. Thus, SVF injection was the sole therapeutic change within the study period, allowing evaluation of safety, tolerability and therapeutic effects.

SVF treatment was associated with an ~50% improvement of CHFS, RCS, hand pain and global disability at 2 and 6 months. Although these scales are clearly subjective evaluations, they remain valid instruments when looking at change within individuals. These endpoints were complemented by improvement of objective criteria such as grip and finger's strength, corners

distances and fingers' circumference, and nailfold capillaroscopy data. When analyses were performed taking into account the cutaneous form of the disease, both patients with diffuse and limited forms benefited from the SVF injections. Kapandji and pad to distal palmar line distances did not improve during the trial but were only faintly altered at start, with little room for improvement. Altogether, these changes suggest that autologous SVF injection can counterbalance the various physiopathogenic factors involved in SSc disease. In the literature, no such large

Table 3 Effect of autologous adipose-derived stromal vascular fraction injection on scleroderma and skin thickness

	Baseline	2 months	Change (2 months–baseline)	p Value*	6 months	Change (6 months—baseline)	p Value†
<i>Mean circumference of F1–F5 (ring size)</i>							
Dominant hand							
Mean±SD	61.9±2.2	60.7±2.3	−1.2±1.4	0.013	59.8±2.4	−2.1±1.1	<0.001
Median (range)	61.8 (58.8–66)	60.8 (57–65)	−0.9 (−4.6–0.6)		60 (56–64.4)	−2.3 (−4.6—0.4)	
Non-dominant hand							
Mean±SD	60.7±2.3	59.3±1.7	−1.3±1.4	0.008	58.1±2.2	−2.5±1.5	<0.001
Median (range)	61 (57–64.2)	59 (56.8–63.4)	−1.2 (−3.4–0.8)		58 (54–62.8)	−2.5 (−4.8–0.4)	
<i>MRSS applied to hand/18</i>							
Whole population							
Mean±SD	10.9±4.9	10.0±5.3	−0.9±1.6	0.067	9.9±6.0	−1.0±2.8	0.246
Median (range)	11.5 (3–18)	10.5 (2–18)	−0.5 (−4–1)		12 (1–18)	−2 (−5–4)	
<i>Global MRSS/51</i>							
Whole population							
Mean±SD	13.9±9.8	11.7±9.8	−2.3±2.5	0.010	11.5±10.1	−2.4±2.8	0.013
Median (range)	12 (3–32)	9 (2–29)	−2 (−6–3)		8.5 (1–29)	−2 (−7–2)	

Bold values correspond to the p value columns. $p \leq 0.05$ (2-sided) was considered significant.

Data are mean±SD.

*p Value from baseline to M2.

†p Value from baseline to M6.

Data are mean±SD.

F1–F5, finger 1–finger 5; MRSS, Modified Rodnan skin score.

Table 4 Effect of autologous adipose-derived stromal vascular fraction injection on vascular manifestations

	Baseline	2 months	Change (2 months–baseline)	p Value*	6 months	Change (6 months–baseline)	p Value†
RCS/10							
Mean±SD	7.2±0.9	4.5±1.9	−2.7±2.2	<0.001	2.9±1.4	−4.3±2.1	<0.001
Median (range)	7.5 (6.5–8.0)	5.0 (4.0–5.5)	−2.0 (−6.5–1.0)		3.0 (2.5–3.0)	−4.5 (−7.0–1.5)	
Hand visual analogue scale/100							
Mean±SD	59.4±17.2	21.6±17.5	−37.8±28.6	0.001	17.8±15.3	−41.7±22.7	<0.001
Median (range)	58.5 (50.0–72.5)	19.0 (7.0–40.0)	−39.0 (−80.0–21.0)		13.6 (9.0–26.0)	−44.0 (−80.0–10.0)	
Capillaroscopy data							
Number of capillary loops							
Dominant hand							
Mean±SD	136.3±78.9	124.1±60.1	−12.2±40.9	0.325	138.5±60.8	2.3±47.9	0.874
Median (range)	124.0 (42–298)	118.0 (30–226)	−12.5 (−98–61)		132.5 (48–232)	8.0 (−97–92)	
Non-dominant hand							
Mean±SD	159.5±96.9	142.1±71.0	−23.5±54.1	0.181	152.3±75.2	−11.7±42.0	0.377
Median (range)	122.0 (45–298)	139.0 (42–277)	−13.0 (−137–33)		131.5 (70–308)	−10.0 (−103–34)	
Number of giant capillaries							
Dominant hand							
Mean±SD	20.3±20.3	17.8±18.0	−2.4±8.4	0.341	15.3±17.9	−4.9±7.8	0.035
Median (range)	12.0 (0–64)	13.0 (1–67)	1.0 (−22–7)		9.0 (0–63)	−1.0 (−17–6)	
Non-dominant hand							
Mean±SD	18.3±11.7	18.2±20.5	−5.5±7.9	0.046	18.1±18.6	−5.1±8.9	0.074
Median (range)	13.0 (0–39)	12.5 (0–77)	−3.0 (−20–3)		13.0 (0–72)	−3.0 (−23–4)	
Number of dystrophic capillaries							
Dominant hand							
Mean±SD	23.3±26.8	13.0±11.6	−10.3±17.9	0.115	11.0±10.3	−12.3±20.2	0.047
Median (range)	14.0 (0–79)	14.0 (0–30)	−2.0 (−53–6)		8.5 (0–29)	−1.0 (−57–5)	
Non-dominant hand							
Mean±SD	23.2±22.8	11.2±9.5	−11.4±14.9	0.030	9.7±6.7	−13.5±19.8	0.047
Median (range)	17.0 (0–58)	8.0 (0–29)	−4.0 (−37–6)		10.5 (0–22)	−6.0 (−45–15)	
Vascular suppression score							
Dominant hand							
Mean±SD	1.7±0.8	1.6±0.6	−0.1±0.4	0.656	1.5±0.7	−0.2±0.3	0.010
Median (range)	1.9 (1.0–2.3)	1.5 (1.1–2.1)	0.0 (−1.0–0.4)		1.4 (0.9–2.0)	−0.2 (−0.4–0.0)	
Non-dominant hand							
Mean±SD	1.6±0.7	1.5±0.7	−0.2±0.2	0.031	1.3±0.7	−0.3±0.3	0.003
Median (range)	1.8 (0.9–2.2)	1.6 (0.8–2.0)	−0.1 (−0.6–0.0)		1.1 (0.8–2.0)	−0.4 (−0.8–0.2)	

Bold values correspond to the p value columns. $p \leq 0.05$ (2-sided) was considered significant.

Data are mean±SD, median (minimum–maximum).

*p Value from baseline to M2.

†p Value from baseline to M6.

Dystrophic capillaries are enlarged loops over four times the normal afferent, transition and efferent limbs width. Giant capillary loops are 10 or more times the normal width of capillary limbs. The avascular score was assessed according to Lee's method¹⁸, in which each finger was rated from 0 to 3: grade 0=no avascular area; 1=one or two discrete avascular areas; 2=more than two discrete avascular areas; 3=extensive and confluent avascular areas. Vascular suppression score are the mean of fingers. RCS, Raynaud's Condition Score.

effect has been reported in patients with SSc: in the two randomised, prospective, placebo-controlled, double-blind studies (RAPIDS-1,⁶ RAPIDS-2⁷), bosentan showed its efficacy on prevention of DUs, but without any significant improvement of the HAQ score.

SVF injection predominantly improved vascular peripheral manifestations such as Raynaud's phenomenon severity, DUs outcome and hand pain, the latter being in major part related to chronic vasospasm and DUs. The decrease of fingers' circumference was probably related to an improvement of finger skin oedema, but we did not measure the hand volume to differentiate oedema from fibrosis. Altogether, these effects could explain the functional improvement observed in this study. These observations also suggest that SVF may improve vasomotor tone and microvascular perfusion. This hypothesis is further substantiated by the significant reduction of avascular areas and dystrophic capillaries evaluated using nailfold capillaroscopy. SVF mainly promotes vascular repair and angiogenesis as documented in various experimental models of tissue ischaemia.^{34–35} Moreover, defective endogenous capacity for vascular repair has been involved in SSc pathogenesis.³⁶ Thus, local supply of high amount of endothelial precursors, mesenchymal-like stem cells, pericytes and various angiogenic factors through SVF injection in damaged fingers can directly contribute to improve vascular deficiency. We found that the main characteristics of the Celution processed SVF from patients with SSc were comparable to previous reports³⁷ and to our personal observations from healthy subjects (data not shown). Consistently, BM-MS-C from patients with SSc were reported to exhibit similar phenotypic, proliferative, differentiation potential and immunosuppressive properties as their healthy counterparts.^{38–39} Given the heterogeneity of SVF and the as-yet unclear mechanisms, it would be difficult to determine whether the properties critical for efficacy are impaired in SVF from patients with SSc compared with that of healthy donors.

No significant correlation was observed between the characteristics of the injected SVF and clinical outcomes. Further study in a larger cohort may help delineate the contribution of each cell subpopulation to the clinical effect.

Although this was a single-centre, phase I study in a limited number of patients and with relatively brief follow-up, it showed that this innovative treatment was not only safe and well tolerated in patients with SSc but also provided encouraging preliminary evidence of efficacy in numerous secondary endpoints. Longer-term, larger and controlled studies will be important to confirm whether this new form of cell therapy can improve the long-term prognosis of this debilitating disease. In addition, future studies may help understand the mechanisms supporting the clinical effects of SVF.

Author affiliations

¹Internal Medicine Department, Assistance Publique Hôpitaux de Marseille (AP-HM), Aix-Marseille University, Marseille, France

²Vascular Research Center Marseille, INSERM UMR-S-1076, Aix-Marseille University, Marseille, France

³CIC- CPCET, Assistance Publique Hôpitaux de Marseille (AP-HM), Aix-Marseille University, Marseille, France

⁴Plastic Surgery Department, Assistance Publique Hôpitaux de Marseille (AP-HM), Aix-Marseille University, Marseille, France

⁵Cell Therapy Facility & Inserm CBT 510, Institut Paoli Calmettes, Marseille, France

⁶Department of Radiology, Assistance Publique Hôpitaux de Marseille (AP-HM), Aix-Marseille University, Marseille, France

⁷Délégation de la Recherche et de l'Innovation, Assistance Publique Hôpitaux de Marseille (AP-HM), Marseille, France

⁸Department of Dermatology, Assistance Publique Hôpitaux de Marseille (AP-HM), Aix-Marseille University, Marseille, France

⁹Culture and Cell Therapy Laboratory, INSERM CBT 510, Assistance Publique Hôpitaux de Marseille (AP-HM), Aix-Marseille University, Marseille, France

¹⁰Hematology and Vascular Biology Laboratory, Assistance Publique Hôpitaux de Marseille (AP-HM), Aix-Marseille University, Marseille, France

Acknowledgements We would like to thank the French Scleroderma Research Group for financial support. We also thank the Fondation de l'Avenir, Estelle Charles-Baumel, who contributed to data management, Didier Samson, who anaesthetised the patients, and Joelle Micallef, for her advices in the redaction of the manuscript.

Contributors All authors contributed to review of this manuscript. BG, AD, EJ, FS, JV and GM designed the protocol. SM did the dermatological assessment, NC did the Doppler US and X-ray and both helped analysing the data. AB did the physiotherapist assessments. BG and AD did the recruitment of patients and the inpatient and outpatient care. GM, PSN and DC performed the surgical procedure. FS, JV, LG and LA were in charge of the SVF preparation and packaging of the SVF for injection. They also did the characterisation and the quality controls of the SVF. JCR monitored the trial. EJ and RT performed the statistical analyses, reviewed charts and data and helped in interpretation of the findings. JRH was at the head of the independent survey committee of the study. CC criticised and reviewed the manuscript.

Competing interests None.

Ethics approval Local Ethic Committee and French Drug Agency.

Provenance and peer review Not commissioned; externally peer reviewed.

Open Access This is an Open Access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially, and license their derivative works on different terms, provided the original work is properly cited and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>

REFERENCES

- Guillemin L, Hunsche E, Denton CP, *et al.*; DUO Registry Group. Functional impairment of systemic sclerosis patients with digital ulcerations: results from the DUO Registry. *Clin Exp Rheumatol* 2013;31:71–80.
- Hachulla E, Clerson P, Launay D, *et al.* Natural history of ischemic digital ulcers in systemic sclerosis: single-center retrospective longitudinal study. *J Rheumatol* 2007;34:2423–30.
- Sandqvist G, Eklund M. Hand Mobility in Scleroderma (HAMIS) test: the reliability of a novel hand function test. *Arthritis Care Res* 2000;13:369–74.
- Avouac J, Guérini H, Wipff J, *et al.* Radiological hand involvement in systemic sclerosis. *Ann Rheum Dis* 2006;65:1088–92.
- Kowal-Bielecka O, Landewé R, Avouac J, *et al.* EULAR recommendations for the treatment of systemic sclerosis: a report from the EULAR Scleroderma Trials and Research group (EUSTAR). *Ann Rheum Dis* 2009;68:620–8.
- Korn JH, Mayes M, Matucci-Cerinic M, *et al.* Digital ulcers in systemic sclerosis: prevention by treatment with bosentan, an oral endothelin receptor antagonist. *Arthritis Rheum* 2004;50:3985–93.
- Matucci-Cerinic M, Denton CP, Furst DE, *et al.* Bosentan treatment of digital ulcers related to systemic sclerosis: results from the RAPIDS-2 randomised, double-blind, placebo-controlled trial. *Ann Rheum Dis* 2011;70:32–8.
- Brueckner CS, Becker MO, Kroencke T, *et al.* Effect of sildenafil on digital ulcers in systemic sclerosis: analysis from a single centre pilot study. *Ann Rheum Dis* 2010;69:1475–8.
- Wigley FM, Seibold JR, Wise RA, *et al.* Intravenous iloprost treatment of Raynaud's phenomenon and ischemic ulcers secondary to systemic sclerosis. *J Rheumatol* 1992;19:1407–14.
- Pope JE, Bellamy N, Seibold JR, *et al.* A randomized, controlled trial of methotrexate versus placebo in early diffuse scleroderma. *Arthritis Rheum* 2001;44:1351–8.
- Jewett LR, Hudson M, Malcarne VL, *et al.* Sociodemographic and disease correlates of body image distress among patients with systemic sclerosis. *PLoS One* 2012;7:e33281.
- Traverse JH, Henry TD, Ellis SG, *et al.* Effect of intracoronary delivery of autologous bone marrow mononuclear cells 2 to 3 weeks following acute myocardial infarction on left ventricular function: the LateTIME randomized trial. *JAMA* 2011;306:2110–19.
- Matoba S, Tatsumi T, Murohara T, *et al.* Long-term clinical outcome after intramuscular implantation of bone marrow mononuclear cells (Therapeutic Angiogenesis by Cell Transplantation [TACT] trial) in patients with chronic limb ischemia. *Am Heart J* 2008;156:1010–18.
- Hare JM, Fishman JE, Gerstenblith G, *et al.* Comparison of allogeneic vs autologous bone marrow-derived mesenchymal stem cells delivered by transendocardial injection in patients with ischemic cardiomyopathy: the POSEIDON randomized trial. *JAMA* 2012;308:2369–79.

- 15 Lu D, Chen B, Liang Z, *et al.* Comparison of bone marrow mesenchymal stem cells with bone marrow-derived mononuclear cells for treatment of diabetic critical limb ischemia and foot ulcer: a double-blind, randomized, controlled trial. *Diabetes Res Clin Pract* 2011;92:26–36.
- 16 Zuk PA, Zhu M, Ashjian P, *et al.* Human adipose tissue is a source of multipotent stem cells. *Mol Biol Cell* 2002;13:4279–95.
- 17 Kapur SK, Katz AJ. Review of the adipose derived stem cell secretome. *Biochimie* 2013;95:2222–8.
- 18 Leto Barone AA, Khalifian S, Lee WP, *et al.* Immunomodulatory effects of adipose-derived stem cells: fact or fiction?. *Biomed Res Int* 2013;2013:383685.
- 19 Bura A, Planat-Benard V, Bourin P, *et al.* Phase I trial: the use of autologous cultured adipose-derived stroma/stem cells to treat patients with non-revascularizable critical limb ischemia. *Cytotherapy* 2014;16:245–57.
- 20 Scuderi N, Ceccarelli S, Onesti MG, *et al.* Human adipose-derived stromal cells for cell-based therapies in the treatment of systemic sclerosis. *Cell Transplant* 2013;22:779–95.
- 21 Subcommittee for Scleroderma Criteria of the American Rheumatism Association Diagnostic and Therapeutic Criteria Committee. Preliminary criteria for the classification of systemic sclerosis (scleroderma). *Arthritis Rheum* 1980;23:581–90.
- 22 LeRoy EC, Black C, Fleischmajer R, *et al.* Scleroderma (systemic sclerosis): classification, subsets and pathogenesis. *J Rheumatol* 1988;15:202–5.
- 23 Van den Hoogen F, Khanna D, Fransen J, *et al.* 2013 classification criteria for systemic sclerosis: an American College of Rheumatology/European League against Rheumatism collaborative initiative. *Arthritis Rheum* 2013;65:2737–47.
- 24 Rannou F, Poiraudou S, Berezne A, *et al.* Assessing disability and quality of life in systemic sclerosis: construct validities of the Cochin Hand Function Scale, Health Assessment Questionnaire (HAQ), Systemic Sclerosis HAQ, and Medical Outcomes Study 36-Item Short Form Health Survey. *Arthritis Rheum* 2007;57:94–102.
- 25 Bourin P, Bunnell BA, Casteilla L, *et al.* Stromal cells from the adipose tissue-derived stromal vascular fraction and culture expanded adipose tissue-derived stromal/stem cells: a joint statement of the International Federation for Adipose Therapeutics and Science (IFATS) and the International Society for Cellular Therapy (ISCT). *Cytotherapy* 2013;15:641–8.
- 26 Kapandji A. Clinical test of apposition and counter-apposition of the thumb. Cotation clinique de l'opposition et de la contre-opposition du pouce. *Ann Chir Main* 1986;5:67–73.
- 27 Mathiowetz V, Kashman N, Volland G, *et al.* Grip and pinch strength: normative data for adults. *Arch Phys Med Rehabil* 1985;66:69–74.
- 28 Lee P, Leung FY, Alderdice C, *et al.* Nailfold capillary microscopy in the connective tissue diseases: a semiquantitative assessment. *J Rheumatol* 1983;10:930–8.
- 29 Clements PJ, Lachenbruch PA, Seibold JR, *et al.* Skin thickness score in systemic sclerosis: an assessment of interobserver variability in 3 independent studies. *J Rheumatol* 1993;20:1892–6.
- 30 Georges C, Chassany O, Mouthon L, *et al.* Validation of French version of the Scleroderma Health Assessment Questionnaire (SSc HAQ). *Clin Rheumatol* 2005;24:3–10.
- 31 Katz AJ, Tholpady A, Tholpady SS, *et al.* Cell surface and transcriptional characterization of human adipose-derived adherent stromal (hADAS) cells. *Stem Cells* 2005;23:412–23.
- 32 Lee SH, Lee EJ, Lee SY, *et al.* The effect of adipose stem cell therapy on pulmonary fibrosis induced by repetitive intratracheal bleomycin in mice. *Exp Lung Res* 2014;40:117–25.
- 33 Ikegami Y, Yamashita K, Hayashi S, *et al.* Comparison of mesenchymal stem cells from adipose tissue and bone marrow for ischemic stroke therapy. *Cytotherapy* 2011;13:675–85.
- 34 Rehman J, Traktuev D, Li J, *et al.* Secretion of angiogenic and antiapoptotic factors by human adipose stromal cells. *Circulation* 2004;109:1292–8.
- 35 Nakagami H, Maeda K, Morishita R, *et al.* Novel autologous cell therapy in ischemic limb disease through growth factor secretion by cultured adipose tissue-derived stromal cells. *Arterioscler Thromb Vasc Biol* 2005;25:2542–7.
- 36 Matucci-Cerinic M, Kahaleh B, Wigley FM. Review: evidence that systemic sclerosis is a vascular disease. *Arthritis Rheum* 2013;65:1953–62.
- 37 Lin K, Matsubara Y, Masuda Y, *et al.* Characterization of adipose tissue-derived cells isolated with the Celution system. *Cytotherapy* 2008;10:417–26.
- 38 Larghero J, Farge D, Braccini A, *et al.* Phenotypical and functional characteristics of in vitro expanded bone marrow mesenchymal stem cells from patients with systemic sclerosis. *Ann Rheum Dis* 2008;67:443–9.
- 39 Cipriani P, Di Benedetto P, Liakouli V, *et al.* Mesenchymal stem cells (MSCs) from scleroderma patients (SSc) preserve their immunomodulatory properties although senescent and normally induce T regulatory cells (Tregs) with a functional phenotype: implications for cellular-based therapy. *Clin Exp Immunol* 2013;173:195–206.

Statistical analysis (see online supplementary text)

Safety analysis was descriptive and based on the reported treatment-emergent adverse events and other safety information. Adverse events (AEs) were coded according to an established and validated adverse reaction dictionary. (MedDRA version 16.1)

Efficacy endpoints were continuous or ordinal variables. Data are shown as mean \pm standard deviation, median and range [minimum-maximum]. Assessments at month 2 (M2) and 6 (M6) were compared from baseline using a single sample test on mean of 0. Normal distribution was examined by Shapiro Wilks test. Normally distributed were analysed by paired t test. Non-parametric data including Jamar score, Kapandji score, number of giant capillaries, number of dystrophic capillaries and vascular suppression score were analysed by Wilcoxon signed-rank test. A nominal $p \leq 0.05$ (2-sided) was considered significant.

As this was the first study with SVF in scleroderma with safety as the primary objective, efficacy endpoints were deemed exploratory and therefore the p-values were considered nominal. Results were not corrected for multiple comparisons. Statistical analyses were performed using SAS (version 9.2)

Table S1: Assessment of safety, tolerability and efficacy of the trial

Assessment of safety and tolerability *
<i>Tolerability was based on clinical examination at Day+1, Day+7, Day+21, and was assessed by the following items:</i>
Skin hands colouring at room temperature
Body temperature
Healing of tissue harvest and SVF injection sites
Visual analog scale (VAS 0-100) for hands and abdomen pain quantification
Raynaud's Condition Score (RCS) evaluating how much the condition affected use of the hands during the preceding week
<i>scale 0=no pain or no functional impact, to 10=maximum pain or functional impact</i>
DU(s) outcome (size, aspect, number...)
Assessment of efficacy *
<i>Efficacy endpoints were assessed by a single evaluator at baseline, 2 and 6 months following SVF injection</i>
1) Hand function:
Cochin Hand Function Scale(CHFS) questionnaire [24]
<i>The questionnaire was completed in the light of the patient's answers to 18 questions concerning daily living activities, each question being scored from 0=performed without difficulty, to 5=impossible to do. Disability was recorded as the total score (range 0–90)</i>
Kapandji Score assessing opposition of the thumb
<i>scale 0=impossible, to 10 =complete [26]</i>
Grip and pinch strength using a Jamar dynamometer under standard conditions [27]
Lateral range of motion of the fingers by measuring the distance between the thumb and index finger (1 st corner) and the sum of the distances between the four fingers (2 nd , 3 rd , and 4 th corners) upon maximum stretch
Fingers' flexion by measuring the finger pad to distal palmar line (mm)
2) Vasculopathy:
Raynaud's Condition Score (RCS)
Hands pain (VAS 0-100)
DU(s) outcome
Nailfold Capillary microscopy [28] <i>by a single, experienced evaluator using a stereomicroscope (SMZ-10 NIKON) at a stable temperature between 20-22°C</i>
3) Skin fibrosis:

Global Modified Rodnan Skin Score (MRSS) (scale 0-51) [29]

MRSS for the hands evaluated skin thickening on the dorsal hand and the first and second phalanges of the most affected finger

scale 0=no skin fibrosis, to 18= maximum fibrosis

Fingers edema assessed by circumference of fingers measured with a jeweler's rings

4) Overall disability: the Scleroderma Health Assessment Questionnaire (SHAQ) [30]

score ranged from 0=no disability, to 3 =severe disability

* Duration of the follow-up was chosen at 6 months because evaluation of safety requires observation of patients for a substantial period to capture the safety profile and preliminary efficacy endpoints

<http://www.fda.gov/downloads/medicaldevices/deviceregulationandguidance/guidancedocuments/ucm279107>.

Table S2: Radiological findings in the 12 patients with systemic sclerosis

Radiological findings	SSc patients (n = 12)
Number of patient with:	
Joint pattern	7
Erosion	3*
Wrists	2/3
MCP	0/3
PIP	0/3*
DIP	2/3*

Joint space narrowing	6*
Wrists	4/6
MCP	0/6
PIP	2/6*
DIP	3/6*
Arthritis (erosion+joint space narrowing)	1/6*
Bone pattern	3
Radiological demineralisation	0/3
Bone resorption	3/3
Acro-osteolysis	3/3
Resorption of distal ulna	0/3
Osteitis	0/3
Soft tissue pattern	7
Calcinosis	6/7
Flexion contracture	5/7

*not evaluable for 2 patients due to severe and irreducible digital flexion

Table S3: Effect of autologous adipose-derived SVF injection on digital ulcers outcome

	Number of DUs at baseline	Number of DUs at 2 months	Number of DUs at 6 months
Patient 1			
Dominant hand	0	0	0
Non dominant hand	1 (2mm diameter)	1 (1mm diameter)	0
Patient 3			
Dominant hand	4 (2mm diameter for 2, 3mm diameter for 2)	0	1 (3mm diameter)
Non dominant hand	1 (3mm diameter)	0	0
Patient 4			
Dominant hand	5 (4mm diameter for 2 and 1mm diameter for the 3 other)	7 (4mm diameter for 2 and 1mm diameter for the 5 other)	3 (1mm diameter for 2 and 2mm diameter for the third)
Non dominant hand	0	0	0
Patient 7			
Dominant hand	1 (necrotic ulcer 9mm x 7mm)	0	1 (healed ulcer 2mm diameter)
Non dominant hand	1 (scab in ulcer 8mm x 4mm)	0	1 (healed ulcer 1mm diameter)
Patient 10			
Dominant hand	1 (2mm diameter)	1 (scab in ulcer 1mm diameter)	0
Non dominant hand	1 (4mm diameter)	1 (2mm diameter)	1 (2mm diameter)