

FRI0319 PREDICTORS OF COMPLETE REMISSION IN POLYMYALGIA RHEUMATICA

D. Birra, A. Zoli, G. Peluso, S.L. Bosello, E. Gremese, G. Ferraccioli.
 Dipartimento di Reumatologia e Scienze Affini, Università Cattolica del Sacro Cuore - Fondazione Policlinico Universitario Agostino Gemelli, Rome, Italy

Background: Polymyalgia rheumatica (PMR) is an inflammatory rheumatic condition characterized by aching and morning stiffness in the shoulders, hip girdle, and neck that typically occurs in adults over the age of 50. A rapid resolution of symptoms with low-dose glucocorticoids is a feature of PMR although some patients may experience a disease flare-up during steroid tapering.

Objectives: The aim of the study was to investigate possible clinical or laboratory prognostic factors of remission during a 12 month follow up in PMR patients treated with a starting low prednisone dose following the 2015 ACR-EULAR guidelines⁽¹⁾.

Methods: From 86 consecutive outpatients, diagnosed with PMR following ACR/EULAR 2012 provisional clinical criteria for PMR⁽²⁾, 79 patients (56 women and 23 men), that achieved a complete follow up of at least 12 months, were selected. Clinical evaluation and laboratory tests were performed every 3 months. Clinical remission was defined as lack of shoulder and hip girdle pain and as levels of ESR \leq 40 mm/h and CRP \leq 0.5 mg/dL.

Results: 37 PMR patients reached a complete remission after twelve months follow-up. We didn't find any significant difference in the mean age and in ESR and CRP values at the beginning of the disease in patients in remission after 12 month of follow up when compared with patients not in remission. Presence of obesity, dyslipidemia, hypertension, diabetes and smoking habits were not significantly different in the two groups of patients. No significant difference in steroid therapy at the beginning and after 6 month of follow up was noted between the two groups of patients. A statistically significant female low clinical remission was seen at the end of 12 month follow up when compared with male complete clinical remission (33.9% versus 78.2%, $p=0.000$). Moreover it was shown that the patients achieving clinical and laboratory remission after six months of therapy were also those who maintained remission at the 12 month. CRP values, instead of ESR ones, were more predictive of remission after one year of therapy. Therefore patients with clinical remission in the absence of normalization of CRP value had greater risk of exacerbation.

Table 1. Demographic and laboratory characteristic of patients

	Remission T12	Non responders T12	p.
Age	75.1±6.7	69.7±5.7	ns
N*	37	42	
Female	19	37	.000
ESR T0 (mm/h)	56±31.6	50.2±29.4	ns
CRP T0 (mg/dL)	3.3±3.3	3.8±4.2	ns
ESR T6 (mm/h)	24.3±16.2	29.9±20.4	ns
CRP T6 (mg/dL)	0.6±0.7	1.5±2.1	.01
ESR T12 (mm/h)	22±15.4	28.7±17.4	ns
CRP T12 (mg/dL)	0.2±0.2	1±0.8	.000
Remission T6	27	7	.000
Prednisone dosage (mg) T0	16.3±8.4	15.5±6.6	ns
Prednisone dosage (mg) T6	5.9±4.8	7.5±5.1	ns
Prednisone dosage (mg) T12	2.3±2.5	4.4±4.7	.04
Dyslipidemia	9	12	ns
Hypertension	21	21	ns
Obesity	9	8	ns
Diabetes	6	8	ns
Smokers	3	3	ns

T0: Baseline; T6: Six months of follow up; T12: One year of follow up; ESR: erythrocyte sedimentation rate; CRP: C-Reactive Protein.

Conclusions: The sixth month of therapy is a crucial target for the management of PMR, because it can help to identify patients at greater risk of exacerbations, which may benefit from a tighter follow-up and a more aggressive therapeutic strategy. Among prognostic factors female sex and high CRP values at six months appear to be associated with higher relapse risk and a longer duration of treatment.

References:

[1] DeJaco C, Singh YP, European League Against Rheumatism; American College of Rheumatology, 2015 Recommendations for the management of polymyalgia rheumatica: a European League Against Rheumatism/American College of Rheumatology collaborative initiative. *Ann Rheum Dis*. 2015 Oct;74(10):1799–807. doi: 10.1136/annrheumdis-2015-207492.

[2] Dasgupta B, Cimmino MA, Matteson EL. 2012 provisional classification criteria for polymyalgia rheumatica: a European League Against Rheumatism/American College of Rheumatology collaborative initiative. *Ann Rheum Dis*. 2012 Apr;71(4):484–92. doi: 10.1136/annrheumdis-2011-200329.

Disclosure of Interest: None declared

DOI: 10.1136/annrheumdis-2017-eular.6262

FRI0320 PREDICTIVE VALUE OF PLATELET TO LYMPHOCYTE RATIO IN RENAL INVOLVEMENT IN PATIENTS WITH GRANULOMATOSIS WITH POLYANGIITIS

H. Kucuk¹, D. Tecer², O. Varan¹, H. Babaoglu¹, S.C. Guven², A. Tufan¹, S. Haznedaroglu¹, M.A. Öztürk¹, B. Goker¹. ¹Department of Internal Medicine, Division of Rheumatology; ²Department of Physical and Rehabilitation Medicine, Division of Rheumatology, Gazi University Faculty of Medicine, Ankara, Turkey

Background: Granulomatosis with polyangiitis (GPA) is a granulomatous necrotizing vasculitis with high morbidity and mortality. Anti-neutrophil cytoplasmic antibody (ANCA) is a valuable diagnostic marker, however its titer lacks predictive value for the severity of organ involvement. Platelet to lymphocyte ratio (PLR) and mean platelet volume has been regarded as a potential marker in assessing systemic inflammation.

Objectives: We aimed to investigate PLR and MPV as inflammatory marker in patients with GPA.

Methods: GPA patients and age-sex matched healthy controls were included. Demographic, clinical and laboratory information were extracted from medical records. Erythrocyte sedimentation rate (ESR), C-reactive protein (CRP), white blood cell (WBC), platelets (PLT), lymphocyte and neutrophils counts and glomerular filtration rate (GFR) were recorded. PLR was calculated. Disease activity was assessed with Birmingham Vasculitis Activity Score for WG vasculitis (BVAS/WG).

Results: 56 patients with GPA and 53 healthy controls were included. Clinical characteristics and laboratory findings of the study population are shown in Table 1. ESR, CRP, MPV and PLR were significantly higher in patients with GPA than controls. PLR positively correlated with ESR and CRP ($r:0.389$, $p:0.005$ and $r:0.512$ $p<0.001$, respectively). In contrast, MPV negatively correlated with ESR and CRP ($r:-0.308$, $p:0.028$ and $r:-0.337$ $p:0.014$, respectively). There were no significant correlation between PLR, MPV and BVAS/WG. Patients with renal involvement had statistically significantly higher PLR than patients without renal involvement (303.01 ± 287.33 vs $177.98 + 75.43$, $p: 0.020$ respectively). Moreover PLR negatively correlated with glomerular filtration rate ($r:-0.266$ and $p:0.009$). Receiver operating characteristic curve of PLR, ESR and CRP for differentiating renal involvement is presented in Figure 1. Area Under Curve (AUCs) for PLR, CRP and ESR were 0.703 (95% confidence interval [CI], 0.558–0.849, $p=0.016$), 0.577 (95% CI: 0.416–0.738, $p=0.362$), 0.508 (95% CI: 0.337–0.678, $p=0.929$), respectively. The cutoff level of PLR was 204 (sensitivity 65.6%, specificity 62.5%, positive predictive value 70%, negative predictive value 57.7%). Patients with alveolar hemorrhage tended to have higher PLR but this difference did not reach statistically significance ($266.60 + 182.90$ vs $240.61 + 252.43$ $p=0.382$, respectively).

Conclusions: Results suggest that PLR exhibit favorable diagnostic performance in predicting renal involvement in patients with GPA.

Table 1. Clinical characteristics and laboratory findings of the patients and healthy controls

	GPA patients	Controls	p value
Age (years)	48.14 ± 14.09	46.77 ± 14.14	0.614
Males (n)	33 (58.9%)	26 (49.1%)	0.034
WBC (x 10 ³ /uL)	12359.96 ± 6604.92	7230.80 ± 1630.33	<0.001
Neutrophils	9267.74 ± 5701.73	4285.73 ± 1597.18	<0.001
Lymphocytes	1805.64 ± 986.24	2311.22 ± 581.93	<0.001
Platelets	343640 ± 174905.10	234939.02 ± 57204.24	<0.001
ESR	54.19 ± 35.42	10.12 ± 7.66	<0.001
CRP	64.18 ± 72.11	3.42 ± 1.59	<0.001
MPV	7.73 ± 0.96	8.76 ± 1.10	<0.001
PLR (mean ±SD)	253.99 ±	104.79 ± 25.23	<0.001

Figure. ROC curve of PLR, ESR and CRP for differentiating renal involvement

