differences. The simple interpretation of the radiology is difficult and one has to be somewhat lenient. Mr. Kates also commented on the changes found at operation, which are usually more severe than one sees radiologically. With this too I would agree.

DR. P. J. L. HOLT (*London*) Some of the younger patients, and these are very often girls, complain of the postoperative scars. Now this is not wound breaking but wound stretching, and we were put off early mobilization because we felt that this was increasing the size of the scar when the tissues were tense.

DR. CAMP I have had no specific complaints of this nature from the patients, but those who were in the very young age groups, of which there are three patients with five knees, were in fact the ones who regained range of movement very slowly and we held back from manipulation or from forcing them. They did not appear to be left with an ugly scar.

Surgical Management of Severe Subluxations of the Rheumatoid Cervical Spine. By R. Q. CRELLIN, J. J. MACCABE, and E. B. D. HAMILTON (*King's College Hospital, London*).

21 patients with classical rheumatoid arthritis and severe subluxations of the cervical spine are reported. Fourteen had atlanto-axial subluxations with displacements ranging from 5 to 15 mm. (mean 9.5 mm.). The next most frequent level was at C4-5; this occurred in five patients with a displacement of from 3 to 10 mm. (mean 5 mm.). Thirteen patients underwent surgery because of symptoms or signs of spinal cord compression, or vertebral artery insufficiency. The operations comprised eight posterior fusions, three anterior fusions, and two laminectomies. The differential diagnosis, the radiological findings, the indications for surgery, and the results of treatment have been analysed.

Discussion

MR. D. O. HANCOCK (*Stoke Mandeville*) Could I ask what your operative indications are? Would you operate on all cases showing subluxation without neurological signs? If you would not, what sort of neurological signs do you look for? For example, would you operate for minimal hyperreflexia or would you prefer a little more? I quite agree about the extreme difficulty in assessing the power in these severely crippled patients, as I find it impossible to differentiate neurological loss from rheumatoid loss. Could I commend to your neurosurgeon's notice the excellent operation described by Newman and Sweetnam (1969) on the use of cancellous bone-chip grafts applied to the raw occiput and the arches of C1 and 2? This is a simple and most effective operation.

MR. CRELLIN I have also noticed this and it is certainly much simpler than passing wires around the laminae, which is why we employ a neurosurgeon. To answer the first point, we do not know when we should operate, and that is why I have suggested that several centres should co-operate to solve this problem. We have not felt that severe subluxation in itself was a reason to operate.

I do not think we shall know until the natural history of this condition is worked out. There are obviously many patients who have severe subluxation who have no evidence of spinal cord compression or vertebral artery trouble. Secondly, as far as the neurological assessment of the patient goes, it is an extremely difficult thing to decide. Our neurosurgeon puts great weight on two-point discrimination in his assessment of these patients, and speaking as an orthopaedic surgeon I probably would not notice anything wrong with them neurologically when he can! I think that if there is evidence of neurological impairment at a very early stage then it is time to operate. It is probably time to operate before that, but it is difficult to find out just when that should be. Are we to be guided by an actual amount of subluxation of x number of millimetres at the atlanto-axial joint, or should we wait for signs to occur? I do not know. I have left it an open question.

DR. W. W. BUCHANAN (*Glasgow*) I noticed the sharpening of the cervical spines. This is quite common in severe rheumatoid arthritis affecting the neck. I wonder if you have any ideas regarding the pathogenesis.

MR. CRELLIN I am afraid I have not. In one of the most severely affected of these patients, when we explored her neck from behind, we could find no evidence of cervical spines at all, at several levels. This is not obviously due to rheumatoid granulations or anything like that. I cannot explain why they disappear, but they just pull out like a piece of bubble gum and vanish.

NEWMAN, P. AND SWEETNAM, R. (1969) *J. Bone Jt. Surg.*, 51B 423

Dermal Arthroplasty. By B. N. BAILEY and S. N. DESAI (*Plastic Surgery Unit, Stoke Mandeville Hospital, Aylesbury*).

A technique of dermal arthroplasty of the finger joints has been developed in the Plastic Surgery Unit at Stoke Mandeville Hospital. The aim has been to achieve dynamic stability and restoration of a useful proportion of lost function and, if possible, of anatomical alignment. The results in 35 hands treated during the past 27 months have been encouraging. Certain aspects of preoperative assessment, surgical technique, and postoperative management have proved to be of paramount importance.

Discussion

DR. B. M. ANSELL (*Taplow*) Do you take any notice of wrist position, and does it in any way influence the end-results? Secondly, have you had any problems fairly late postoperatively—say after 6 months—with carpal tunnel and flexor tendon involvement causing a patient, who had previously been doing quite well, to lose movement, say by one year? Thirdly, what is the advantage of skin over silastic skid pads or prostheses?

MR. BAILEY To answer these questions one at a time:

Of course wrist position is terribly important. If we have a patient with a weak hand and weak flexors because of a painful wrist, then we do a fusion first—after testing as a preliminary step with splintage. If splintage improves

the power of the flexors then we go ahead and do an orthopaedic fusion of the wrist. We subsequently carry out arthroplasty or any other procedure which is indicated.

The second question was about carpal tunnels or flexor synovitis in the later stages. We have been fortunate so far in that we have not yet encountered this, although it is obviously something that may happen some time. If the patient has any suggestion of carpal tunnel compression or synovial proliferation on the flexor side, we deal with this as a preliminary before undertaking the arthroplasty.

The final question touches on the crux of the whole thing: has skin any advantage over silastic skid pads? Quite obviously it has or we should not use it; at least we think it has or we should not use it! In the first place the use of skin is founded on general principles. As plastic surgeons we have seen and removed over the course of years various materials from the human body—polyurethane, silastic, Tantalum gauze, steel and various lumps of different types of material; and although silastic is reputed to have been put into and to have stayed in the breast, the nose, and even the fingers, for years, we do see reactions. On general principles we feel that to put in autogenous tissue (quite apart from being cheaper, which it is by a long, long chalk) is probably a good thing to do, and so we started off by using it and have found that advantages accrued. One must have a spacer. I think this is quite obvious. If you cut off the ends of the bones something must separate them. What you choose may be the Brannan Klein prothesis—which is a complicated and expensive piece of metal and which even Adrian Flatt, its chief protagonist, admits never gives more than about two-fifths of the central range of movement. Alternatives like silastic protheses and other forms of artificial material require first of all a spacer and secondly a prong to go down the medulla of the metacarpal and proximal phalanx in order to provide rigidity. We prefer dermis on first principles and we prefer not to put anything down the shaft to provide rigidity if we can achieve this by tendon transfers, and this is the purpose of the indicis transfer, the abductor section, the plicating of the head, and the crossed intrinsic transfer. So overall we feel dermis has the advantage. With the few silastic implants of the Swanson type which we have done we have had rather poor results, not in any way comparable with those achieved by dermal arthroplasty.

DR. A. G. S. HILL (*Stoke Mandeville*) I should like to echo what Mr. Bailey has said. If one is thinking in terms of autogenous versus man-made materials, one should frame the question: 'What advantages has silastic over skin?', rather than the other way around, even though in terms of time the use of skin in this particular situation came after the use of silastic.

DR. G. NUKI (*Glasgow*) In your list of alternatives I think you missed out the Vainio arthroplasty in which the tendon is used as the interposing material. What is your opinion of this operation?

MR. BAILEY It is a splendid form of arthroplasty and we have used it. Like the Fowler operation it is very difficult, but those who often practise it achieve some quite magnificent results with it. We have found that by the time we have cut off the end of the bone, put the

tendon down, reunited the tendon, transferred the collateral ligaments and sewn it up again, 2 to 3 hours of tourniquet time has gone by. Unless we were prepared perhaps to sacrifice our first dozen, fifteen, or twenty cases knowing that the tourniquet time would be too long, I can not see how we should gain the expertise to do the job as rapidly as Vainio does by now. He is one of the few people who do many Vainio arthroplasties; it is a very good one based on sound principles, but it is very difficult and not quick enough. One can get through a dermal arthroplasty in an hour and 10 minutes, doing all the metacarpophalangeal joints of one hand.

DR. A. ST. J. DIXON (*Bath*) One very simple question. Why must you have a spacer at all? Mr. Kates, who is here, does a somewhat (to me as a physician) similar-looking operation in the feet, yet no spacer is put in and the toes can come back to very good function afterwards.

MR. BAILEY One very simple answer. You have not the 90° range of movement at the metatarsophalangeal joints that you have at the metacarpophalangeals.

MR. K. M. BACKHOUSE (*London*) I would agree with Dr. Dixon. In fact you do not need a spacer if you have good bone ends and a good muscle transfer as you are doing, or in the modification that we do, if the bone ends are not totally removed. I should, however, like to challenge you on this problem of the removal of pain by taking out all the synovium. In fact, if you have a lot of nasty bone-end and grating occurs, then patients may be in pain simply because the synovial pad has been removed, if you don't give a spacer. What you have done in fact, is to put in a beautiful surface lining which gives freedom of movement and then, which is far more important, allows the muscles to act effectively to carry out that movement.

Assessment of Clinical Methods of Evaluation and Isotope Techniques in Analgesic Trials in Rheumatoid Arthritis. By W. CARSON DICK, R. NEUFELD, M. GRAYSON, A. PRENTICE, A. WOODBURN, K. WHALEY, G. NUKI, and W. WATSON BUCHANAN (*The Centre for Rheumatic Disease and the University Department of Medicine, Royal Infirmary, Glasgow*).

A double-blind triple crossover study of indomethacin, aspirin, and placebo was undertaken. Ten patients took part in the trial for a period of 3 weeks, each receiving one of the treatments each week: thus each patient experienced all three treatments during the trial. Six methods of assessment were used: the Ritchie articular index, knee score, ring sizes, grip strength, radio-technetium peak uptake, and radioactive Xenon clearance rate. Analysis of variance techniques was employed to determine which methods best discriminated between the three treatments.

Discussion

DR. J. S. LAWRENCE (*Manchester*) You did not mention the erythrocyte sedimentation test at all. I presume this test was carried out?

DR. DICK It was not carried out. We were forced to draw the line somewhere. This was more a methodologi-