Therapeutic value of arthroscopy

H. A. BIRD AND E. F. J. RING

From the Royal National Hospital for Rheumatic Diseases, Bath

SUMMARY Thermography before and after joint irrigation at arthroscopy showed no significant improvement of inflammation in 14 patients. Subjectively, those with chondrocalcinosis did best.

Joint lavage has been advocated for years in the treatment of arthritis, though a controlled trial on 24 patients with classical rheumatoid arthritis using a 1.45 mm diameter cannula showed this to be of no more value than joint aspiration (Lindsay *et al.*, 1971). In diagnostic arthroscopy of the knee the joint can be irrigated with over one litre of dextrose saline through a 5 mm diameter arthroscope sheath. Observers have been impressed by the apparent subjective improvement after this procedure (Burman *et al.*, 1934; Jayson and Dixon, 1968; Jayson, 1976) and we decided to assess this using objective infrared thermography as a quantitative assessment of joint inflammation.

Accepted for publication May 24, 1977
Correspondence to Dr H. A. Bird, Rheumatism Research Unit, University of Leeds School of Medicine, 36 Clarendon Road, Leeds LS2 9PJ.

Patients and methods

A group of 14 patients referred for diagnostic arthroscopy of the knee were studied. The final clinical diagnosis is shown in the Table. After intra-articular injection of up to 20 ml of 2% lignocaine, a Storz arthroscope with a cannula diameter of 5 mm was inserted into one knee. All knee joints examined were irrigated with at least 1.0 litre of dextrose saline solution, run in through a separate polyethylene cannula inserted on the opposite side of the joint. Synovial biopsies were taken during the procedure.

Infrared thermography was performed on both knees 2 hours before arthroscopy and 8 days later. Thermographic index (TI) was calculated by the method of Collins *et al.* (1974) and change in TI compared between the two knees. The patients were assessed for pain improvement of the irrigated knee.

Table Details of 14 patients referred for arthroscopy

Case no.	TI change in irrigated knee	TI change in nonirrigated knee	Subjective improvement at 4 weeks			
			Much better	Better	Same	Worse
<i>Rheumatoid arthritis</i>						
1	+0.24	+0.14			1	
2	-1.76	-2.62				1
3	+0.54	-1.42		1		
<i>Seronegative arthritis</i>						
4	-0.01	-0.24	1			
5	-0.16	+0.55		1		
6	-0.46	+0.17			1	
7	-0.78	-1.28	1			
8	+0.43	-0.47	1			
<i>Osteoarthritis</i>						
9	+0.05	+0.14			1	
10	-0.43	+0.15		1		
<i>Chondrocalcinosis</i>						
11	-0.18	-0.54	1			
12	+0.08	+0.20	1			
13	-1.69	-1.11		1		
14	-0.91	-1.71	1			

P>0.5*

*Student's *t* test, not significant.
TI=thermographic index.

compared with the nonirrigated knee as much better, better, same, or worse at 1 and 4 weeks after arthroscopy. A single suture was removed at 1 week.

Results

13 out of 14 patients reported subjective improvement at 1 week but this was not always maintained. Subjective improvement at 4 weeks is shown in the Table. The patients with chondrocalcinosis and seronegative arthritis thought themselves better than those with seropositive rheumatoid arthritis and osteoarthritis. The mean fall in TI of arthroscoped knees was 0.36 units compared with 0.57 units for nonarthroscoped knees. This is not significant (Student's *t* test, $P > 0.5$). 2 patients, both with chondrocalcinosis, requested a further arthroscopy on the other knee.

Discussion

Some inflammation following tissue damage is inevitable in the arthroscoped knee. For this reason TI was assessed after 1 week when it was judged improvement would be maximal and traumatic inflammation minimal. In spite of this there was no significant improvement in TI after joint irrigation. Intra-articular steroid, for example, produces a constant improvement of about 1.00 units by this time (Esselinckx *et al.*, 1978).

Subjectively, patients with chondrocalcinosis, and to a lesser extent seronegative arthritis, fared best.

This is in keeping with the assumption that the therapeutic value of arthroscopy, if any, lies only in the evacuation of joint debris. Patients with seronegative arthritis often have large amounts of intra-articular fibrin that is broken down and removed (unpublished) and in chondrocalcinosis the value may lie in the removal of debris or alteration of solubility equilibrium. Further controlled evaluation of therapeutic arthroscopy in crystal deposition disease is required.

We thank Drs J. A. Cosh, A. St. J. Dixon, and P. A. Bacon for permission to report on their patients.

References

- Burman, M. S., Finkelstein, H., and Mayer, L. (1934). Arthroscopy of the knee joint. *Journal of Bone and Joint Surgery*, **16**, 255–268.
- Collins, A. J., Ring, E. F. J., Cosh, J. A., and Bacon, P. A. (1974). Quantitation of thermography in arthritis using multi-isothermal analysis. *Annals of the Rheumatic Diseases*, **33**, 113–115.
- Esselinckx, W., Bacon, P. A., Ring, E. F. J., Crooke, D., Collins, A. J., and Demottaz, D. (1978). A thermographic assessment of three intra-articular prednisolone analogues given in rheumatoid synovitis (in preparation).
- Jayson, M. I. V. (1976). Arthroscopy. *Clinics in Rheumatic Diseases*, **1**, 77–91.
- Jayson, M. I. V., and Dixon, A. St. J. (1968). Arthroscopy of the knee in rheumatic diseases. *Annals of the Rheumatic Diseases*, **27**, 503–511.
- Lindsay, D. J., Ring, E. F. J., Coorey, P. F. J., and Jayson, M. I. V. (1971). Synovial irrigation in rheumatoid arthritis. *Acta Rheumatologica Scandinavica*, **17**, 169–174.