Prostaglandin (PG) E₂ generation by cultured canine synovial fibroblasts exposed to microcrystals containing calcium

DANIEL J McCARTY AND HERMAN S CHEUNG

From the Division of Rheumatology, Department of Medicine, Medical College of Wisconsin, 8700 W Wisconsin Avenue, Milwaukee, Wisconsin 53226, USA

SUMMARY Immunoreactive prostaglandin (PG) E₂ was released into the ambient medium in a dose dependent fashion when either hydroxyapatite (HA) or calcium pyrophosphate dihydrate (CPPD) crystals were added to canine synovial fibroblasts in tissue culture. PGE₂ release peaked 6 to 9 hours after HA or CPPD crystals were added in the presence of serum but at 24 hours if they were added in the presence of lactalbumin hydrolysate. PGE₂ release correlated with crystal endocytosis estimated qualitatively by serial phase contrast microscopy and time lapse photography. As postulated previously by others for monosodium urate crystals, prostaglandin production by synovial cells may also be related to the pathogenesis of the destructive arthropathies associated with HA or CPPD crystals.

Key words: hydroxyapatite, calcium pyrophosphate.

Both acute inflammation and chronic destructive arthropathy have been described in association with basic calcium phosphate (BCP=hydroxyapatite, octacalcium phosphate, tricalcium phosphate) or calcium pyrophosphate dihydrate (CPPD) crystals.¹⁻⁵ Such tissue reactions resemble in some respects the acute inflammatory attack and chronic (tophaceous) destructive arthropathy associated with gout, the hallmark of which is the formation in tissues of monosodium urate (MSU) crystals. Each of these crystal species undergoes endocytosis by neutrophils and to a lesser extent by monocytes in acute inflammatory exudates.^{1,3,6} Products of both the lipoxygenase and cyclo-oxygenase pathways of arachidonic acid metabolism have been identified when MSU crystals are exposed to these cells in vitro (reviewed by McCarty⁷), and products of both pathways have been identified when cultured human synovial fibroblasts⁸ or human neutrophils⁹ were exposed to MSU crystals. Cyclo-oxygenase products, mostly prostaglandin (PG)E₂ were elaborated in impressive quantities when MSU crystals were

added to murine macrophages¹⁰ or lapine synovial fibroblasts.^{11,12} Microcrystalline, but not amorphous, calcium oxalate produced effects similar to that of MSU crystals when incubated with cultured lapine synovial fibroblasts.¹³

PGE₂, and to a lesser extent, PGF_α, generation by cultured human or canine synovial fibroblasts and by lapine chondrocytes in primary culture¹⁵ was augmented when cells were incubated with hydroxyapatite or CPPD crystals in concentrations greater than those actually found in synovial fluid from patients. As prostaglandins have been implicated as mediators in both acute inflammation¹⁶ and in bone resorption,¹⁷ they may relate to the pathogenesis of both the acute and chronic reactions to microcrystals.

The work reported here was performed with highly characterised synthetic hydroxyapatite (HA) or CPPD crystals in concentrations actually found in human joint fluid.

Materials and methods

General. All reagents were prepared in double distilled, glass distilled, charcoal filtered water.

Cultures. Canine synovial cells were obtained from anaesthetised animals by a modification of the

Accepted for publication 31 July 1984.

Correspondence to Professor Daniel J. McCarty, MD, Department of Medicine, Medical College of Wisconsin, 8700 W Wisconsin Avenue, Milwaukee, Wisconsin 53226, USA.

method of Clarris *et al.*¹⁸ Visually confluent second passage cultures from four mongrel dogs were used experimentally. There were about 1.5 to 2 × 10⁶ cells/plate. Cells were grown in 100 mm plastic dishes (Scientific Products, McGaw Park, Ill.) containing 10 ml of Dulbecco's modified Eagle's essential medium (DMEM) supplemented with 10% horse serum and 1% penicillin, streptomycin, and amphotericin B (all from Gibco, Grand Island, NY). At the end of each experiment the cells were washed three times with Hanks's balanced salt solution, released with 2 ml of 20 mM EDTA, and precipitated with an equal volume of 20% trichloroacetic acid. After dissolution of the centrifuged precipitate in 2 ml 0.1 N NaOH, the protein content was measured.¹⁹

Crystals. Hydroxyapatite crystals were a kind gift from M D Francis, PhD, the Proctor and Gamble Co., Cincinnati, Ohio. They had a molar Ca/P of 1.67 and showed only HA adsorption bands by Fourier transform infrared spectrophotometry.²⁰ Crystal aggregates were sieved to 10–20 μm with Endicott test sieves (London, England) and heated to 200°C for 2 h to assure sterility and to eliminate possible pyrogen contamination. Calcium pyrophosphate dihydrate crystals were synthesised, characterised, and sieved to 10–20 μm and heated as described previously.²¹ Monosodium urate monohydrate crystals were prepared and characterised as described previously.²² They were sonicated to 10–20 μm and then heated to 170°C for 2 h, which does not alter crystal structure.²³

Prostaglandin (PG) E₂ assay. This was done by radioimmunoassay by the method of Dunn *et al.*²⁴ From 0.5 to 3 ml of conditioned medium was extracted for each determination to give the desired sensitivity. The higher values required dilution to give results that fell on the standard curve.

Experiments. Culture medium was replaced with

DMEM alone, DMEM with 10% serum, or 0.2% lactalbumin hydrolysate (LA) with or without HA crystals, 100 μg/ml (mg/l) PGE₂ generation by synovial cells was determined. All cultures were examined at intervals up to 24 hours by phase-contrast microscopy. Several cultures containing crystals were examined continuously by time lapse photography with a microscope equipped with a camera (Nikon).

Samples of conditioned media were removed for PGE₂ assay at periodic intervals up to 24 hours, at which time the cells were released and protein content measured. Similar experiments were performed with MSU or CPPD crystals. Dose response was determined by addition of HA or CPPD crystals in concentrations varying from 0 to 500 μg/ml. Indomethacin was added to at least one plate containing crystals and serum in each experiment at final concentration of 10⁻⁶M. PGE₂, 45 500 pg/ml (ng/l) and 2400 pg/ml (ng/l) were incubated in DMEM–10% horse serum that had been conditioned for 24 hours by incubation with synovial cells, and samples were removed for radioimmunoassay at hourly intervals for the first six hours and then at 24 hours.

Results

All data are expressed as pg of PGE₂ per mg of cell protein. Cell protein per plate varied from 350 to 900 μg. Except for occasional clumps, nearly all crystals appeared to have undergone endocytosis within 4 to 6 hours when serum was present. In the presence of DMEM alone or when 0.2% LA was substituted, intracellular crystals were noted in large numbers only at 24 hours. These observations were confirmed by time lapse photography.

PGE₂ concentration in the conditioned medium increased in all cultures (Table 1). The increased

Table 1 PGE₂ (pg/mg cell protein) release from cultured canine synovial cells by hydroxyapatite (HA) crystals

Time (h)	DMEM	DMEM +HA [*]	DMEM +HA [†]	DMEM 10% HS‡ mean±SEM (n=4)	DMEM 0.2% LA§	DMEM 10% HS +HA [†]	DMEM 0.2% LA+HA [*]	DMEM 10% HS+HA+Ind
0	0	0	0	114	0	120	0	
3	24	36	19	2480±588	159±77	5,734±1,166	155±67	
6	31	87	78	3659±1068	3565±1789	18,907±1,790	2,609±1,564	
9	124	471	714	4629±954	3424±1540	23,939±4,057	5,227±2,978	
24	821	30,285	88,109	2080±468	3238±1496	46,149±18,911	70,884±49,098	100±33

^{*} HA=hydroxyapatite crystals, opsonised in 10% horse serum in DMEM washed 3 times; 100 μg/ml (mg/l) used throughout 10–20 μm aggregate size.

[†] HA=not opsonised.

[‡] HS=horse serum.

[§] LA=lactalbumin hydrolysate.

^{||} Ind=indomethacin 10⁻⁶M.

level in the presence of DMEM alone was just detectable; in the presence of 10% serum or 0.2% LA, PGE₂ levels increased after 3 to 6 hours but remained stable. HA crystals added with 10% serum produced progressively greater PGE₂ levels up to 24 hours; when incubated with DMEM alone or with 0.2% LA the PGE₂ rise was delayed but showed a very large rise at 24 hours, sometimes exceeding that noted with crystals in the presence of serum (Fig. 1). Repeated experiments with cells from different animals gave widely variable results in terms of absolute numbers, but the shapes of the curves representing the time course of PGE₂ release were nearly identical.

The time course of release of PGE₂ from canine synovial cells by triclinic CPPD crystals and by MSU crystals is shown in Table 2. In all cultures MSU crystals resulted in a massive and progressive release of PGE₂ into the ambient medium. CPPD behaved like HA crystals, producing an early (3–6 h) rise in the presence of serum. The PGE₂ concentrations at 9 and 24 hours were similar with opsonised crystals in DMEM alone or in DMEM with 10% serum.

The PGE₂ released into the culture medium 9 hours after addition of varying concentrations of HA or CPPD crystals is shown in Table 3. The levels increased in proportion to the concentration of CPPD crystals and to the concentration of HA crystals at 50, 100, and 500 µg/ml (mg/l). Concentrations of HA crystals of 10 µg/ml (mg/l) or less were not associated with PGE₂ generation greater than that found in 10% horse serum alone.

Immunoreactive PGE₂ levels decreased gradually over time when incubated in conditioned media. Only 71% and 48% of the higher concentration

Fig. 1 Time course of PGE₂ released from cultured canine synovial cells after addition of HA crystals (---) 100 µg/ml (mg/l) in DMEM+10% HS, or (—) in DMEM+0.2% lactalbumin hydrolysate. Data shown as mean±SEM, n=4. At the lower concentration 66% and 48% remained at 6 and 24 hours respectively. At the higher concentration 66% and 48% remained at 6 and 24 hours respectively. At the lower concentration 66% and 48% remained at these times (data not shown).

Table 2 PGE₂ (pg/mg cell protein) released from cultured canine synovial cells by monosodium urate (MSU) or calcium pyrophosphate dihydrate (CPPD) crystals

Time (h)	MSU (500 µg/ml)			CPPD (100 µg/ml)		
	DMEM*	DMEM 0.2% LA†	DMEM 10% HS‡	DMEM OP§	DMEM 10% HS	DMEM 10% HS+Ind
0	0	ND¶	ND	0	324	—
3	20,031	15,535	11,888	0	742	—
6	39,891	28,469	62,598	2,244	11,904	—
9	152,036	84,201	200,618	21,083	19,641	—
24	304,581	ND	ND	4,822	4,636	128

* DMEM=Dulbecco's modified Eagle's medium.

† LA=lactalbumin hydrolysate.

‡ HS=horse serum.

§ OP=opsonised crystals.

|| Ind=indomethacin 10⁻⁶M.

¶ ND=not measured.

Table 3 PGE₂ released from cultured canine synovial cells by various concentrations of crystals after nine hours of incubation

HA* (µg/ml)	PGE ₂ [†] (pg/mg protein)	CPPD [‡] (µg/ml)	PGE ₂ [†] (pg/mg protein)	PGE ₂ [§] (pg/mg protein)
0	4.630	0	3.454	333
1	2.734	5	6.784	1.058
10	3.834	10	8.494	1.063
50	13.704	100	10.623	2.100
100	18.526	500	20.893	14.344
500	24.457			

* HA=hydroxyapatite 10–20 µm aggregate size.

† DMEM=10% horse serum.

‡ CPPD=calcium pyrophosphate dihydrate—10–20 µm.

§ DMEM=opsonised crystals only.

SI conversion: µg/ml=mg/l.

Discussion

The results of these experiments with canine cells confirm previous reports that monosodium urate crystals stimulate PGE₂ production by cultured human⁸ or lapine^{11–13} synovial fibroblasts. Although precise comparison is not possible because of the variability of response when cells from different animals are used, it seems evident that hydroxyapatite or calcium pyrophosphate dihydrate crystals are much less potent stimulators of PGE₂ production than are MSU crystals (Tables 1 and 2). PGE₂ levels in the ambient culture medium showed a definite lag period after addition of either HA or CPPD crystals in serum, with a marked increase between 3 and 6 hours and a further rise at 9 and/or 24 hours. PGE₂ release correlated with qualitative assessment of endocytosis estimated by serial phase contrast microscopy and time lapse photography, as noted previously with MSU crystals.⁸ This release was found when synthetic HA crystals were used with properties similar to natural BCP crystals but only in concentrations of 50 µg/ml (mg/l) or greater, levels not generally found in patients joint fluid.²⁵ PGE₂ release by CPPD crystals was noted even at 5 µg/ml (mg/l), far less than the mean of 72 µg/ml (mg/l) found in chronic and 280 µg/ml (mg/l) found in acute effusions.²⁶

Both endocytosis and PGE₂ generation occurred but were delayed when either opsonised or uncoated crystals were added to cultured cells in the absence of serum. Wigley *et al.* found that serum was required for PGE₂ generation from human synovial cells and that this was not accompanied by cell death as estimated by lactate dehydrogenase (LDH) release into the culture fluid.⁸ Hasselbacher and colleagues found no need for serum when MSU crystals were added to murine macrophages or

rabbit synovial fibroblasts, but LDH was released at 2 hours and this became massive after 20 hours.^{10–13} We had previously reported no significant LDH release when either synovial cells¹⁴ or lapine chondrocytes¹⁵ were exposed to HA or CPPD crystals in the presence of serum.

MSU, calcium oxalate, BCP, or CPPD crystals are all capable of releasing both PGE₂ and collagenase from cultured cells. The latter event was originally demonstrated by Werb and Reynolds by incubating cultured lapine cells with latex beads²⁷ and the former by Fine *et al.*, who exposed human synovial fibroblasts and murine macrophages to MSU crystals.²⁸ The mechanism of prostaglandin generation by cells stimulated with any of these particulates remains to be determined.

It is possible that crystal-induced release of prostaglandins from synovial cells contributes to the destructive arthropathies which occur in some patients with BCP or CPPD crystal deposition diseases. Determinations of the types and amounts of eicosinoids in synovial fluid from patients with these conditions may shed further light on this possibility.

This work was supported by USPHS AM26062, a grant from Kroc Foundation, and Research Career Development Award AM01065.

References

- 1 McCarty D J, Gatter R A. Recurrent acute inflammation associated with focal apatite crystal deposition. *Arthritis Rheum* 1966; **9**: 804–19.
- 2 Schumacher H R, Smolyo A P, Tse R L, Maurer K. Arthritis associated with apatite crystals. *Ann Intern Med* 1977; **87**: 411–6.
- 3 McCarty D J, Kohn N N, Faires J S. The significance of calcium phosphate crystals in the synovial fluid of arthritis patients: the 'pseudogout syndrome'. I. Clinical aspects. *Ann Intern Med* 1962; **56**: 711–37.

- 4 Menkes C J, Simon F, Delrieu F, Forest M, Delbarre F. Destructive arthropathy in chondrocalcinosis articularis. *Arthritis Rheum* 1976; **19**: 329-48.
- 5 McCarty D J, Halverson P B, Carrera G F, Brewer B J, Kozin F. 'Milwaukee shoulder': Association of microspheroids containing hydroxyapatite crystals, active collagenase and neutral protease with rotator cuff defects. I. Clinical aspects. *Arthritis Rheum* 1981; **24**: 464-73.
- 6 McCarty D J. Phagocytosis of urate crystals in gouty synovial fluid. *Am J Med Sci* 1962; **243**: 288-95.
- 7 McCarty D J. Pathogenesis and treatment of crystal induced inflammation. In: McCarty D J ed. *Arthritis and allied conditions*. 10th ed. Philadelphia: Lea and Febiger, in press.
- 8 Wigley F M, Fine I T, Newcombe D S. The role of the human synovial fibroblast in monosodium urate crystal-induced synovitis. *J Rheumatol* 1983; **10**: 602-11.
- 9 Schran C N, Lundberg U, Abramson S, Samuelsson B, Weissmann G. Formation of leukotrienes (LT) and hydroxy acids by human leukocytes exposed to monosodium urate (MSU). *Arthritis Rheum* 1983; **26**: 59 (abstr).
- 10 McMillan R M, Hasselbacher P, Hahn J, Harris E D Jr. Interactions of murine macrophage with monosodium urate crystals: stimulation of lysosomal enzyme release and prostaglandin synthesis. *J Rheumatol* 1981; **8**: 555-62.
- 11 McMillan R M, Vater C A, Hasselbacher P, Hahn J, Harris E D Jr. Induction of collagenase and prostaglandin synthesis in synovial fibroblasts treated with monosodium urate crystals. *J Pharm Pharmacol* 1981; **33**: 382-3.
- 12 Hasselbacher P, McMillan R M, Vater C A, Hahn J, Harris E D Jr. Stimulation of collagenase and PGE₂ by synovial fibroblasts in response to crystals of monosodium urate monohydrate: a model for joint destruction in gout. *Trans Assoc Am Physicians* 1981; **94**: 243-52.
- 13 Hasselbacher P. Stimulation of synovial fibroblasts by calcium oxalate and monosodium urate monohydrate. A mechanism of connective tissue degradation in oxalosis and gout. *J Lab Clin Med* 1982; **100**: 977-85.
- 14 Cheung H S, Halverson P B, McCarty D J. Release of collagenase, neutral protease and prostaglandins from cultured mammalian synovial cells by hydroxyapatite and calcium pyrophosphate dihydrate crystals. *Arthritis Rheum* 1981; **24**: 1338-44.
- 15 Cheung H S, Halverson P B, McCarty D J. Phagocytosis of hydroxyapatite or calcium pyrophosphate dihydrate crystals by rabbit articular chondrocytes stimulates release of collagenase, neutral protease and prostaglandins E₂ and F_{2α}. *Proc Soc Exp Biol Med* 1983; **173**: 181-9.
- 16 Moncada S, Vane J R. Pharmacology and endogenous roles of prostaglandin endoperoxides, thromboxane A₁ and prostaglandin. *Pharmacol Rev* 1979; **30**: 293-335.
- 17 Robinson D R, Tashjian A H Jr, Levine J. Prostaglandin-stimulated bone resorption by rheumatoid synovia. *J Clin Invest* 1975; **56**: 1181-8.
- 18 Clarris B J, Fraser J R E, Moran C J, Muirden K D. Rheumatoid synovial cells from intact joints. *Ann Rheum Dis* 1977; **36**: 293-301.
- 19 Lowry O H, Rosebrough N J, Farr A L, Randall J R. Protein measurement with the folin phenol reagent. *J Biol Chem* 1951; **193**: 265-75.
- 20 McCarty D J, Lehr J R, Halverson P B. Crystal populations in human synovial fluid. Identification of apatite, octacalcium phosphate and tricalcium phosphate. *Arthritis Rheum* 1983; **26**: 1220-4.
- 21 McCarty D J, Palmer D W, James C. Clearance of calcium pyrophosphate dihydrate (CPPD) crystals in vivo II. Studies using triclinic crystals doubly labelled with ⁴⁵Ca and ⁸⁵Sr. *Arthritis Rheum* 1979; **22**: 1122-31.
- 22 Cheung H S, Story M, McCarty D J. Mitogenic effects of hydroxyapatite and calcium pyrophosphate dihydrate crystals on canine synovial cells in culture. *Arthritis Rheum* 1984; **27**: 668-74.
- 23 Mandel N S. Structural changes in sodium urate crystals on heating. *Arthritis Rheum* 1980; **23**: 772-6.
- 24 Dunn M J, Laird J F, Dray F. Basal and stimulated rates of renal secretion and excretion of prostaglandins E₂, F_{2α}, and 13-14 dihydro 15 beta F_{2α} in the dog. *Kidney Int* 1978; **13**: 136-43.
- 25 Halverson P B, Cheung H S, McCarty D J, Garancis J, Mandel N. 'Milwaukee shoulder': association of microspheroids containing hydroxyapatite crystals, active collagenase and neutral protease with rotator cuff defects. II. Synovial fluid studies. *Arthritis Rheum* 1981; **24**: 474-83.
- 26 Silcox D C, McCarty D J. Elevated inorganic pyrophosphate concentrations in synovial fluid in osteoarthritis and pseudogout. *J Lab Clin Med* 1974; **83**: 518-31.
- 27 Werb Z, Reynolds J J. Stimulation by endocytosis of the secretion of collagenase and neutral proteinase from rabbit synovial fibroblasts. *J Exp Med* 1974; **140**: 1482-97.
- 28 Fine I, Sannes P, Newcombe D S. Human synovial fibroblasts and murine macrophages: urate phagocytosis and prostaglandin release. *Clin Res* 1980; **28**: 146A (abstr).