Case reports

Acute transverse myelopathy complicating systemic lupus erythematosus

D J PROPPER AND R C BUCKNALL

From the Rheumatic Diseases Unit, Royal Liverpool Hospital, Liverpool

SUMMARY A sixteen year old girl with systemic lupus erythematosus developed acute transverse myelopathy. She was treated with high dose steroids, cyclophosphamide, and plasma exchange and regained partial neurological function. Previous descriptions of transverse myelopathy complicating systemic lupus erythematosus are reviewed, with particular reference to the efficacy of high dose steroid treatment.

Transverse myelopathy is a rare complication of systemic lupus erythematosus, and the prognosis is generally poor.¹ Although it has been suggested that steroids confer no benefit,² there are several reports of complete or partial recovery associated with high dose steroid treatment. We describe a patient with transverse myelopathy associated with systemic lupus erythematosus treated, uniquely, with steroids, cyclophosphamide, and plasma exchange. Moreover, we review the published work concerning the treatment and prognosis of this condition, and the usual cerebrospinal fluid findings.

Case report

A sixteen year old negroid girl presented with a three month history of polyarthralgia and Raynaud's phenomena. Examination was normal apart from cyanotic mottling over the finger tips and florid splinter haemorrhages affecting the fingernails.

Initial investigations were as follows: antinuclear antibody titre 1/1024, DNA binding 21% (normal <20%), C3 0.59 g/l (normal 0.7–1.7 g/l), C4 0.12 g/l (normal 0.15–0.55 g/l). Despite treatment with prednisolone 60 mg daily the digital vasculitis progressed. Hence treatment with intravenous cyclophosphamide 1 g weekly was started, and she underwent three plasma exchanges, totalling nine litres. The digital lesions subsequently healed. Over the next eight months apart from mild arthralgia she was well, and treatment was reduced to prednisolone 2.5 mg daily. Subsequently she developed active

synovitis, associated with rising antinuclear antibody titres and falling complement concentrations, and was readmitted. Three days after admission she complained of mid-scapular pain and numbness affecting the legs, and became paraplegic over four hours. She had a sensory level at T7, below which pain and temperature sensation were absent, though vibration sense and proprioception were preserved. She developed urinary retention and faecal incontinence. Lumbar puncture showed normal cerebrospinal fluid pressure. Cerebrospinal fluid analysis showed protein 1100 mg/l (normal <600 mg/l), glucose 2.0 mmol/l (normal 3.9–6.1 mmol/l), and white cell count $21 \times 10^9/l$. Myelography was normal. A diagnosis of transverse myelopathy was made, and on the day of onset she received methylprednisolone 1 g and cyclophosphamide 1 g, both intravenously. Prednisolone was increased to 80 mg daily. Over the next three days she underwent three plasma exchanges, totalling nine litres.

No further neurological deterioration occurred. After two months movement returned, initially to the right foot, and after four months bladder and bowel control returned. Eighteen months after the onset she had sufficient lower limb power to walk with the aid of crutches. Throughout the illness antiphospholipid antibodies were not detected in the sera.

Discussion

There are 44 reported cases of transverse myelopathy occurring in SLE,^{1–30} though the prevalence is probably higher. Twenty six,^{1–18} including the present case, are reported in sufficient detail to enable meaningful comparison (Table 1). We wish

Accepted for publication 25 October 1988.

Correspondence to Dr D J Propper, Department of Medicine and Therapeutics, Polwarth Building, Foresterhill, Aberdeen AB9 2ZD.

Table 1 Comparison of treatments and outcome in transverse myelopathy associated with systemic lupus erythematosus

Reference number	Age and sex	Diagnosis before onset of myelopathy	Time from onset of myelopathy to start of treatment	Drug treatment	Neurological outcome	Time from onset of myelopathy to neurological outcome
1	24 F	Rheumatoid disease	1 day	Decadron 8 mg/day	No change	3½ months
1	22 F	None	1 day	Prednisolone 100 mg/day	No change	3 months
1	56 F	SLE*	1 day	Prednisolone 60 mg/day	Death	40 days
2	24 F	SLE	Not known	Prednisolone 20 mg/day	No change	9 years
2	35 M	SLE	Not known	Prednisolone 20 mg/day	No change	12 years
3	47 F	SLE	Same day	Prednisolone 120 mg/day	Partial recovery	4 months
4	21 F	SLE	Same day	Prednisolone 20 mg/day, azathioprine 100 mg/day	Death	1 day
5	16 F	SLE	Same day	Prednisolone 100 mg/day	Partial recovery	4 weeks
6	27 F	None	3 weeks	Prednisolone	No change	5 months
6	53 F	None	3 weeks	Prednisolone	No change	10 months
7	35 F	SLE	None	None	Death	4 days
8	19 F	SLE	Receiving treatment before	Cortisone 75 mg/day	Death	45 days
9	40 M	None	Not known	Antibiotics	Recovery	—
10	28	SLE	Not known	Cortisone 200 mg/day	Recovery	80 days
11	26 M	SLE	Receiving treatment before	Steroids, ACTH*	Death	3 days
12	43 F	SLE	Same day	Cortisone 100 mg/day	Death	1 year
13	11 F	SLE	Receiving treatment before	Cortisone 50–100 mg/day	Death	14 days
13	12 F	Pyrexia of unknown origin	4 days	Dexamethasone 24 mg/day	Partial recovery	6 months
13	35 F	None	8 months	Prednisolone 80 mg/day	No change	2 years
14	33 M	Idiopathic thrombocytopenic purpura	20 days 30 days	Steroids, cyclophosphamide	Death	35 days
15	41 M	Rheumatoid disease	Not known	Prednisolone	Death	20 days
16	51 F	SLE	Not known	Prednisolone	Death	20 months
17	23 F	SLE	Same day	Steroids-high dose	Recovery	1 month
18	29 F	None	—	None	Continued deterioration	—
18	33 F	SLE	Not known	Not known	Continued deterioration	—
This case	16 F	SLE	Same day	Prednisolone 80 mg/day, cyclophosphamide 1 g, plasma exchange	Partial recovery	18 months

*SLE=systemic lupus erythematosus; ACTH=corticotrophin.

to review two aspects of this condition: firstly, the effect of high dose steroid treatment and, secondly, cerebrospinal fluid glucose concentrations at the onset of myelopathy.

Of the 26 cases, seven recovered either full or partial neurological function, in nine the neurological deficit remained static or deteriorated further, and the remaining 10 patients died. For

eight patients treatment with high dose steroids was started at the onset of myelopathy, for seven of these within 24 hours and for the eighth within four days. Of these, five recovered either full or partial neurological function, one died, and in two the neurological deficit was unchanged. Eighteen patients were treated at the onset of myelopathy with either low dose steroids or a variety of drugs, or given no drug treatment at all. Of this group, two recovered, seven experienced no change in neurological deficit, and nine died.

Although it has been suggested that steroids are ineffective in transverse myelopathy due to systemic lupus erythematosus,² this review suggests that significant benefit may be gained from their early administration. Furthermore, the published work contains a further four cases where high dose steroids were administered with subsequent improvement in neurological function,²⁵⁻²⁷ but as insufficient detail was given in these reports they are not included in this review. We can only postulate the mode of action of high dose steroids in this condition. Vasculitis is a prominent feature in the spinal cord at postmortem examination,¹ which suggests that steroids may act by suppressing the vasculitic process. If this is the case then plasma exchange is a logical treatment as it may reduce

concentrations of circulating immune complexes and inflammatory mediators.

A review of the cerebrospinal fluid findings in patients with transverse myelopathy due to systemic lupus erythematosus shows a significantly reduced glucose concentration in many of the cases (Table 2), though it must be noted that simultaneous blood sugars were not measured. In fact of the 10 patients known to have undergone lumbar puncture on the day of onset of myelopathy, all had low cerebrospinal fluid glucose concentrations. Low cerebrospinal fluid glucose is not, however, a common feature of central nervous system lupus. Gibson and Myers, on reviewing the cerebrospinal fluid findings of 80 patients with central nervous system lupus, found only four instances, one of whom had a transverse myelopathy.²⁸ Duffy *et al* reviewing 21 cases of central nervous system lupus found one instance,³¹ and Feinglass found none in 44 cases.²⁴ Furthermore, low cerebrospinal fluid glucose concentrations are not noted to accompany idiopathic transverse myelopathy.³² As 10 of the 26 cases in this review were not known to have systemic lupus erythematosus at the onset of transverse myelopathy the finding of a low cerebrospinal fluid glucose in cases of transverse myelopathy arising afresh may have important diagnostic implications.

In conclusion, we suggest that the early administration of high dose steroids for transverse myelopathy due to systemic lupus erythematosus may decrease mortality and reduce the eventual neurological deficit in survivors.

Table 2 Summary of cerebrospinal fluid findings in transverse myelopathy associated with systemic lupus erythematosus

Reference number	Time from onset of myelopathy to lumbar puncture	Glucose (mmol/l)	Protein (mg/l)	Number of cells ($\times 10^6/l$)
1	Same day	0.8	1150	—
1	Same day	1.0	3250	—
1	Same day	1.4	1150	—
2	24 hours	Normal*	1280	430
2	Unknown	Normal	—	1600
3	Same day	2.2	4050	940
4	Same day	1.2	1200	48
5	Same day	1.3	220	0
6	Same day	1.6	880	29
6	Same day	1.8	5850	3
7	Unknown	—	400	7
8	Unknown	0.9	1750	16
11	Same day	1.1	530	39
12	Unknown	—	400	7
13	Unknown	3.8	430	16
13	Unknown	—	3400	—
15	Unknown	—	1400	413
16	Unknown	—	Normal*	27
17	1 month	—	1500	49
This case	Same day	2.0	1100	—

*Normal ranges: glucose 3.9–6.1 mmol/l; protein <600 mg/l.

References

- 1 Andrianakos A A, Duffy J, Suzuki M, Sharp J T. Transverse myelopathy in systemic lupus erythematosus: report of three cases and review of the literature. *Ann Intern Med* 1975; **83**: 616–24.
- 2 Hachen H J, Chantaine A. Spinal cord involvement in systemic lupus erythematosus. *Paraplegia* 1979; **17**: 337–46.
- 3 Henry A K, Brunner C M. Relapse of lupus transverse myelitis mimicked by vertebral fractures and spinal cord compression. *Arthritis Rheum* 1985; **28**: 1307–10.
- 4 Scharf I, Nahir M, Hemli J. Transverse myelitis with systemic lupus erythematosus. *J Neurol* 1977; **215**: 231–2.
- 5 Warren R W, Kredich D W. Transverse myelitis and acute central nervous system manifestations of systemic lupus erythematosus. *Arthritis Rheum* 1984; **27**: 1058–60.
- 6 Kewalrami L S, Saleem S, Bertrand D. Myelopathy associated with systemic lupus erythematosus (erythema nodosum). *Paraplegia* 1978; **16**: 282–94.
- 7 Fisher J H, Gilmour J R. Encephalomyelitis following administration of sulphonamide. *Lancet* 1939; **ii**: 301–5.
- 8 Piper P G. Disseminated lupus erythematosus with involvement of the spinal cord. *JAMA* 1953; **153**: 215–7.
- 9 Siekert R G, Clark E C. Neurological signs and symptoms as early manifestations of systemic lupus erythematosus. *Neurology* 1955; **5**: 84–8.
- 10 Mustata N, Trica L. Lupus eritematos diseminat cu manifestari neurologice. *Viata Medicala* 1960; **7**: 445.

- 11 Orthner H, Rossner R. Chronisch rezidivierender lupus erythematosus visceralis mit akut todlicher zentralnervösen exacerbation. *Deutsche Zeitschrift für Nervenheilkunde* 1965; **187**: 1–24.
- 12 Johnson R T, Richardson E P. The neurological manifestations of systemic lupus erythematosus. A clinical-pathological study of 24 cases and review of the literature. *Medicine (Baltimore)* 1968; **47**: 337–9.
- 13 Penn A S, Rowan J R. Myelopathy in systemic lupus erythematosus. *Arch Neurol* 1968; **18**: 337–40.
- 14 Sinkovics J G, Gyorkey F, Thoma G W. A rapidly fatal case of systemic lupus erythematosus: structure resembling viral nucleoprotein strands in the kidney and activity of lymphocytes in culture. *Texas Reports on Biology and Medicine* 1969; **27**: 887–908.
- 15 Castaing R, Vital C, Cardinaud J P, et al. Ramollissement médullaire au cours d'un lupus erythemateux disseminé. Etude anatomo-clinique d'une observation. *Bordeaux Medicine* 1970; **3**: 707–18.
- 16 Andrews J M, Cancilla P A, Kunin J. Progressive spinal cord signs in a patient with disseminated lupus erythematosus. *Bulletin of the Los Angeles Neurological Society* 1970; **35**: 78–85.
- 17 Vitale C, Kahn M F, De Sèze M, De Sèze S. Névrite optique myélite et maladie lupique. A propos de deux observations. *Ann Med Interne (Paris)* 1973; **124**: 211–6.
- 18 Tharkarar P, Greenspun B. Transverse myelopathy in systemic lupus erythematosus. *Arch Phys Med Rehabil* 1979; **60**: 323–4.
- 19 Nasonova V A, Konchakova M I. The course of systemic lupus erythematosus with neurological syndromes. *Voprosy Reumatizma* 1965; **5**: 12–16.
- 20 Bas H, Vachtenheim J. Neurologische erscheinungen beim systematischen lupus erythematosus. *Deutsche Zeitschrift für Nervenheilkunde* 1963; **185**: 244–51.
- 21 Poch G F. *Enfermedades del Colageno. Manifestations neurologicas, musculares y psiquiatricas*. Buenos Aires: Lopez Libreros Editores SRL, 1960.
- 22 Armas-Cruz R, Harnecker J, Ducach G. Clinical diagnosis of systemic lupus erythematosus. *Am J Med* 1958; **25**: 409–19.
- 23 Dubois E L. The effect of the LE cell test on the clinical picture of systemic lupus erythematosus. *Ann Intern Med* 1953; **38**: 1265–94.
- 24 Feinglass E J, Arnett F C, Dorsch A C, Zizic T M, Stevens M B. Neuropsychiatric manifestations of systemic lupus erythematosus: diagnosis, clinical spectrum, and relationship to other features of the disease. *Medicine (Baltimore)* 1976; **55**: 323–39.
- 25 Goldman M, Herode A, Borenstein S, Zonen A. Optic neuritis, transverse myelitis and anti-DNA antibodies nine years after thymectomy for myasthenia gravis. *Arthritis Rheum* 1984; **27**: 701–3.
- 26 Small P, Mass M F, Kohler P, Harbeck R J. Central nervous system in SLE—diagnostic profile and clinical features. *Arthritis Rheum* 1977; **20**: 869–78.
- 27 Sergeant J S, Locksin M D, Klempner M S. Central nervous system disease in systemic lupus erythematosus: therapy and prognosis. *Am J Med* 1975; **58**: 645–54.
- 28 Gibson T, Myers A R. Nervous system involvement in systemic lupus erythematosus. *Ann Rheum Dis* 1976; **35**: 398–406.
- 29 Clinicopathological conference. *Bulletin of the Johns Hopkins Hospital* 1966; **118**: 423–37.
- 30 Granger D P. Transverse myelitis with recovery: the only manifestation of systemic lupus erythematosus. *Neurology* 1960; **10**: 325–9.
- 31 Duffy J, Andrianakos A, Lidsky M. Low spinal fluid glucose in systemic lupus with spinal cord disease [Abstract]. *Clin Res* 1974; **22**: 610.
- 32 Lipton H L, Teasdale R D. Acute transverse myelopathy in adults. *Arch Neurol* 1978; **4**: 51–9.