SCIENTIFIC PAPERS

Autoantibodies in patients with juvenile chronic arthritis and their immediate family relatives

T R Southwood, P J Roberts-Thomson, M J Ahern, K Shepherd, R McEvoy, J B Ziegler, J Edmonds

Abstract

Antibodies to nuclear antigens were assessed in 23 children with juvenile chronic arthritis (JCA) and 66 of their first degree relatives. Serum samples from 16 patients with JCA (70%) and nine relatives (14%) had antinuclear antibodies by indirect immunofluorescence. Antibodies against nuclear antigens in rabbit thymus extract or an erythroblastoid cell line (K562) were detected by counterimmunoelectrophoresis and immunoblotting in 16 patients (70%) and 39 family relatives (59%). Immunoblotting did not show any banding patterns common to all patients with JCA, though bands in the 43-45 kD range were detected in 5/23 patients. Anticardiolipin antibodies were found in 7/23 patients. In total, 18/20 families (90%) had members other than the probands with detectable autoantibodies. In five families immunoblotting showed common banding patterns between the probands and other members. This suggests that there might be an inherited trend towards autoimmune responses in some families of patients with JCA.

The pathogenesis of juvenile chronic arthritis (JCA) is poorly understood. The presence of antinuclear antibodies,¹⁻⁴ anticollagen antibodies,⁵⁻⁸ anti-T-cell antibodies,⁹⁻¹¹ rheumatoid factors,¹²⁻¹⁴ and other autoantibodies^{15 16} in patients with JCA suggests that an autoimmune process is taking place. In other autoimmune diseases, such as systemic lupus erythematosus and progressive systemic sclerosis, there seems to be an inherited tendency towards autoantibody formation,¹⁷⁻²¹ but, as far as we are aware, this has not been previously investigated in JCA.

Studies of familial patterns and HLA associations²²⁻²⁹ have provided evidence for a genetic predisposition to JCA. There also seems to be an increased prevalence of autoimmune disease in the families of patients with JCA,³⁰ but the prevalence of autoantibodies in these families has not been reported. In this study our first aim was to report the prevalence of antinuclear antibodies among 66 first degree relatives of 20 patients with JCA, and to determine if there were common patterns of these antibodies among family members.

The antigenic specificities of most antinuclear antibodies associated with JCA are undetermined, though there is evidence that some of the antinuclear antibodies are directed against histones.^{31 32} Our second aim was to clarify some of the cellular constituents to which

autoantibodies are directed in patients with JCA by immunoblotting techniques. Our third aim was to investigate antibodies to cardiolipin in our patients with JCA by an enzyme linked immunosorbent assay (ELISA). This family of autoantibodies has a close clinical association with the presence of antinuclear antibodies.

Patients and methods

Twenty three patients with juvenile chronic arthritis³³ were arbitrarily selected from patients attending paediatric arthritis clinics. Eighteen children had pauciarticular onset JCA (14 girls, of whom nine also had chronic anterior uveitis, and four boys, only one of whom had chronic anterior uveitis). One girl had systemic onset disease and four children had polyarticular onset (three girls and one boy); none of these five patients had chronic anterior uveitis. Four patients were HLA-B27 positive (three boys and one girl with pauciarticular onset disease). All patients had had active disease for at least one year before the study, and most were taking non-steroidal anti-inflammatory drugs; several were also taking slow acting antirheumatic drugs (gold or penicillamine).

Serum samples from 66 immediate family members (20 families) of 20 patients with JCA were also studied. Samples were not available from any family members of the other three patients with JCA. Only one of the family members had an overt autoimmune disease (rheumatoid arthritis in an older sister of a patient). Blood was obtained by antecubital venepuncture in all instances, and the serum samples were stored in small aliquots at -70°C.

Antinuclear antibodies were measured by indirect immunofluorescence on HEp-2 cell line substrates by standard techniques.³⁴ All serum samples were screened at a dilution of 1/40 and a positive reaction recorded if nuclear fluorescence was seen at this dilution. Antibodies to extractable nuclear antigens were measured by counterimmunoelectrophoresis in 1% agarose using either rabbit thymus extract or the cell line K562 to provide nuclear antigenic material. Rheumatoid factor was assessed by latex agglutination (Behring). An ELISA was used to measure anticardiolipin IgG antibodies, as previously described.³⁵

Antinuclear antibodies were characterised by a modification of the immunoblotting technique described by Towbin.³⁶ Briefly, nuclear proteins from rabbit thymus extract or K562 were subjected to sodium dodecyl sulphate/polyacrylamide (12%) gel electrophoresis and the separated proteins transferred to nitrocellulose

Department of
Clinical Immunology,
Flinders Medical Centre,
Bedford Park,
Adelaide,
South Australia,
Australia

T R Southwood
P J Roberts-Thomson
M J Ahern
K Shepherd
R McEvoy

Department of
Immunology, Allergy
and Rheumatology,
University of New
South Wales, Sydney,
Australia
J B Ziegler
J Edmonds

Correspondence to:
Dr T R Southwood,
Department of
Rheumatology, Rheumatism
Research Wing,
The Medical School,
University of Birmingham,
Birmingham B15 2TJ.

Accepted for publication
31 January 1990

Table 1 Prevalence of autoantibodies in families of patients with juvenile chronic arthritis*

	Immunofluorescence†	CIEP‡		Immunoblotting§	
	HEp-2	RTE	K562	RTE	K562
Number of families with one positive¶ member	5	4	7	5	9
Number of families with two positive members	2	2	6	0	3
Number of families with three positive members	0	0	1	0	0
Total No (%) positive of 66 relatives	9 (14)	8 (12)	22 (33)	5 (8)	15 (23)

*Excluding the probands with juvenile chronic arthritis.

†Indirect immunofluorescence using HEp-2 substrate.

‡Counterimmunoelectrophoresis using either RTE or K562 substrate.

§Immunoblotting using either RTE or K562 substrate.

||RTE=rabbit thymus extract.

¶A positive family member had serological evidence of at least one autoantibody.

paper. After blocking with horse serum the nitrocellulose paper was cut into strips according to individual lanes of separated proteins, incubated with patients' serum samples, and washed. Any fixed IgG antibody was visualised with an avidin-biotin, alkaline phosphatase substrate system (Vector).

All the above immunological techniques were standardised and were in regular use in our department. Positive and negative internal controls were included in each assay, and regular participation in an external quality assurance programme in immunology (Royal College of Pathologists of Australasia) ensured close monitoring of the accuracy and precision of the techniques. Fisher's exact test (two tailed) was used for statistical analysis where appropriate.

Results

Eighteen of 20 families had at least one additional member (other than the proband with JCA) with serological evidence of autoimmunity (table 1). In one family four members had positive tests for autoantibodies; three members were positive in five families; two were positive in eight families; and one was positive in four families. In all, 39/66 family members (59%) had one or more positive tests for autoantibodies. No positive tests for autoimmunity in the families could be significantly correlated with positive tests for autoimmunity in the patients.

Serum samples from 16 of the 23 patients with JCA were antinuclear antibody positive in

this study. Nuclear fluorescence was homogeneous in 13 patients and speckled in three. Thirteen antinuclear antibody positive patients had pauciarticular onset disease (12 girls, one boy) and the other three (all girls) had polyarticular onset JCA. Only one of four B27 positive patients had antinuclear antibodies, and she also had antibodies to cardiolipin (table 2). Two female patients with polyarticular onset disease had low titre rheumatoid factor (1/20), and 7/23 patients (30%) had antibodies to cardiolipin.

Fifteen of 23 patients (65%) had antibodies to nuclear antigens by either counterimmunoelectrophoresis or immunoblotting against either rabbit thymus extract or K562 extract. The greatest prevalence of positive reactions (10/23, 43%) was elicited by immunoblotting using K562 (table 3). In only one patient could the antibodies to extractable nuclear antigens be characterised further; he had anti-Sm and later developed systemic lupus erythematosus. None of the other patients had antibodies directed against the nRNP, La, Ro, or ribosomal P antigens. Autoantibodies to nuclear antigens by counterimmunoelectrophoresis were not found in the absence of antinuclear antibodies, and positive reactions by immunoblotting or with cardiolipin were more commonly found in antinuclear antibody positive patients, but the only association which reached statistical significance was antinuclear antibody positivity and positive reactions detected with counterimmunoelectrophoresis using K562 ($p=0.046$).

Precipitin lines were seen on immunoblotting

Table 2 Autoantibody profiles of probands by sex, juvenile chronic arthritis onset type, and presence of chronic anterior uveitis. Figures show the number of patients who were positive for a particular investigation

	Immunofluorescence	CIE*		Immunoblotting		ELISA*
	HEp-2 (16)†	RTE* (4)	K562 (7)	RTE (4)	K562 (10)	ACL* (7)
Female (n=19)	15	4	6	4	8	6
Male (n=4)	1	0	1	0	2	1
Pauciarticular onset (n=18)						
B27 negative (n=14)	12	2	5	4	8	6
B27 positive (n=4)	1	0	0	0	1	1
Polyarticular onset (n=4)						
B27 negative (n=3)	3	2	2	0	1	0
Presence of CAU* (n=10)	8	1	5	3	5	5

*CIE=counterimmunoelectrophoresis; ELISA=enzyme linked immunosorbent assay; RTE=rabbit thymus extract; ACL=anti-cardiolipin antibodies; CAU=chronic anterior uveitis.

†Substrate (number positive) is shown.

Table 3 Comparison of tests for autoantibodies in patients with juvenile chronic arthritis. Figures show the number of patients who were positive for the investigations indicated.†

Test	Substrate	Immu- fluorescence	CIE*		Immunoblotting		ELISA*
		HEp-2 (16)†	RTE* (4)	K562 (7)	RTE (4)	K562 (10)	ACL* (7)
Immunofluorescence CIE	HEp-2		4	7	3	7	6
	RTE			3	1	2	0
	K562				2	3	4
Immunoblotting	RTE					1	1
	K562						4

*CIE=counterimmunoelectrophoresis; ELISA=enzyme linked immunosorbent assay; RTE=rabbit thymus extract; ACL=anti-cardiolipin antibodies.

†Substrate (number positive) is shown.

‡None of the associations reached statistical significance.

in 13 of 23 (57%) patients with JCA. Five of 23 patients (22%) had bands in the 43–45 kD range, and 4/23 (17%) had bands in the 30–35 kD range using K562 substrate (total range of banding 25–50 kD). No common banding patterns were seen on rabbit thymus extract immunoblotting (range of banding 29–80 kD). In total, only three patients (13%) (two of whom were HLA-B27 positive) did not have any evidence of autoantibodies to nuclear antigens or cardiolipin.

Immunoblotting showed common banding patterns between family members and probands with JCA in five families (rabbit thymus extract: one family, 60 kD bands; K562: four families, 33, 35, 37, and 45 kD bands respectively). The nature of the bands was not further characterised. None of these five families had banding patterns in common between the father and the mother. There were also similarities in antinuclear antibody patterns on indirect immunofluorescence (three families), antibodies to rabbit thymus extract (counterimmunoelectrophoresis; three families), and antibodies to K562 (counterimmunoelectrophoresis; five families). Lines of identity were seen between two members of one family on counterimmunoelectrophoresis using K562.

Discussion

The presence of autoantibodies in 39/66 (59%) of the immediate family relatives of patients with JCA, with shared family patterns of banding, immunofluorescence, and precipitation, provides evidence for an inherited basis to the immunoregulatory abnormalities seen in JCA. This finding has not been previously reported as far as we know, though the autoantibodies associated with other autoimmune diseases have been found in asymptomatic family members^{19 21 37 38} and related to the inheritance of certain HLA haplotypes. Anti-Ro, anti-DNA, and rheumatoid factor were associated with HLA-DR2 or HLA-DR3 in patients with systemic lupus erythematosus,¹⁷ and anticentromere antibody was associated with DR1 in patients with progressive systemic sclerosis and their immediate family relatives.²⁰ Environmental factors alone are unlikely to account for the familial prevalence of autoantibodies noted in our study because none of the parental partners had common banding patterns on immunoblotting, including those from the five families who had banding patterns in common with the proband with JCA. In addition,

the prevalence of antinuclear antibodies noted in the families from our study (14%) was much higher than that expected in the general Australian population (3.5%).³⁹ Clemens *et al* reported that HLA-DR5 was associated with antinuclear antibodies in sibling pairs with JCA.²⁸ The subjects in our study did not have full HLA typing. The relation between autoantibodies and the HLA haplotypes associated with the families of children with chronic arthritis remains to be elucidated.

There is other evidence that families of arthritic children are predisposed to autoimmunity. A survey of 888 first degree relatives of 64 probands with juvenile rheumatoid arthritis found a significantly increased prevalence of autoimmune diseases such as systemic lupus erythematosus and adult rheumatoid arthritis.³⁰ Juvenile arthritis was associated with a high prevalence of insulin dependent diabetes mellitus in a large inbred kindred in Canada.²⁴ There have been several reports of concurrence of childhood arthritis among twins,^{25 40} siblings,^{28 29 41} and other family members.²³ The association of juvenile arthritis with HLA-A2, DR5, DRw8, and DPw2^{2 26 27} also provides support for a genetic basis to JCA. It should be noted, however, that multiple sibships with JCA are uncommon⁴² and concordance for juvenile rheumatoid arthritis was found in only 3/10 monozygotic twin pairs,⁴³ suggesting that other precipitating factors play a part.

We found serological evidence of autoimmunity in 20/23 patients (87%). Seventy per cent (16/23) of our patients were antinuclear antibody positive in this study. This percentage was similar to those previously reported in other studies of children with chronic arthritis.^{2–4 8} Comparisons of this nature might be improved if one set of diagnostic/classification criteria for children with arthritis was used instead of the current diverse definitions.^{33 44}

Countercurrent immunoelectrophoresis with rabbit thymus extract showed few positive reactions (4/23), in keeping with results reported by Alspaugh,⁴⁵ Moore,¹³ and Malleon.³² Positive reactions using the K562 substrate were more commonly found (7/23). The use of this substrate to assess autoimmunity in patients with JCA has not been previously reported. Substrates such as calf thymus extract have been shown to be cheaper and superior to rabbit thymus extract in precipitation line resolution, and stability,⁴⁶ and perhaps K562 represents a similar substrate.

Immunoblotting seems to be a more sensitive technique than counterimmunoelectrophoresis for characterising antinuclear antibodies,⁴⁷ but despite using immunoblotting and K562 substrate we were unable to show any banding patterns common to all or even most of our patients. Precipitation lines were most commonly seen in the 43–45 kD (22%) and 30–35 kD (17%) ranges. Using an HEp-2 extract, Malleson *et al* found common IgG precipitation lines at 33/35 and 45 kD (24% and 14% of patients respectively).³² The doublet at 33/35 was thought to represent antihistone H1 reactivity, and we found similar doublets in two of our patients. Antibodies to histone have also been shown in about 50% of serum samples from children with chronic arthritis by immunoblotting,⁴⁸ and by an ELISA technique.³¹ Reactivity to a 70 kD component of a saline extract of beef spleen was reported in 59% of patients with juvenile arthritis,⁴⁹ and more recently the same group reported that 63% of antinuclear antibody positive serum samples from patients with juvenile rheumatoid arthritis react with two bands (50–40 kD) from a saline extract of HeLa cells.⁵⁰ These studies further illustrate the influence of the immunoblotting substrate on the molecular weight of the nuclear antigens recognised by JCA autoantibodies. The detection of common antigenic specificities of antinuclear antibodies associated with JCA may require testing of different nuclear substrates, refinement of immunological techniques, and possibly an improvement of the clinical subclassification of patients with JCA.⁵¹

Anticardiolipin antibodies were detected in 30% (7/23) of our patients, a percentage similar to the 42% (5/12 patients) reported by Shergy *et al*.⁵² The significance of these autoantibodies in JCA is unclear. None of our anticardiolipin antibody positive patients has had any clinically apparent thrombotic episodes, the feature which most commonly links adult patients with antiphospholipid antibodies.^{53–54} Interestingly, 33% of a series of adult patients with rheumatoid arthritis had anticardiolipin antibodies, but no significant correlation with recurrent thrombosis could be found.⁵⁵

Our results have highlighted the occurrence of autoantibodies in other family members of patients with JCA, though we have not been able to define a specific autoantibody common to all of our patients. Future research in this area may increase our understanding of the genetic basis of JCA.

The authors gratefully acknowledge all the patients with juvenile chronic arthritis and their families who contributed blood for this study, the Arthritis Foundation of Australia and the National Health and Medical Research Council for financial assistance, Dr A Hohmann for the results from the anticardiolipin assay, Dr M, MacKinnon of the Statistical Advisory Service of British Columbia's Children's Hospital Research Centre, and B Sharpe and Petra for secretarial assistance.

- 1 Petty R E, Cassidy J T, Sullivan D B. Clinical correlates of antinuclear antibodies in juvenile rheumatoid arthritis. *J Pediatr* 1976; 83: 386–9.
- 2 Leak A M. Autoantibody profile in juvenile chronic arthritis. *Ann Rheum Dis* 1988; 47: 178–82.
- 3 Osborn T G, Patel N J, Moore T L, Zuckner J. Use of the Hep-2 cell substrate in the detection of antinuclear antibodies in juvenile rheumatoid arthritis. *Arthritis Rheum* 1984; 27: 1286–9.
- 4 Rosenberg A M. The clinical associations of antinuclear

- antibodies in juvenile rheumatoid arthritis. *Clin Immunol Immunopathol* 1988; 49: 19–27.
- 5 Rosenberg A M, Hunt D W C, Petty R E. Antibodies to native type I collagen in childhood rheumatic diseases. *J Rheumatol* 1984; 11: 421–4.
- 6 Rosenberg A M, Hunt D W C, Petty R E. Antibodies to native and denatured type II collagen in children with rheumatic diseases. *J Rheumatol* 1984; 11: 425–31.
- 7 Petty R E, Hunt D W C, Rosenberg A M. Antibodies to type IV collagen in rheumatic diseases. *J Rheumatol* 1986; 13: 246–53.
- 8 Haynes D C, Gershwin M E, Robbins D L, Miller III J J, Cosca D. Autoantibody profiles in juvenile arthritis. *J Rheumatol* 1986; 13: 358–63.
- 9 Borel Y, Morimoto C, Cairns L, Mantzouranis E, Strelkauskas A J, Schlossman S F. Anti-T cell antibody in juvenile rheumatoid arthritis. *J Rheumatol* 1984; 11: 56–61.
- 10 Barron K S, Lewis D E, Brewer E J, Marcus D M, Shearer W T. Cytotoxic anti-T cell antibodies in children with juvenile rheumatoid arthritis. *Arthritis Rheum* 1984; 27: 1272–80.
- 11 Morimoto C, Reinhertz E L, Borel Y, Mantzouranis E, Steinberg A D, Schlossman S F. Autoantibody to an immunoregulatory inducer population in patients with juvenile rheumatoid arthritis. *J Clin Invest* 1981; 67: 753–61.
- 12 Bianco N E, Panush R S, Stillman J S, Schur P H. Immunologic studies in juvenile rheumatoid arthritis. *Arthritis Rheum* 1971; 14: 685–96.
- 13 Moore T L, Osborn T G, Weiss T D, *et al*. Autoantibodies in juvenile arthritis. *Semin Arthritis Rheum* 1984; 13: 329–36.
- 14 Schlumpf U, Howard A, Ansell B M. IgG-Anti-IgG antibodies in juvenile chronic arthritis. *Scand J Rheumatol* 1985; 14: 65–8.
- 15 Petty R E, Hunt D W C, Rollins D F, Schroeder M-L, Puterman M L. Immunity to soluble retinal antigen in patients with uveitis accompanying juvenile rheumatoid arthritis. *Arthritis Rheum* 1987; 30: 287–93.
- 16 Moore T L, Dorner R W. 19S IgM Forssman-type heterophile antibodies in juvenile rheumatoid arthritis. *Arthritis Rheum* 1980; 23: 1262–7.
- 17 Ahearn J M, Provost T T, Dorsch C A, Stevens M B, Bias W B, Arnett F C. Interrelationships of HLA-DR, MB, and MT phenotypes, autoantibody expression and clinical features in systemic lupus erythematosus. *Arthritis Rheum* 1982; 25: 1031–40.
- 18 Teitsson I, Thorsteinsson J, Arnason A, Valdimarsson H. Rheumatic diseases in an Icelandic family. *Scand J Rheumatol* 1985; 14: 109–18.
- 19 Adamashvili I M, Makarova O V, Kvachadze I M, Trubnikov V I, Zamchuk L A. Antibodies to nucleic acids in systemic lupus erythematosus and rheumatoid arthritis patients and their relatives. *Immunol Lett* 1983; 6: 33–8.
- 20 Ruffarri A, Artifone L, Glorioso S, *et al*. Prevalence of antientromere antibody in blood relatives of antientromere positive patients. *J Rheumatol* 1985; 12: 940–2.
- 21 Lipman S M, Arnett F C, Conley C L, Ness P M, Meyers D A, Bias W B. Genetic factors predisposing to autoimmune diseases: autoimmune hemolytic anemia, chronic thrombocytopenic purpura, and systemic lupus erythematosus. *Am J Med* 1982; 73: 827–40.
- 22 Howard J F, Sigbee A, Glass D N. HLA genetics and inherited predisposition of JRA. *J Rheumatol* 1985; 12: 7–12.
- 23 Rossen R D, Brewer E J, Sharp R M, Ott J, Templeton J W. Familial rheumatoid arthritis: linkage of HLA to disease susceptibility locus in four families where proband presented with juvenile rheumatoid arthritis. *J Clin Invest* 1980; 65: 629–42.
- 24 Jaworski M A, Slater J D, Severini A, *et al*. Unusual clustering of diseases in a Canadian old colony (Choritz) Mennonite kindred and community. *Can Med Assoc J* 1988; 138: 1017–25.
- 25 Aho K, Koshenvuo M, Tuominen J, Kaprio J. Occurrence of rheumatoid arthritis in a nationwide series of twins. *J Rheumatol* 1986; 13: 899–902.
- 26 Hoffman R W, Shaw S, Francis L C, *et al*. HLA-DP antigens in patients with pauciarticular juvenile rheumatoid arthritis. *Arthritis Rheum* 1986; 29: 1057–62.
- 27 Oen K, Petty R E, Schroeder M-L. An association between HLA-A2 and juvenile rheumatoid arthritis in girls. *J Rheumatol* 1982; 9: 916–20.
- 28 Clemens L E, Albert E, Ansell B M. Sibling pairs affected by chronic arthritis of childhood: evidence for a genetic predisposition. *J Rheumatol* 1985; 12: 108–13.
- 29 Rosenberg A M, Petty R E. Similar patterns of juvenile rheumatoid arthritis within families. *Arthritis Rheum* 1980; 23: 951–3.
- 30 Church J L, Bernstein B, Church J A, Hanson V. Familial prevalence of immunologically-mediated diseases (IMD) in juvenile rheumatoid arthritis (JRA) [Abstract]. *Arthritis Rheum* 1983; 26 (suppl): S85.
- 31 Ostensen M, Fredriksen K, Kass E, Rekvig O-P. Identification of antihistone antibodies in subsets of juvenile chronic arthritis. *Ann Rheum Dis* 1989; 48: 114–7.
- 32 Malleson P N, Petty R E, Fung M, Candido E P M. Reactivity of antinuclear antibodies with histones and other antigens in juvenile rheumatoid arthritis. *Arthritis Rheum* 1989; 32: 990–4.
- 33 Wood P H. Nomenclature and classification of arthritis in children. In: Munthe E, ed. *The care of rheumatic children*. Basel: EULAR, 1978: 47–50.
- 34 Friou G J. Fluorescent spot test for anti-nuclear antibodies. *Arthritis Rheum* 1962; 5: 407–10.
- 35 Sutjita M, Hohmann A, Comacchio R, Bradley J. Poly-

- specific human and murine antibodies to diphtheria and tetanus toxoids and phospholipids. *Clin Exp Immunol* 1988; 73: 191-7.
- 36 Towbin H, Gordon J. Immunoblotting and dot immunobinding—current status and outlook. *J Immunol Methods* 1984; 72: 313-40.
  - 37 Aho K, Sistonen P, Takala J, Sievers K. Genetics of autoantibodies in relation to disease. An epidemiological study of two population series. *Acta Med Scand* 1982; 211: 213-8.
  - 38 Fennell R H Jr, MacLachlan M J, Rodnan G P. The occurrence of antinuclear factors in the sera of relatives of patients with systemic rheumatic disease. *Arthritis Rheum* 1962; 5 (suppl): 296.
  - 39 Hawkins R B, O'Connor K J, Dawkins R L, et al. Autoantibodies in an Australian population. 1. Prevalence and persistence. *J Clin Lab Immunol* 1979; 2: 211-5.
  - 40 Husby G, Williams R C, Tung K S, et al. Immunologic studies in identical twins concordant for juvenile rheumatoid arthritis but discordant for monoclonal gammopathy and amyloidosis. *J Lab Clin Med* 1988; 111: 307-14.
  - 41 Fujita K, Kobayashi K, Okino F. Juvenile rheumatoid arthritis in two siblings with congenital leucocyte adhesion deficiency. *Eur J Pediatr* 1988; 148: 118-9.
  - 42 Ansell B M. Juvenile chronic arthritis-families. *Dis Markers* 1986; 4: 163.
  - 43 Baum J, Fink C. Juvenile rheumatoid arthritis in monozygotic twins: a case report and review of the literature. *Arthritis Rheum* 1968; 11: 33-6.
  - 44 Cassidy J T, Levinson J E, Bass J C, et al. A study of classification criteria for a diagnosis of juvenile rheumatoid arthritis. *Arthritis Rheum* 1986; 29: 274-81.
  - 45 Alsbaugh M A, Miller III J J. A study of specificities of antinuclear antibodies in juvenile rheumatoid arthritis. *J Pediatr* 1977; 90: 391-5.
  - 46 Collins R J, Neil J C, Druery L N, Wilson R J. Detection of antibodies to extractable nuclear antigens using calf thymus and rabbit thymus. A comparative study of 1000 consecutive antinuclear antibody positive patients. *J Immunol Methods* 1989; 116: 53-7.
  - 47 Westgeest A A A, van den Brink H G, de Jong J, Swaak A J G, Smeenk R J T. Routine testing for antinuclear antibodies: a comparison of immunofluorescence, counter immunoelectrophoresis and immunoblotting. *Journal of Autoimmunity* 1988; 1: 159-70.
  - 48 Pauls J D, Silverman E, Laxer R M, Fritzler M J. Antibodies to histones H1 and H5 in sera of patients with juvenile rheumatoid arthritis. *Arthritis Rheum* 1989; 32: 877-83.
  - 49 Osborn T G, Smith K R, Haber P L, Moore T L. Characterisation of a saline extractable antigen reactive with juvenile rheumatoid arthritis sera [Abstract]. *Arthritis Rheum* 1987; 30 (suppl): 126.
  - 50 Haber P L, Osborn T G, Moore T L. Antinuclear antibody in juvenile rheumatoid arthritis sera reacts with 50-40 kDa antigen(s) found in HeLa nuclear extracts. *J Rheumatol* 1989; 16: 949-54.
  - 51 Southwood T R, Petty R E, Malleson P N, et al. Psoriatic arthritis in children. *Arthritis Rheum* 1989; 32: 1007-13.
  - 52 Shergy W J, Kredich D W, Pisetsky D B. The relationship of anticardiolipin antibodies to disease manifestations in pediatric systemic lupus erythematosus. *J Rheumatol* 1988; 15: 1389-94.
  - 53 Hughes G R V, Asherson R A, Khamashta M A. Antiphospholipid syndrome: linking many specialties. *Ann Rheum Dis* 1989; 49: 355-6.
  - 54 Alarcon-Segovia D. Pathogenetic potential of antiphospholipid antibodies. *J Rheumatol* 1988; 15: 890-3.
  - 55 Fort J G, Cowchock F S, Abruzzo J L, Smith J B. Anticardiolipin antibodies in patients with rheumatic diseases. *Arthritis Rheum* 1987; 30: 752-60.