DISPATCH

From India

V R Joshi

Annual conference

When Chandigarh was chosen as the venue for the annual conference (1989) many had misgivings about its suitability. Normally, Chandigarh, the first planned modern city of India, is a tourist attraction, but for the medical community there is the added inducement of visiting the postgraduate institute. The prevailing political atmosphere kept many away, but it was a pleasant surprise to see that Chandigarh itself was quiet and peaceful. Conference venues are often hotels, but this conference took place on the postgraduate institute campus, permitting interaction between those at the conference and faculty members. This institute has excellent conference facilities for up to 200 participants.

The conference drew attention to certain rare diseases. The Chandigarh group led by Professor D S Deodhar presented data on Wegener's disease, a condition not often diagnosed and, possibly, sometimes missed. A similar contribution was made by Dr Malaviya's group from Delhi, on Behçet's disease. It seems that with greater awareness, and ability to diagnose, more of these rare but important conditions will come to light. Lyme disease may be one such condition—one case was reported recently.

The conference the following year (1990) was held in the historic city of Hyderabad. Details will be given in the next dispatch.

Tuberculosis and rheumatology

Tuberculosis poses many challenges to a rheumatologist. Being a common disease it can never be ignored. Owing to a lack of bacteriological facilities diagnosis is often based not on demonstration of acid fast bacilli, but on the clinical picture and the histopathological appearance. If tuberculosis is suspected antituberculous drugs are given and the therapeutic results observed. The real dilemma arises when the tuberculosis is polyarticular when diagnosis is difficult and often delayed.

The following case report illustrates this point. A female patient had arthritis in her right foot, which slowly spread over one to two years to her right knee, left elbow, and left wrist. She was diagnosed and treated for rheumatoid arthritis until a discharging sinus developed in her elbow.

An excision biopsy specimen from her elbow and wrist was consistent with tuberculosis. Acid fast bacilli could not be shown. Standard treatment was given and she is improving.

Another variant of the disease is disseminated haematogenous tuberculosis of bones. A natural differential diagnosis is that of metastatic lesions. The availability of a bone scan has shown the presence of multiple lesions in more than one case seen by me and by others. Apparently, articular and osseous tuberculosis is often a multifocal disorder.

Tuberculosis is an ever present danger to patients receiving corticosteroids and immuno-suppressive treatment. There is no consensus on the use of chemoprophylaxis for tuberculosis. Should all such patients receive it? If, yes, which drugs and for how long? These questions need to be answered. One study on renal transplant recipients from Vellore (India) concluded that chemoprophylaxis is of no help for such patients. Whether the same is true for patients with rheumatic disorders is not clear.

Atypical tubercle bacilli pose a problem as illustrated by the following case. A female patient with systemic lupus erythematosus, who had received steroids for a long time, developed fever and nodules on her legs and, subsequently, on the abdominal wall. Histological and bacteriological tests of the nodule at an early stage were inconclusive. She was given further steroids and even cyclophosphamide. Eventually, one such lesion softened and examination confirmed *Mycobacterium chelonii* infection, which responded to treatment with appropriate chemotherapeutic agents.

Socioeconomic status and disease patterns

Interesting differences in disease patterns are apparent among the poor and the affluent. Ulcerative colitis, until recently thought to be uncommon, is diagnosed quite often among the more affluent. Interestingly, articular manifestations are as common as in the West.

In contrast, in public hospitals (in Bombay), where most of the patients come from the poorer section of the community, seronegative (spondyl-)arthritis, the reactive variety, is as prevalent as rheumatoid arthritis. Ulcerative colitis is seen here, but less commonly, and arthritis secondary to inflammatory bowel disease is distinctly uncommon.

In a private hospital, drawing patients from the affluent section, rheumatoid arthritis is the most common inflammatory arthritis, but extra-articular features are uncommon. The overall pattern of disease here is thus more like that in the West. It would be interesting to know whether such a pattern exists in other developing countries.

P D Hinduja National
Hospital and Medical
Research Centre,
Veer Savarkar
Marg, Mahim,
Bombay 400 016, India
V R Joshi