Plasma von Willebrand factor antigen concentrations. Each point represents one patient or control. The bar is the median value.

of a higher von Willebrand factor antigen in patients with diffuse disease suggests endothelial cell damage may be more extensive and active in this group. The extent of vascular injury as shown by von Willebrand factor antigen concentrations may be regarded as a prognostic marker in these disorders.

A D BLANN
Department of Surgery
Research and Teaching Building
University Hospital of South Manchester
Nell Lane, Didsbury
Manchester M20 8LR, UK

KAREN J ILLINGWORTH
M I V JAYSON
University of Manchester Rheumatic Diseases Centre
Clinical Sciences Building
Hope Hospital
Eccles Old Road
Salford, Manchester M6 8HD, UK

- Belch J J F, Zoma A A, Richards I M, McLaughlin K, Forbes C D, Sturrock R D. Vascular damage and factor-VIII-related antigen in the rheumatic diseases. *Rheumatol Int* 1987; 7: 107-11.
- Kahaleh M B, Osborn I, Leroy E C. Increased factor VIII/von Willebrand factor antigen and von Willebrand factor activity in scleroderma and in Raynaud's phenomenon. *Ann Intern Med* 1981; 94: 482-4.
- Sheeran T P, Blann A D, Emery P, Bacon P A. Factor VIII:Rag in primary Raynaud's syndrome and scleroderma. Is it of diagnostic or prognostic value? *Arthritis Rheum* 1990; 33 (suppl): S157.
- Greaves M, Malia R G, Milford Ward A, et al. Elevated von Willebrand factor antigen in systemic sclerosis: relationship to visceral disease. *Br J Rheumatol* 1988; 27: 281-5.
- LeRoy E, Black C, Fleischmajer R, et al. Scleroderma (systemic sclerosis): classification, subsets and pathogenesis. *J Rheumatol* 1988; 15: 202-5.
- Short P E, Williams C E, Picken A M, Hill F G H. Factor VIII related antigen: an improved enzyme immunoassay. *Med Lab Sci* 1982; 39: 351-5.

Anticardiolipin antibodies in drug addicted patients with AIDS

Sir: The presence of anticardiolipin antibodies and lupus anticoagulant has been described in up to 90% of homosexual patients with AIDS.^{1 2} There have been some attempts to

relate anticardiolipin antibodies with *Pneumocystis carinii* infection,³ other opportunistic infections, and with neoplasm,⁴ but up to the present, anticardiolipin antibodies in patients with AIDS have only been associated with a poor outcome of the illness.^{5 6} In these patients anticardiolipin antibodies do not have the same clinical significance as they have in systemic lupus erythematosus.¹

As drug addicted patients with AIDS have a different spectrum of rheumatic manifestations, with less reactive arthritis, Reiter's syndrome, and HIV related arthritis, and more septic arthritis than homosexual patients with AIDS (Monteagudo I *et al*, unpublished data), we studied the presence of anticardiolipin antibody in drug addicted patients with AIDS and its relation with clinical manifestations.

Anticardiolipin antibody was determined (Cheshire Diagnostic QACA enzyme linked immunosorbent assay (ELISA) kit) in 55 drug addicted patients. Forty three of these had been admitted to hospital because of acute problems related to AIDS (mainly infections) and the remaining 12, without acute problems, were seen at the outpatient AIDS clinic in the hospital. Anticardiolipin antibodies were positive in 42 (76%)—31/43 (72%) of the hospital patients and 11/12 (92%) of the outpatients. Clinical problems in the hospital patients were diverse, and we found no significant relation between infection and the presence of anticardiolipin antibodies. Additionally, there was no relation between anticardiolipin antibodies and the antecedent and kind of infection in these patients. Of the 55 patients, 43 were in stage IV AIDS, with 34 (79%) being positive for anticardiolipin antibodies. The remainder were in other stages and anticardiolipin antibodies were found in 10/12 (83%).

Our results show that the presence of anticardiolipin antibodies in drug addicted patients with AIDS is similar to that found in homosexual patients with AIDS. We found no relation between anticardiolipin antibodies and clinical manifestations (mainly infections) nor pathological antecedents in these patients. Similarly, there were no differences between outpatients and hospital patients. As with homosexual patients with AIDS, we found no correlation between anticardiolipin antibodies and recurrent thrombosis and thrombopenia. Probably, anticardiolipin antibodies are associated with HIV infection itself or with an abnormal immune response which is not yet well defined.

JAVIER RIVERA
INDALECIO MONTEAGUDO
JAVIER LOPEZ-LONGO
EDUARDO MALDONADO
LUIS CARREÑO
Servicio de Reumatología
Hospital General Gregorio Marañón
Madrid, Spain

Correspondence to: Dr J Rivera, Servicio de Reumatología, Hospital Gregorio Marañón, Doctor Esquerdo 46, 28007 Madrid, Spain.

- Canoso R T, Zon L I, Groopman J E. Anticardiolipin antibodies associated with HTLV-III infection. *Br J Haematol* 1987; 65: 495-8.
- Mizutani W T, Woods V L, McCutchan J A, Zvaifler N J. Anticardiolipin antibodies in human immunodeficiency virus (HIV) infected gay males may be associated with a deteriorating clinical course. *Arthritis Rheum* 1987; 30: S35.

Disabling ossification of the patellar tendon

Sir: A 42 year old lorry driver without any medical or traumatic history had had for two years pains in the right knee when walking. They had appeared simultaneously with the use of a new truck, in which the accelerator pedal was particularly stiff. After some months of limping clinical examination showed that the right patellar tendon was diffusely thickened and tender; a bulge sign was found, as well as atrophy of the quadriceps muscle and a marked reduction of the knee flexion. Erythrocyte sedimentation rate was 1 mm/1 st h; fasting blood sugar, calcium, phosphorus, and alkaline phosphatase were normal. Synovial fluid contained 0.1×10^9 leucocytes/l without crystals.

Lateral plain radiographs and tomograms (without signs of patella alta: length of the patellar tendon equal to the diagonal length of the patella¹) showed ossification of the tendon, which did not affect its distal third (fig 1). Sonography showed that ossification was mostly in the lateral part and that the non-ossified distal third was thickened in comparison with the tendon on the left side (anteroposterior thickness 9 mm *v* 6 mm). Radiographs and computed tomography also showed irregular enthesitic osteophytes of the upper part of patellae on both knees and, in addition, a femoral and tibial osteopoikilosis. Radiographs of cervical, thoracic, and lumbar spine showed no abnormalities.

Arthroscopic examination showed the joint cavity was normal. During operation the orthopaedic surgeon found that the right patellar tendon was wider than normal. The ossified mass was not adherent to the adjacent patella or tibia and could be easily dissected and removed. The patient recovered slowly and was able to resume his work three months later.

Pathological examination showed compact remodelled lamellar bone. In the mass of the bone, and on the proximal end, strips of fibrous cartilage were seen, which might be considered as an abnormal metaplasia of the tendon (figs 2A and B). Despite its distance from any insertion site this intermingling of bone and fibrous cartilage resembles that seen in enthesopathic hyperostosis—that is, in a location in which fibrocartilaginous bundles are normally found.² On the distal end tendinous bundles intermingled with scar tissue associated with some degree of scar

Figure 1 (A) Plain radiograph; (B) tomogram.

- Stimmler M M, Quismorio F P, McGehee W G, Boylen T, Sharma O P. Anticardiolipin antibodies in acquired immunodeficiency syndrome. *Arch Intern Med* 1989; 149: 1833-5.
- Bloom E J, Abrams D I, Rodgers G. Lupus anticoagulant in the acquired immunodeficiency syndrome. *JAMA* 1986; 256: 491-3.
- Gold J E, Haubenstock A, Zalusky R. Lupus anticoagulant and AIDS. *N Engl J Med* 1986; 314: 1252-3.
- Cohen A J, Phillips T E, Kessler G M. Circulating coagulation inhibitors in the acquired immunodeficiency syndrome. *Ann Intern Med* 1986; 104: 175-80.