DISPATCH

From India

V R Joshi

The Indian Rheumatism Association holds elections every two years. In 1991 Professor A N Maliviya, AIIMS, New Delhi, was elected as president for the years 1992–1993.

The annual conference of the Indian Rheumatism Association was held at Lucknow from 17–19 December 1991. The conference had its problems owing to a break down in air services. Nevertheless, Dr S K Das, the organising secretary, did an excellent job. Some of the conference highlights are given below.

Handigodu disease

Professor S S Agarwal gave a lecture on Handigodu disease based on his observations as a part of an ICMR study project. This disease has attracted a lot of attention. It was first reported as endemic familial arthritis of Maland by Bhat and Krishnamachari; it was not noted before 1965. Initially, over 400 cases were reported, most coming from only two villages, though some 35 villages had these cases. Now only sporadic cases are being reported. The disease is characterised by back and hip pain, leading to flexion deformity and a prominent lordotic gait. There is no evidence of growth retardation. Radiologically there is osteoarthritic change affecting vertebrae and hips. Initial observations implicated environmental factors as three generations were affected 'simultaneously'. The incriminated environmental factors included pesticides and heavy metals (cadmium).

Professor Agarwal thinks otherwise. He considers it to be a genetic disorder because the disease appears to affect members of the Harijan caste; the affected villages are widely separated;

presence of familial aggregation, often inter-related parents, and evidence of skeletal abnormalities in other family members. Professor Agarwal has noted two other skeletal abnormalities in the same group: (a) women with long arms and flat vertebral bodies and (b) dwarfs. Further, he notes that men and women are equally affected; one of the parents is affected in most cases, and all the three skeletal abnormalities can be seen in one family. Dwarfs are often born when both parents are affected. He postulates that it is an autosomal dominant disorder or metaphysal dysplasia with variable, age modified, expression. The low calcium and high fluorine intake noted may act as permissive factors.

SSA–rheumatoid arthritis overlap

We must all occasionally have come across patients who seem to have SSA–rheumatoid arthritis overlap, Dr Maliviya's group presented their observations on 14 such cases. Briefly, the disease is seen commonly in male patients who have ankylosing spondylitis. Over the years they develop peripheral arthritis resembling rheumatoid arthritis clinically and radiologically (7/14 showed erosions). Rheumatoid factors were detected in 5/14, B27 in 10/14, DR4 in 10/14, while both DR4 and B27 were present in 7/14.

It seems that the disease is basically an SSA (ankylosing spondylitis). What leads to peripheral arthritis resembling rheumatoid arthritis is not clear. Does DR4 have some role? Study of these cases is important as it may offer some insights into disease mechanisms.

Other topics

The Boots oration was delivered by Dr Nirupa E Borges on 'A decade with seronegative spondarthritides'.

Epidemiology and standardisation of databases for patient evaluation and outcome assessment were also discussed. In India work is beginning on both these important aspects of rheumatology.

P D Hinduja National
Hospital and Medical
Research Centre,
Veer Savarkar Marg,
Mahim, Bombay 400 016,
India
V R Joshi