baby girl was delivered (650 g; small for date: Apgar 4). Examination of the placenta disclosed focal decidual infarctions. The mother progressed well after birth, but although the baby was intensively treated, she died within a month owing to complications associated with rupture of the intestine.

Anticardiolipin antibody (IgG) was retrospectively determined by ELISA with or without the addition of β_2 -GPI to the ELISA system.²⁻⁴ The concentration of ACA with β_2 -GPI increased from the 20th week of gestation when retardation of placental growth was noticed and the TAT concentration showed an extraordinary increase, whereas the level of ACA without β_2 -GPI remained within the normal range (figure). Furthermore, the level of thrombomodulin, which has been reported as a marker for endothelial cell injury,⁸ increased from the 24th week of gestation. There were no remarkable changes in other serological data, except circulating immune complexes, which increased transiently during the 20th week of gestation.

Thus ACA must be determined with the addition of β_2 -GPI. Furthermore, serial examination of TAT or thrombomodulin concentrations, or both, will provide additional information as to whether subclinical thrombotic conditions exist or not because ACA is not always associated with a pathological role⁹ and nor does a marked increase of ACA always induce thrombosis.¹⁰

S KOBAYASHI
M TANAKA
H TSUDA
H HASHIMOTO
S HIROSE

Department of Rheumatology
Juntendo University School of Medicine
Hongo, Bunkyo-ku
Tokyo 113, Japan

T SAIKAWA
K YOSHIDA
Department of Obstetrics and Gynecology
Juntendo University School of Medicine
Tokyo 113, Japan

Correspondence to: Dr Shigeto Kobayashi, Department of Rheumatology, Juntendo University School of Medicine, 2-1-1 Hongo, Bunkyo-ku, Tokyo 113, Japan.

- McNeil P H, Simpson R J, Chesterman C N, Krilis S A. Anti-phospholipid antibodies are directed against a complex antigen that includes a lipid-binding inhibitor of coagulation: β_2 -glycoprotein I (apolipoprotein H). *Proc Natl Acad Sci USA* 1990; 87: 4120-4.
- Matsuura E, Igarashi M, Igarashi Y, et al. Molecular definition of human β_2 -glycoprotein I (β_2 -GPI) by cDNA cloning and inter-species differences of β_2 -GPI in alternation of anticardiolipin binding. *International Immunology* 1991; 3: 1217-21.
- Chamley L W, McKay E J, Pattison N S. Co-factor dependent and co-factor independent anticardiolipin antibodies. *Thromb Res* 1991; 61: 291-9.
- Matsuura E, Igarashi Y, Fujimoto M, Ichikawa K, Koike T. Anticardiolipin co-factor(s) and differential diagnosis of autoimmune disease. *Lancet* 1990; 336: 177-8.
- Harris E N, Gharavi A E, Patel P, Hughes G R V. Evaluation of the anti-cardiolipin antibody test: report of an international workshop held on 4 April 1986. *Clin Exp Immunol* 1987; 68: 215-22.
- Kobayashi S, Tamura N, Tsuda H, Mokuno C, Hashimoto H, Hirose S. Immunoabsorbent plasmapheresis for a patient with antiphospholipid syndrome during pregnancy. *Ann Rheum Dis* 1992; 51: 399-401.
- Pelzer H, Schwartz A, Heimburger N. Determination of human thrombin-antithrombin III complex in plasma with an enzyme-linked immunosorbent assay. *Thromb Haemost* 1988; 59: 101-6.
- Ishii H, Nakano M, Tsubouchi J, et al. Establishment of enzyme immunoassay of human thrombomodulin in plasma and urine using monoclonal antibodies. *Thromb Haemost* 1990; 63: 157-62.

- Caporali R, Ravelli A, De Gennaro F, Montecucco C, Martini A. Prevalence of anticardiolipin antibodies in juvenile chronic arthritis. *Ann Rheum Dis* 1991; 50: 599-601.
- Asherson R A, Baguly E, Pal C, Hughes G R. Antiphospholipid syndrome: five year follow up. *Ann Rheum Dis* 1991; 50: 805-10.

Sulphoxidation status in rheumatoid arthritis

SIR: In a recent issue of this journal Emery *et al*¹ found an increased prevalence of poor sulphoxidation in patients with rheumatoid arthritis (RA). Using the same methods as Emery's group, we have found similar results.

Sulphoxidation status was assessed in 116 of 200 patients enrolled in a comparative study of sulphasalazine and D-penicillamine as second line drugs in RA (table). Follow up data are available on these patients for between 5 to 7 years from entry into the trial. Surprisingly, no increase in toxicity from either drug was observed in the poor sulphoxidisers group (figure).

Previous studies have suggested that toxicity from intramuscular gold² and D-penicillamine³ is related to sulphoxidation status. In contrast with gold and D-penicillamine, however, sulphasalazine is metabolised predominantly by acetylation rather than sulphoxidation, and one would therefore have predicted that toxicity from sulphasalazine would not be influenced by sulphoxidation status. In this study we failed to find any association between toxicity from D-penicillamine and sulphoxidation status. Although the period of follow up in this study was longer than in previous studies, this finding cannot be explained by an excess of late toxicity in the good sulphoxidisers group.

A second possible explanation is that the group in whom sulphoxidation status was measured was not representative of the group as a whole. To consider this point we noted toxicity in those patients in whom sulphoxidation status was not assayed. Of the patients receiving sulphasalazine in whom sulphoxida-

tion status was known, 35% developed toxicity compared with 30% of those in whom sulphoxidation status was not known. In those receiving D-penicillamine the figures were 34% and 38% respectively. Thus although there was a small excess in toxicity in those receiving D-penicillamine who did not have sulphoxidation status assayed, this was not statistically significant (χ^2). We cannot rule out the possibility that the ratio of good and poor sulphoxidisers might have been different in those who were not assayed. There were 12 deaths (five sulphasalazine, seven D-penicillamine) in the group where sulphoxidation status could not be evaluated compared with two in the group where sulphoxidation status was assayed (one sulphasalazine, one D-penicillamine). None of these deaths was directly attributable to drug toxicity.

The consistent finding that poor sulphoxidisers are overrepresented in RA compared with the normal population is of great interest, and the effect of sulphoxidation status on susceptibility to RA or disease expression, or both, merits further study.

E A MURPHY
R MADHOK
H A CAPELL
Centre for Rheumatic Diseases
Glasgow Royal Infirmary
Glasgow G4 0SF
United Kingdom

R WARING
Department of Biochemistry
University of Birmingham
Birmingham
United Kingdom

J A HUNTER
Gartnavel General Hospital
Glasgow G12
United Kingdom

Correspondence to: Dr Elizabeth Murphy, University Department of Medicine, Glasgow Royal Infirmary, Glasgow G3 7ER, United Kingdom.

- Emery P, Bradley H, Gough A, Arthur V, Jubb R, Waring R. Increased prevalence of poor sulphoxidation in patients with rheumatoid arthritis: effect of changes in the acute phase response and second line drug treatment. *Ann Rheum Dis* 1992; 51: 318-20.
- Ayesh R, Mitchell S C, Waring R H, Withington R H, Seifert M H, Smith R L. Sodium aurothiomalate toxicity and sulphoxidation capacity in rheumatoid arthritis patients. *Br J Rheumatol* 1987; 26: 197-202.
- Madhok R, Zoma A, Torley H I, Capell H A, Waring R, Hunter J A. The relationship of sulphoxidation status to efficacy and toxicity of penicillamine in the treatment of rheumatoid arthritis. *Arthritis Rheum* 1990; 33: 574-7.

Numbers (percentages) of patients receiving sulphasalazine or D-penicillamine who were good or poor sulphoxidisers

	Poor	Good
Sulphasalazine (n=63)	49 (78)	14 (22)
D-penicillamine (n=53)	36 (68)	17 (32)
Expected	26 (22)	90 (78)

Toxicity due to sulphasalazine or D-penicillamine in the good and poor sulphoxidisers groups.