samples were tested by both solid phase assays. None of the 25 serum samples tested was positive for the anti-29 kilodalton ELISA. Four out of 27 serum samples were positive for the anti-myeloperoxidase ELISA and negative for antinuclear antibody testing. Anti-myeloperoxidase positive serum samples were P-ANCA positive (n=3) or immunofluorescence negative (n=1). No marked hypergammaglobulinaemia could explain 'false' positive results by non-specific Fc receptor binding. No association was found between the presence of ANCA and either vasculitis or glomerulonephritis. The serum of one patient with relapsing polychondritis-Wegener's granulomatosis overlap contained P-ANCA at a titre of 1/20. Among the eight patients positive for ANCA, relapsing polychondritis was active in seven, whereas it was active in only 12 out of 25 ANCA negative patients (p=0.1, two tailed Fisher's test).

These results suggest that low titres of C-ANCA are not specific for Wegener's granulomatosis and that ANCA (either diffuse or perinuclear) may be present in 24% of serum samples from patients with relapsing polychondritis, especially during the active phase of the disease.

T PAPO
J C PIETTE
LE THI HUONG DU
P GODEAU
Service de Médecine Interne
Groupe Hospitalier Pitié-Salpêtrière
75651 Paris Cedex 13, France

O MEYER
M F KAHN
Service de Rhumatologie
Hôpital Bichat
75018 Paris, France

P BOURGEOIS
Service de Rhumatologie
Groupe Hospitalier Pitié-Salpêtrière
75651 Paris Cedex 13, France

Correspondence to: Dr Thomas Papo, Service de Médecine Interne, Groupe Hospitalier Pitié-Salpêtrière, 83 bd de l'Hôpital, 75651 Paris Cedex 13, France.

- 1 Ludeman J, Utecht B, Gross W. Anti-neutrophil cytoplasm antibodies in Wegener's granulomatosis recognize an elastolytic enzyme. *J Exp Med* 1990; 171: 357-62.
- 2 Hagen E C, Ballieux B E, Dahan M R, van Hes L A, van der Woude F J. Fundamental and clinical aspects of antineutrophil cytoplasmic antibodies. *Autoimmunity* 1992; 11: 199-207.
- 3 Specks U, Wheatley C L, McDonald T J, Rohrbach M S, Deremee R A. Anticytoplasmic autoantibodies in the diagnosis and follow-up of Wegener's granulomatosis. *Mayo Clin Proc* 1989; 64: 28-36.
- 4 Michet C J, McKenna C H, Luthra H S, O'Fallon W M. Relapsing polychondritis. *Ann Intern Med* 1986; 104: 74-8.
- 5 Wiik A. Delineation of a standard procedure for indirect immunofluorescence detection of ANCA. *Acta Pathol Microbiol Immunol Scand* 1989; 97: 12-13.

Two sisters with ANCA positive vasculitis

Antineutrophil cytoplasmic antibodies (ANCA) have been shown to be associated strongly with microscopic polyarteritis nodosa and Wegener's granulomatosis.^{1,2} These antibodies have also been found in patients with other vasculitic disorders, including Kawasaki disease, Churg-Strauss syndrome, and relapsing polychondritis.^{3,4} Factors governing production of ANCA

remain unknown, though infection has been implicated as a possible trigger for onset of relapse of Wegener's granulomatosis.^{5,6} We describe two sisters with ANCA positive vasculitic diseases.

The first patient, aged 55 years, presented with arthralgia and subsequently developed uveitis and scleritis, swelling of the nasal cartilage, chondritis of her ear, and costochondritis. She had an episode of transient diplopia and a Bell's palsy. A diagnosis of relapsing polychondritis was made. She was initially treated with steroids and a non-steroidal anti-inflammatory drug but subsequently required azathioprine to control her symptoms. She remains well six years later, having had two minor relapses in the interim. There has been no evidence of renal disease.

The second sister presented at the age of 54 years with nasal stuffiness, impaired hearing, episcleritis, myalgia, abnormal liver function, and proteinuria. A diagnosis of Wegener's granulomatosis was made. She has responded well to cyclophosphamide and remains in remission 18 months later.

Investigations have shown that both sisters have been repeatedly positive for ANCA. Patient 1 had a perinuclear pattern of staining (pANCA) with a maximum titre of 1/64 (on treatment), whereas patient 2 had a cytoplasmic pattern of staining (cANCA) with a maximum titre of 1/512. The sisters are HLA identical—A2, A10, B27, Bw4, Cw1, DR8, DR9.

As far as we know, this is the first report of ANCA positive vasculitis in siblings. There is a striking similarity between the age of onset and the clinical features of the two cases despite the different diagnoses. They are HLA identical. These cases suggest a role for genetic factors in the development of ANCA positive vasculitis, which merits further study. They also highlight the fact that within the spectrum of vasculitis there is considerable overlap in the clinical features of patients with different vasculitic disorders.

It is now recognised that the immunofluorescent staining patterns obtained when testing for ANCA reflect different antibody specificities—cANCA being associated with antibodies to proteinase-3 and pANCA being associated with antibodies to myeloperoxidase in some cases. It is therefore of interest that our two cases had different staining patterns.

There have been two recent reports of an HLA association in patients with ANCA positive vasculitis. Spencer *et al* found an association between HLA-DQw7 and susceptibility to ANCA related disease, while possession of HLA-DR5 or DR6 seemed to prolong the duration of ANCA synthesis.⁷ Papiha *et al* reported an association between Wegener's granulomatosis and HLA-DR1.⁸ It is therefore of interest that these two sisters are HLA identical, though they do not possess any of the HLA antigens previously associated with ANCA positive vasculitis.

E A MURPHY
R D STURROCK
Centre for Rheumatic Diseases
University Department of Medicine
Glasgow Royal Infirmary
Glasgow G3 7 2ER
United Kingdom

J G FOX
J M BOULTON-JONES
Renal Unit
University Department of Medicine
Glasgow Royal Infirmary
Glasgow G3 7 2ER
United Kingdom

- 1 Van der Woude F J, Rasmussen N, Lobatto S, *et al*. Autoantibodies against neutrophils and monocytes: tool for diagnosis and marker of disease activity in Wegener's granulomatosis. *Lancet* 1985; i: 425-9.
- 2 Lockwood C M, Bakes D, Jones S, Whitaker K B, Moss D W, Savage C O S. Association of alkaline phosphatase with an autoantigen recognised by circulating anti-neutrophil antibodies in systemic vasculitis. *Lancet* 1987; i: 716-9.
- 3 Venning M C, Arfeen S, Bird A G. Antibodies to neutrophil cytoplasmic antigen in systemic vasculitis. *Lancet* 1987; ii: 850.
- 4 Savage C O S, Lockwood C M. Autoantibodies in primary systemic vasculitis. *Ann Intern Med* 1990; 35: 73-92.
- 5 Pinching A J, Rees A J, Pussell B A, *et al*. Relapses in Wegener's granulomatosis: the role of infection. *BMJ* 1980; 281: 836.
- 6 Csernok E, Schmitt W H, Ernst M, Bainton D, Gross W L. Translocation of proteinase-3 on the cell membrane of polymorphonuclear leucocytes in Wegener's granulomatosis: correlation with disease activity. *Clin Rheumatol* 1992; 11: 124.
- 7 Spencer S J, Burns A, Gaskin G, Pusey C, Rees A J. Influence of HLA class II genes on susceptibility to vasculitis. *Nephrol Dial Transplantation* 1991; 6: 900.
- 8 Papiha S S, Murty G E, Ad'Hia A, Mains B T, Venning M. Association of Wegener's granulomatosis with HLA antigens and other genetic markers. *Ann Rheum Dis* 1992; 51: 246-8.

Acquired Brown's syndrome in a patient with SLE

Sir: We read with interest the recent article by Alonso-Valdivielso *et al*¹ describing a patient with systemic lupus erythematosus (SLE) who later developed Brown's syndrome. In view of the authors' statement that this association has not previously been described we wish to draw your attention to two previously published reports.

In 1990 we published a case report² in which the patient, a 30 year old man, initially sought medical advice as a result of his ocular symptoms of variable double vision on upgaze. Only the left eye was affected and on examination restricted elevation of the adducted left eye produced diplopia associated with pain and a palpable click over the trochlea. A diagnosis of left Brown's syndrome was made following Hess chart assessment. The condition settled during treatment with ibuprofen 1200 mg daily. A five month history of arthralgia affecting the hands, wrists, elbows, shoulders, hips, and knees and acute alopecia one year before was noted. Later examination during febrile flare disclosed rash, lymphadenopathy, and synovitis of the hands, left elbow, shoulders, and knees. Investigations, including antinuclear antibody, dsDNA antibody, complement and immune complex levels, confirmed SLE.

In our discussion we made reference to an earlier review of Brown's syndrome which included a description of the condition occurring in a patient with established SLE.³ We agree with Alonso-Valdivielso *et al* that Brown's syndrome should be considered in patients with diplopia and SLE. We consider that the true incidence of clinical disease may well be higher than these three reports suggest, perhaps because of mild or transient symptomatology and difficulty in diagnosis without orthoptic assistance.

JAMES MCGALLIARD
Royal Liverpool University Hospital
Liverpool 7
United Kingdom

AUBREY L BELL
Rheumatology Unit
Musgrave Park Hospital
Stockmans Lane
Belfast BT9 7JB
and the School of Clinical Medicine
Queens University of Belfast
United Kingdom

Correspondence to: Dr Bell.

1 Alonso-Valdivielso J L, Alvarez Lario B, Alegre López J, Sedano Tous M J, Buitrago Gómez A. Acquired Brown's syndrome in a patient with systemic lupus erythematosus. *Ann Rheum Dis* 1993; 52: 63-4.
2 McGalliard J, Bell A L. Acquired Brown's syndrome in systemic lupus erythematosus: another ocular manifestation. *Clin Rheumatol* 1990; 9: 399-400.
3 Trimble R B, Kelly V, Mitchell M. Acquired Brown's syndrome. In: Ravault A P, Len M, eds. *Transactions of the fifth international orthoptic congress; Lyon*. 1983: 267-72.

**Femoral head
osteonecrosis in primary
hypertrophic
osteoarthropathy**

Primary hypertrophic osteoarthropathy is rare. One patient with primary hypertrophic osteoarthropathy has been described with bilateral hip osteoarthritis,¹ but as far as we are aware osteonecrosis of the femoral head has not previously been reported in this condition. Of 52 patients with primary hypertrophic osteoarthropathy seen in our department, three presented with symptoms relating to femoral head osteonecrosis. In all three cases osteonecrosis developed in both hips over a 3-12 month period, two out of three patients requiring replacements of both hips.

The table presents principal clinical and investigational features of these three patients. Abnormal liver function tests were present in patient 2 (raised transaminases, γ -glutamyltransferase, and bilirubin), which were thought to relate to excessive alcohol intake. Patients 1 and 3 had no other risk factor in their history that might have related to osteonecrosis. Interestingly, all three patients had peptic ulceration. In all three primary hypertrophic osteoarthropathy appeared in childhood or puberty, while osteonecrosis developed in middle age.

The femoral head is a common site of osteonecrosis² and as far as we are aware this is the first report of concurrence of

*Clinical, radiographic and scintigraphic
features, and laboratory findings of three male
patient with aseptic necrosis of the femoral head
(ANFH)*

	Patients		
	1	2	3
History			
Familial history	+	+	-
Age of onset PHO* (years)	11	5	12
Age of onset ANFH (years)	50	36	41
Skeletal pain	+	+	+
Symptoms			
Clubbing of hands	+	+	+
Clubbing of feet	+	+	+
Seborrhoea	+	+	+
Hyperhidrosis	+	+	+
Thickening of distal legs	+	+	+
Periarticular swelling of ankles	+	-	+
Radiography			
Periostitis of long bones	+	+	+
Demarcation of anterolateral part of femoral heads	+	+	+
Scintigraphy			
High uptake of radiotracer in both hips	+	+	+
High uptake of radiotracer in long bones	+	+	+

*PHO=primary hypertrophic osteoarthropathy.

osteonecrosis with primary hypertrophic osteoarthropathy. Although alcohol excess may be incriminated in one of the three cases, in the other two no obvious predisposing factors were apparent. It is generally postulated that the stimulus to new bone formation in primary hypertrophic osteoarthropathy relates to increased local blood flow, possibly related to an endocrine mechanism³ or neurogenic reflex arc.⁴ Alteration in bone perfusion associated with primary hypertrophic osteoarthropathy might thus in some way have predisposed to osteonecrosis in these patients. The three patients described were otherwise typical of primary hypertrophic osteoarthropathy.⁵ Occurrence of peptic ulceration in primary hypertrophic osteoarthropathy has previously been described,⁶ though the mechanism remains unclear.

IVO JAJIC
Lovencka 100
41000 Zagreb
Croatia

1 Pineda C, Fosceca C, Martinez-Lavin M. The spectrum of soft tissue and skeletal abnormalities of hypertrophic osteoarthropathy. *J Rheumatol* 1990; 17: 626-32.
2 Davidson J K. *Aseptic necrosis of bone*. Amsterdam-Oxford: Excerpta Medica, and New York: American Elsevier Publishing Company, 1976.
3 Ginsburg J, Brown J B. Increased oestrogen excretion in hypertrophic pulmonary osteoarthropathy. *Lancet* 1961; ii: 1274-6.
4 Holling H E, Brodey R S, Boland H C. Pulmonary hypertrophic osteoarthropathy. *Lancet* 1961; ii: 1269-74.
5 Jajić I, Pećina M, Krstulović B, et al. Primary hypertrophic osteoarthropathy and changes in the joint; clinical, X-ray, scintigraphic, arteriographic and histologic examination of 19 patients. *Scand J Rheumatol* 1980; 9: 89-96.
6 Skiu Kum Lam, William K K, Kui K, et al. Pachydermoperiostosis, hypertrophic gastropathy, and peptic ulcer. *Gastroenterology* 1983; 84: 834-9.