Annals of the

RHEUMATIC DISEASES

Foreword

Investigators pondering the cause of inflammatory rheumatic diseases have been largely preoccupied with immunology since the discovery of autoantibodies over 30 years ago. The ability to initiate organ specific autoimmune diseases by the experimental presentation of organ specific autoantigens to the immune system gave a further incentive to pursue this theme. An analysis of the major rheumatological journals shows that on average some 10–15% of all articles published over the past 20 years have a major immunological component, and this takes no account of immunological investigations into the rheumatic diseases appearing in other scientific journals. Much of this massive body of data has informed us about immunopathological mechanisms. For example, it is highly probable that cytokines synthesised in the rheumatoid synovial membrane make a major contribution to cartilage degradation. Similarly, it is likely, though less securely established, that HLA associations with rheumatic diseases operate through T cell recognition of genetically determined, topographical variations of oligopeptides presented in the antigen-presenting grooves. However, not unreasonably, rheumatologists and their patients ask relatively simple questions about cause, effect, and interdiction, and it is in this context that the contribution of immunology to rheumatological understanding has also to be assessed.

There are two fundamentally different approaches to interpreting immunological aberrations in the rheumatic diseases. The first argues that these aberrations are primary events. Persistent inflammation in the rheumatoid synovial membrane and autoantibody production in systemic disorders like systemic lupus erythematosus result from a breakdown in tolerance. A rigorous analysis of events such as peptide processing in antigen-presenting cells, the genes encoding T cell antigen receptors, and the genes determining variable sequences in autoantibody synthesis can eventually be expected to disclose abnormalities predisposing to immunoproliferative disorders and immune mediated tissue damage. The strongest argument favouring this view is the contribution of genetic factors to human rheumatic diseases and animal models of rheumatic diseases.

From an alternative viewpoint, the immunological abnormalities in the rheumatic diseases are mainly epiphenomena which may on occasion inflict damage on tissues or circulating cells but do not initiate these diseases. The wealth of microbial disorders accompanied by similar abnormalities is the most cogent reason for adopting this stance. Rational research in a disease such as rheumatoid arthritis should be directed at seeking the ghost in the immunopathological machine. Many microbes are adept at acting

as agent provocateurs and it is vital that their evolutionary patience is matched by those who bestow research grants.

These seemingly contrasting interpretations are not mutually exclusive. 'Some infectious diseases are dangerous to us not because the body fails to defend itself against them but—paradoxically—because it does defend itself: in a sense, the remedy is the disease'.¹ As a cogent example, immunodeficiency predisposes to rheumatic disorders, through atypical infections in common variable hypogammaglobulinaemia and putatively from infections in complement deficiency associated with connective tissue diseases. Severe forms of immunodeficiency, such as defects in HLA molecule assembly, may well prove to be gross examples of commoner defects predisposing to persistent infections with immunological abnormalities. In these subjects immune mediated disease may be the price of host defence. Furthermore, postinfectious reactive disease, including arthritis, is one of the few situations in which there is some intellectual security about the validity of immunological reasoning.

This Heberden issue is a selection of contributions from scientists who have approached these problems in very different ways. The diversity of these studies illustrates the justifiably broad spectrum of immunological studies in the rheumatic diseases. Indeed there are few areas of scientific enterprise in which dogmatism about research strategy is so misplaced; a moment of youthful insight may have the same investigative potential as the efforts of a respected team working on a well defined project.

Avrion Mitchison examines the problem from the standpoint of tolerance and its breakdown. As a scientist who played a major part in the classical debate on the nature of tolerance, his interest in this field has been sustained in the current era when transgenic technology has allowed earlier theories to be subjected to rigorous experimental scrutiny. The study of immunity to collagen and its control is a plausible model for probing the link between immunological reasoning and an autoimmune aetiology for rheumatoid arthritis.

Complementing Avrion Mitchison's paper, Hans Acha-Orbea shows how microbial infections may trigger polyclonal T and B cell reactivity by acting as "superantigens", rather than conventional antigens. This process could abrogate tolerance and thereby initiate autoimmune diseases. The technical elegance of the experiments he describes holds the promises of reviving decades of speculation about the link between common infectious agents and the rheumatic diseases. His paper also illustrates the complex manner in which retroviruses subvert the control of immune response. However, ironically, the extent to which the abrogation of classical

tolerance is relevant to the cause of a disease such as rheumatoid arthritis remains debatable; indeed this is still a disease in search of a universally accepted autoantigen.

The T cell infiltration characteristic of rheumatoid synovitis has been a fertile source of material for immunological experiments; these cells serve as the nearest approach to an intelligent, dolphin-like probe, able to relay aetiological information to the investigator.² Many theories about the immunopathogenesis of the disease presuppose that there are dominant clones, inducing disease because they react with relevant autoantigens or microbial antigens. Hill Gaston examines the evidence for this belief and rightly stresses the practical difficulties in identifying, isolating, and characterising such clones, even granted that they do exist.

Many immunorheumatologists have been preoccupied with assessing the pathological significance of rheumatoid factors and some observers have considered the generation of these self reactive antibodies to be the major autoimmune mechanism responsible for the disease. The discovery of seemingly defective IgG glycosylation in rheumatoid arthritis has prompted speculation that this structural anomaly may make some immunoglobulin sequences abnormally antigenic. Protein glycosylation is involved in many forms of protein interaction, immune regulation, and control of endothelial cells in the microvasculature. The paper by Elizabeth Hounsell and Michael Davies examines these complexities and underlines the danger of oversimplification in this field.

The passing years and scientific sophistication have not dispelled the suspicion that rheumatoid arthritis, and even multisystem connective tissue diseases, are really forms of reactive arthritis initiated and maintained by bacterial infection. The dilemma is obvious; many decades of intensive effort have not revealed agents with the credentials to satisfy an increasingly cynical rheumatological community yet theories based on 'hit and run' mechanisms are more likely to convince theoretical physicists than hard nosed microbiologists. The paper by Graham Rook, Peter Lydyard, and J L Stanford reminds us that bacteria, and particularly mycobacteria, are also accomplished survivors in the human host and have their

own strategy for confusing immune defences. This paper draws the important conclusion that the issues can be resolved by the techniques now available and are unlikely to run unresolved into the next century.

Retroviruses cause neoplastic, immunodeficiency, and immune mediated degenerative diseases in many animal species. With the exception of the last category, analogous diseases have been discovered in man. Faith in a retroviral cause for rheumatoid arthritis, systemic lupus erythematosus, and other connective tissue diseases is at least as rational as any other belief based on theory rather than incontrovertible evidence. There is the additional twist that the distinction between many exogenous retroviral sequences and some endogenous retroviral sequences in the human genome is almost semantic. Steffen Gay, Renate Gay, and William Koopman review the mechanisms by which retroviral infection could account for the immunopathological features of rheumatoid arthritis and the evidence supporting this hypothesis.

Finally, Shozo Izui, Thierry Berney, Takanori Shibata, and Thierry Fulpius show how immunochemical definition can account for the distinctive pattern of disease in a murine model of systemic lupus erythematosus and, by inference, in the human disease. Their contribution emphasises the potential of analytical immunochemistry for explaining disease phenomena. Moreover, their studies show how genetically determined selection of the antibody repertoire could interact with environmental factors such as infections to produce many late manifestations of human connective tissue diseases.

*Clinical Research Centre,
Division of Immunological Medicine,
Watford Road,
Harrow,
Middlesex HA1 3UJ,
United Kingdom*

A M DENMAN

1 Medawar P B. *The threat and the glory*. Oxford: Oxford University Press, 1990: 175-6.

2 Denman A M. Lymphocytes: dolphins for enlightening rheumatological investigators? *J Rheumatol* 1986; 13: 9-12.