Bacterial and viral superantigens: roles in autoimmunity?

Hans Acha-Orbea

Abstract

Superantigens are bacterial, viral, or retroviral proteins which can activate specifically a large proportion of T cells. In contrast with classical peptide antigen recognition, superantigens do not require processing to small peptides but act as complete or partially processed proteins. They can bind to major histocompatibility complex class II molecules and stimulate T cells expressing particular T cell receptor V_β chains. The other polymorphic parts of the T cell receptor, which are crucial for classical antigen recognition, are not important for this interaction. When this strategy is used a large proportion of the host immune system can be activated shortly after infection. The activated cells have a wide variety of antigen specificities. The ability to stimulate polyclonal B (IgG) as well as T cell responses raises possibilities of a role for superantigens in the induction of autoimmune diseases. Superantigens have been a great tool in the hands of immunologists in unravelling some of the basic mechanisms of tolerance and immunity.

(*Ann Rheum Dis* 1993; 52: S6-S16)

Classical antigen recognition by T cells occurs by interaction of self major histocompatibility complex (MHC) proteins, which bind small fragments of proteins, with T cell receptor (TCR). These three components are briefly described below. More detailed information can be found in excellent reviews and the references therein.¹⁻⁶

The MHC encodes many different proteins. Among these the MHC class I and class II molecules are important in this context. These dimeric molecules are highly polymorphic in the population. Of the order of 50 alleles have been defined for the different MHC class I and class II molecules. The major function of these molecules is the presentation of antigens to the T cells by acting as receptors able to bind thousands of 8-9 (MHC class I) or 9-16 (MHC class II) amino acid long processed peptides. Such peptides are generated from proteins in the cytoplasm (for MHC class I presentation) or in endosomes (for MHC class II presentation) by specialised proteases. Most of the antigens entering the cell from the outside will be associated with MHC class II molecules, whereas proteins produced within

the cell are presented by MHC class I molecules. Despite the high numbers of peptides which can be bound to a particular MHC molecule, these receptors show striking specificity for peptides they bind. This specificity is due to differences in the amino acid sequence in the N-terminal part of MHC molecules which forms the peptide binding pocket that can accommodate two to four amino acids of the peptide. Specific amino acids could be implicated at specific localisations in the bound peptides for several different MHC class I as well as class II molecules. Different MHC class I and class II proteins bind different sets of peptides. Single amino acid differences in this peptide binding region can heavily influence the peptides which can bind to a particular MHC class I or class II molecule.

The formation of a trimolecular complex between MHC, peptide, and TCR is highly specific in that a particular TCR can interact only with specific MHC molecules which have bound a particular peptide. For this interaction TCR and MHC sequences as well as peptide sequences are important. As recognition occurs almost exclusively with self MHC molecules, and the same peptides cannot be presented in the context of another MHC molecule, it is called MHC restricted. This MHC restricted antigen recognition by T lymphocytes leads to formation of a trimolecular complex between short antigenic peptides bound to MHC molecules and recognition by a TCR (fig 1A).

In addition to these molecules, a series of other receptors and coreceptors intensify this interaction. Most of these intensifying molecules do not contribute to the specificity of the interaction and are not discussed here. One pair of these coreceptors intensifies the interaction of the TCR with MHC class I molecules (CD8) or MHC class II molecules (CD4). Cytotoxic T cells are found amongst the former, and helper T cells amongst the latter.

The TCR is the end product of a series of recombination events during the maturation and differentiation of T cells. This brings together the constant region which is shared between all the receptor molecules with the two or three highly polymorphic parts: variable (V), diversity (D) (in the case of the β chain only), and junctional (J) elements. These elements are encoded as V and J clusters, separated but in the same chromosomal localisation as the constant region. In addition, a tremendous heterogeneity is introduced by

Figure 1 Classical peptide and superantigen recognition. (A) In classical antigen recognition a processed peptide is presented by major histocompatibility complex (MHC) class I or class II molecules to the T cell receptor (TCR). The CD4 or CD8 molecule intensifies this interaction. (B) In superantigen recognition the superantigen crosslinks the TCR V β with MHC class II. It is not known whether antigenic peptides are bound to MHC and whether CD4 and CD8 are of importance.

random introduction and deletion of short sequences in the V-D, V-J, and D-J regions during this recombination process. The functional TCR is composed of an α and a β chain (which are encoded on different chromosomes and use different elements), and, as only one type of TCR molecule is expressed on one T cell clone, is responsible for giving the T cell the high specific recognition capability. In mice there are about 25 V β , 12 J β , 75 V α , and 75 J α elements encoded. With the introduced random sequences of the order of 10^{15} different TCR molecules can theoretically be produced, by far exceeding the potential repertoire of a single mouse. In humans about 100 V β elements exist, whereas the other TCR elements are similar in number in mice and humans. In humans the V β families, which are defined by 70% amino acid homology, are much larger than in the mouse. The number of families is comparable. In a classical MHC restricted peptide specific T cell response about one in 100 000 T cells can interact with one particular immunogenic peptide-self MHC complex with significant affinity.

Biology of superantigens

The term superantigen has been given to antigens which can activate of the order of one in 10 T cells, allowing the activation of about 10 000 times more T lymphocytes with a single antigen than in classical MHC-peptide recognition.⁷⁻¹² In contrast with T cell mitogens, superantigens have a high degree of specificity.

It became clear recently that superantigens have a new way of interaction with MHC molecules and the TCR. Instead of interacting with the most polymorphic part of the receptor, they recognise one or several different TCR V β regions which make only a small contribution to the overall heterogeneity.¹³⁻¹⁸ They can interact with MHC molecules and cross link the TCR to form an aberrant trimolecular complex (fig 1B). Neither the TCR nor the trimolecular complex has yet been crystallised, but structural homology with previously

crystallised structures allows us to suggest a tentative structure.¹⁻⁴ These superantigens are not processed to small peptides like the classical antigens but function as entire molecules¹⁹⁻²¹ (or possibly partially cleaved molecules in the case of retroviral superantigens, see below). Based on the analysis of MHC class II and TCR mutants which lost or gained superantigen binding, as well as the recent crystallisation and mutagenesis analysis on the bacterial superantigen staphylococcal enterotoxin B, it is now thought that superantigens cross link the TCR with MHC class II molecules on the lateral side of the complex (fig 1B).^{9-11 21-30}

Recently, with the help of superantigens and transgenic mice, direct evidence for two basic tolerance mechanisms (clonal deletion of self reactive cells in the thymus^{13 14 31-36} and induction of unresponsiveness (anergy) in the periphery^{37 38}) was produced. In this review only the results obtained with superantigens are discussed.

T cells mature in the thymus, where they learn to distinguish between self and foreign. Early in life the immune system is plastic and can adapt to learn what belongs to self and what is foreign. The thymus plays an important part in this decision. Cells which are strongly self reactive are negatively selected (deleted), whereas T cells which can interact weakly with self MHC molecules are positively selected for export into the periphery.^{33 39-41} Owing to positive selection the cells are trained to recognise antigens bound to self MHC molecules, as otherwise a large percentage of T cells which cannot interact with self MHC molecules would be wasted. Negative selection in the thymus (deletion) of 'self reactive' T cells has been found in mice expressing superantigens at birth.^{7 12-16 42} With the help of monoclonal antibodies specific for V β regions of the TCR an almost complete elimination of the superantigen reactive V β -bearing T cells has been noted in mature thymocytes as well as in the peripheral T cell repertoire.

A first encounter with a superantigen in an adult mouse in vivo leads to an initial expansion of the reactive T cells bearing

one or several specific TCR V_{β} regions.^{7 8 12-14 23 43 44} Shortly after this encounter the T cells become unresponsive (anergic).^{37 38 45-47} In the case of anergy the cells are still present but have lost the ability to respond a second time to the same antigen; this is in striking contrast with a second challenge with a conventional antigen where a secondary response is more intense. One of the important findings is that these anergic T cells lose the ability to produce one of the important lymphokines for T cell survival and division—namely, interleukin 2.^{37 38 45 48 49} In some but not all cases this state of anergy can be reversed by addition of exogenous interleukin 2. With the availability of monoclonal antibodies specific for TCR V_{β} and V_{α} regions it became easy to measure these changes in the frequencies of the responding T cells. As they make up about 10% of the total mouse T cell repertoire they can easily be followed. Negative selection in the thymus can eliminate 90–99% of T cells expressing a TCR V_{β} element that can interact with a particular superantigen.

Negative selection in the thymus can also be induced with an exogenous antigen, such as a bacterial superantigen injected at birth.⁷ Normally, however, negative selection in the thymus (as well as tolerance) is only maintained if the antigen is present continuously and will fade out with the disappearance of the antigen. For longlasting tolerance the antigen has to be present continuously, as in the case of retroviral infection (see below).

Nature of superantigens

Superantigens have been found in different infectious agents. The recently described superantigens originate from bacteria (staphylococci,^{7 11 44 50-52} streptococci,^{53 145-147} mycoplasma,^{54 148 149} yersinia⁵⁵), retroviruses (mouse mammary tumour virus (MMTV),^{9 56-66} possibly, human immunodeficiency virus (HIV),^{67 68} possibly, murine leukaemia virus⁶⁹), and other viruses (rabies virus⁷⁰). The exclusive distribution of these molecules in microbes raises the question of how important the expression of these structures is for microbes. Probably, the strategy of activation of the host immune response leads to an overstimulation of the immune system which allows the pathogens more efficient infection.^{71 72} Up to now, however, only few results show such an advantage directly.

Bacterial superantigens

Overall, the described bacterial superantigens are of similar size (20–30 kilodaltons) with high to very low amino acid sequence homology (20–80%) (for review see ref 11). Most likely many of them developed by convergent evolution towards the same aim. These bacterial superantigens have been shown to interact strongly with MHC class II but not MHC class I molecules.^{19-21 73 74} Their affinities are of the same order of magnitude as

the binding affinities of antibodies to their antigen (10^{-6} – 10^{-8} M). They can bind to mouse, rat, and human MHC class II antigens. Moreover, for T cell stimulation to occur, class II expressing cells are required. Table 1 provides a summary of the differences between superantigen and classical peptide recognition. Normally, T cells expressing the surface marker CD4 (amongst which helper T cells are found) react with antigens presented by MHC class II molecules, whereas cells expressing CD8 (which include cytotoxic T cells) react with antigens presented by MHC class I molecules. Bacterial superantigens, however, break this rule; both T cell subpopulations are easily stimulated by MHC class II expressing cells in the presence of bacterial superantigens.⁷⁴ In addition, no MHC restriction is seen. For example, mouse superantigen presenting cells can stimulate human T cells. The only requirement is MHC class II expression of the presenting cell and expression of a specific TCR V_{β} chain by the T cell.

Different MHC class II isotypes have different affinities for various bacterial superantigens.^{19-21 73} In general, they interact better with human than mouse MHC class II proteins. Superantigens are the most potent T cell mitogens known. Their affinity constants are in the picomolar to nanomolar range. Table 2 lists the currently known bacterial superantigens and their TCR V_{β} specificities in mice and humans. In mice many more TCR V_{β} specific monoclonal antibodies are available, which makes it much easier to define these specificities. Therefore the TCR specificities of bacterial superantigens in mice are more completely documented than in humans.

One of these superantigens, staphylococcal enterotoxin B (SEB), has recently been crystallised.³⁰ In addition, introduction of mutations in the SEB coding sequence allows localisation of the MHC and the TCR interaction residues.²⁹ For all the analysed superantigens the affinity for the TCR seems to be much lower than for MHC molecules. In addition to contact residues on the superantigen, amino acid contact residues on the TCR and the MHC class II molecules have been characterised for both retroviral (see below) and bacterial superantigens.²¹⁻²⁹ The overall conclusions of these studies suggest that unprocessed superantigen molecules can interact with the lateral side of the TCR and the MHC class II molecules (see fig 1B).

As described above, injection of bacterial superantigens into newborn mice leads to thymic clonal deletion of the reactive T cells.⁷ Injection into adult mice leads to an expansion of the reactive T cells, which thereafter leads to induction of non-responsiveness (anergy).^{48 49} For a limited time period the T cells expressing the responsive TCR V_{β} are reduced in number in vivo. Anergy can also be induced in T cell clones with bacterial superantigens.⁷⁵

A recent report showed that injection of

Table 1 Characteristics of superantigens

	Classical peptide antigen	Superantigen	
Origin	Protein	Bacterial	Viral
Processing	Yes (small peptides)	No	Partial or no
Presentation	MHC* class I, or class II MHC restricted	MHC class II, not MHC restricted	MHC class II on B cells, not MHC restricted
Frequency of responding cells	1 in 10 ⁴ to 10 ⁶	1 in 5 to 1 in 20	1 in 5 to 1 in 20
TCR* elements	V _α J _α , V _β J _β	V _β	V _β

*MHC=major histocompatibility complex; TCR=T cell receptor.

Table 2 Bacterial superantigens

TCR*V _β	In humans	In mice	Reference
1	—	SEA*	134,135, Acha-Orbea, unpublished
2	TSST-1*, streptococcal M protein	—	52,145
3	SEB	SEA, SED, TSST-1, <i>Yersinia enterocolitica</i> extract, streptococcal pyrogenic exotoxin A	7,51,52,55,134,135,146, Acha-Orbea, unpublished
4	Streptococcal M protein	—	145
5	SEC3, SED, SEE	—	51,52
6	SEE	MAM, * <i>Yersinia enterocolitica</i> extract	51,52,55,130,149
7	—	SEB, SEC3, SED	7,135,136
8-1	SEE, streptococcal M protein	SEB, SEC3, SED, MAM	7,51,52,130,135,136,145
8-2	SEE, streptococcal M protein	SEB, SEC(1,2,3), SED, MAM, streptococcal pyrogenic exotoxin A	7,51,52,130,135,136,145,146
8-3	SEE, streptococcal M protein	SEB, SEC1, SED, MAM	7,51,52,130,135,136,145
10	—	SEA, SEC2	23,135
11	—	SEA, SEC1, SED, SEE, <i>Yersinia enterocolitica</i> extract, streptococcal pyrogenic exotoxin A	55,134,135,146, Acha-Orbea, unpublished
12	SEB, SEC(1,2,3) SED	SEA	51,52,134,135, Acha-Orbea, unpublished
13	SEC2	—	51,52
14	SEB, SEC2	—	51,52
15	SEB, SEC2	SEE, TSST-1	51,52,135
17	SEB, SEC2	SEA, SEC(1,2), SED, TSST-1	51,52,135
18	SEE	—	51,52
20	SEB, SEC2	—	51,52

Note that the most homologous TCR V_β elements do not have the same numbers in mice and humans.¹³⁷ The molecular structures of *Yersinia enterocolitica*, *Mycoplasma arthritidis* superantigens have not yet been characterised.
*TCR=T cell receptor; TSST-1=toxic shock syndrome toxin; SEA, SEB, SEC, SED, SEE=staphylococcal enterotoxin A, B, C, D, E; MAM=*Mycoplasma arthritidis* supernatant.

superantigens can lead to a profound inhibition of conventional antigen responses.⁷¹ It is still not clear, however, whether it is this effect that forced bacteria to develop superantigens. Well known effects of superantigens include food poisoning after ingestion of staphylococcal exotoxins,⁷⁶⁻⁷⁷ the toxic shock syndrome,⁷⁸⁻⁸⁰ arthritis after mycoplasma infections,⁸¹ possibly Kawasaki disease,⁸² and possibly rheumatoid fever after streptococcal infections^{83-85 148 149} (see below). Table 3 lists the diseases and the causing superantigens.

Retroviral superantigens in mice

In mice, genes have been mapped which encode endogenous superantigens.^{9 56-63 66 86 87} They were originally described as minor lymphocyte stimulating (Mls) antigens by Festenstein some 20 years ago.⁴³ He discovered that a large proportion of cells responsive to Mls antigens are stimulated in an MHC non-

restricted fashion when T cells from mice lacking a particular Mls antigen, such as Mls-1^a, are mixed with B cells expressing the Mls-1^a antigen. It was clear that the genes encoding these proteins are dominant and, surprisingly, always have an Mls allele and a null allele which are unable to stimulate such a vigorous T cell response.^{8 9 88-91} Many such biallelic systems have been defined in laboratory mouse strains and map different chromosomal localisations. It was discovered later that the T cells reacting with Mls antigens share expression of particular TCR V_β chains.^{9 13-18 42 65 92-99} Mice expressing such an Mls gene delete most of the T cells expressing the responsive TCR V_β chain during thymic maturation. After an intensive search for over 20 years it became clear recently that these Mls antigens are encoded by endogenous proviruses of the MMTV family.^{9 56-64 66 86 87} Figure 2 shows the typical buildup of such a virus. In the 3' long terminal repeat, an open reading frame (ORF) was found which represented a puzzle to retrovirologists.^{9 100-102} They had difficulties ascribing a function to this potential protein, which was never found in normal or tumour cells. It was thought that this protein had an effect on gene regulation and both positive and negative gene regulation were described.^{103 104} It turned out that it is this protein which represents a superantigen.^{9 59 60} As viral genes often have more than one function it will be interesting to see if other functions can be

Table 3 Bacterial superantigens and disease

Superantigen	Disease
Staphylococcal enterotoxins	Food poisoning, shock
Staphylococcal TSST-1*	Toxic shock syndrome
<i>Mycoplasma arthritidis</i> supernatant (MAS)	Arthritis, shock
Streptococcal superantigens: M protein, pyrogenic exotoxins	Rheumatic fever, shock, scarlet fever

*TSST-1=toxic shock syndrome toxin.

Figure 2 Mouse mammary tumour virus. The superantigen encoded in the open reading frame (ORF) molecule has not yet been identified biochemically. Possibly, therefore, the potential cleavage sites in the figure are cleaved before the superantigen functions. Clearly, however, the superantigen is expressed on the B cell surface.^{152 153} LTR=long terminal repeat.

found for this ORF. Most laboratory as well as wild mice have two to 10 copies of different MMTV proviruses integrated into their genome.¹⁰⁵ The different MMTVs are about 95% homologous in their amino acid sequence. Table 4 lists the currently known endogenous MMTVs and their TCR V_{β} specificities.

When ORF molecules interacting with the different TCR V_{β} elements are aligned a striking correlation between the C-terminal amino acid sequence and the TCR V_{β} specificity is observed.^{9 59 60 86} At present some 15 ORF sequences are known and in all cases a striking correlation between C-terminal sequence and TCR V_{β} specificity exists. The last 10–30 amino acids seem to determine the V_{β} specificity. Figure 2 shows the presumed structure of an ORF molecule.

These endogenous superantigens also lead to thymic deletion of the reactive cells when encountered at birth. Thymic deletion can also be induced after injection of cells bearing a superantigen into adult mice.⁸⁶ As described above for bacterial superantigens, immune stimulation of the reactive T cells occurs when the first encounter happens in adult life

followed by induction of unresponsiveness (anergy). Whether the loss of the reactive T cells after this initial stimulation is due to peripheral deletion or to dilution of the cells which can no longer divide is not clear at the moment.

These endogenous superantigens originate from infectious viruses which are still found in wild and laboratory mice (for reviews on MMTV see refs 102, 106–110). In contrast with endogenous transmission, which follows the principles of mendelian inheritance, infectious viruses are transmitted through milk from mother to offspring. The following scenario for infection has been suggested by experiments in newborn or adult mice: the virus is taken up through milk from birth and contacts lymphocytes in the Peyer's patches (Karapetian O, Shakhov A N, Kraehenbuhl J P, *et al*, unpublished data). As soon as the immune system matures sufficiently (around two to three days after birth), infection of B cells is followed by superantigen (ORF) mediated T-B interaction with T cells expressing the responsive TCR V_{β} elements.^{86 111} A strong augmentation of the infected B cells which express the superantigen occurs¹¹¹; at the same time a continuous stimulation (which would be harmful for the host) is prevented by anergisation of the responsive T cells. The superantigen expression allows about 1000 times more efficient infection (Held W, Waanders G A, Shakhov A N *et al*, unpublished data). We have recently shown that the virus infects lymphocytes exclusively in Peyer's patches and the lymph nodes of the small intestine (Karapetian O, Shakhov A N, Kraehenbuhl J P, *et al*, unpublished data). The infection of lymphocytes represents a crucial step in the life cycle of the virus. Mice lacking responsive T cells or mice with a defective immune system cannot propagate the virus efficiently.¹¹² Later in life the virus jumps from lymphocytes to the mammary gland epithelium where large scale virus production occurs which is secreted into the milk. This is the only place where free virus particles have been found. When the mothers transmit the virus to their offspring the vicious circle is closed. Figure 3 shows a typical life cycle of MMTV.

As the name implies, these viruses are the major causative agent of mammary tumours in mice. Integration close to a mouse proto-oncogene can activate it and induce the first steps towards carcinogenesis.^{113 114}

This virus has very cleverly profited from the weaknesses of the mouse's immune system. In doing so it has given us tools which can help us to understand better its normal function. The finding of an 'infectious superantigen' (MMTV), especially, allows experiments which help towards understanding the immune system, the retroviral life cycle, and the interplay between the two.^{64 86 111 115} These experiments are still in progress. One important point for this discussion is the effect on B cells after local infection of adult mice with MMTV. Adult mice cannot eliminate MMTV; they remain infected for the rest of

Table 4 Retroviral superantigens

Mouse	Human	Retroviral superantigen	References
2	?	Mtv-MA, MMTV(C4)	Wei and Acha-Orbea, and Marche <i>et al</i> , in preparation
3	?	Mtv-1, -3, -6, -13, -44, -MAI	58,59,63,138–140
5	?	Mtv-6, MuLV?	57,69
5-1	?	Mtv-8, -9	56–58
5-2	?	Mtv-8, -9	56–58
6	12	Mtv-7, -43, -44, -SW, MMTV(SW)	Shakhov and Acha-Orbea, in preparation
7	?	Mtv-7, -29, -43, -SW, -29, MMTV(SW)	59,63,87,141, Shakhov and Acha-Orbea, in preparation, Marrack, personal communication
8-1	?	Mtv-7, -43, -SW, MMTV(SW)	59,63,87,141, Shakhov and Acha-Orbea, in preparation
9	?	Mtv-7, -43, -SW, MMTV(SW)	59,63,87,141, Shakhov and Acha-Orbea, in preparation
11	?	Mtv-8, -9, -11	62
12	?	Mtv-8, -9, -11	97–99
14	?	Mtv-2, MMTV(GR), MMTV(C3H)	27,59,144
15	?	Mtv-2, MMTV(GR), MMTV(C3H)	27
16	?	Mtv?	97–99,143
17	?	{ Mtv-3, -8, -9 Mtv-6, -MAI	140, 154
19	?	New Mtv?	Hodes, personal communication
20	?	Mtv-?	65

Note that the most homologous TCR V β elements do not have the same numbers in mice and humans. Mtv-n stands for integrated proviruses, MMTV stands for infectious viruses. Most human TCR V β -MMTV ORF interactions have not yet been measured. TCR=T cell receptor; MMTV=mouse mammary tumour virus; ORF=open reading frame.

Figure 3 Life cycle of mouse mammary tumour virus. TCR=T cell receptor.

their life and transmit the virus maternally to their offspring. The first cells which seem to be infected with MMTV are the B cells. Expression of the superantigen leads to strong activation of helper T cells (but not cytotoxic cells in vivo) expressing the reactive TCR V β element. At the same time this T-B interaction leads to terminal activation of the B cells. These B cells start secreting large amounts of IgG antibody and increase in number.¹¹¹ Such an induction of polyclonal IgG production might be part of autoimmune reactions (see below). Similar observations have been made with bacterial superantigens.^{54 71 116 117} Figure 4 provides a schematic summary of these observations.

In the case of MMTV superantigen the answer to the question of why these viruses have adopted such a strategy seems answered. The superantigen allows about 1000 times more efficient infection of the host lymphocytes, which seems crucial for the survival of the virus.

There are still gaps in our knowledge about the interplay between virus and host, but so far (and surely even more so in the near future) retroviral and bacterial superantigens have been great tools in the hands of immunologists, allowing consideration of questions about the normal immune response as well as retrovirology.

Table 5 compares current knowledge on bacterial and retroviral superantigens.

Human viral superantigens

During the last year two reports suggested that HIV may encode a superantigen.^{67 68} The first analysed the expression of mRNA of the different TCR V β elements in the peripheral blood of patients at late stages of AIDS. Several TCR V β elements were found to be reduced strongly in patients with late stage disease, after the drop in the CD4+ T cell population. A semiquantitative polymerase chain reaction was used to measure mRNA expression in unseparated blood isolates.⁶⁷ Analyses on purified CD4+ T cell populations are required to consider this question more directly as, possibly, the V β s which were reduced are underrepresented in the CD8+ T cell population.

In another study it was shown that fresh T cell lines expressing TCR V β 6-7, 8, and 17 or TCR V β 12 could be infected similarly with HIV but, after addition of antigen presenting cells, TCR V β 12 expressing cells had an approximately 100-fold higher HIV titre than the other three, suggesting a superantigen effect in HIV amplification.⁶⁸ More needs to be done to show convincingly that HIV encodes a superantigen. If it does, it opens new strategies for vaccination and would make the MMTV system a model of choice for new vaccination studies directed at the superantigen effects.

In another study it was clearly shown that the rabies virus encodes a superantigen in its nuclear protein.⁷⁰ Recombinant protein showed strong binding to MHC class II

Figure 4 Fate of T and B cells after mouse mammary tumour virus superantigen interaction *in vivo*. TCR=T cell receptor.

Table 5 Comparison of retroviral and bacterial superantigens

	Bacterial superantigens	Retroviral superantigens
Stimulation	Yes	Yes or no
Deletion	Yes	Yes
Anergy	Yes	Yes
Cytotoxicity	Yes	No
APC†: B*	Yes	Yes
APC: class II expressing transfectants	Yes	No
APC: macrophages	Yes	No
Sequence homology to host proteins	None	None
Sequence homology between members‡	20–90%	95%
Binding to MHC‡ class II	Strong	? (Most likely)
Binding to TCR‡	Weak	? (Most likely)
Processing required for presentation	No	No or partial cleavage

*B cells as antigen presenting cells
†Amino acid sequence homology between different bacterial superantigens or between different retroviral superantigens.
‡APC=antigen presenting cell; MHC=major histocompatibility complex; TCR=T cell receptor.

molecules and stimulated T cells expressing V β 12 in humans and V β 6 (the closest murine homologue) in mice. It will be interesting to see the effect of this superantigen on the efficiency of infection with rabies virus. Once this virus reaches the nerve cells it is no longer accessible to the immune system. In addition, it will be interesting to see if and how this superantigen plays a part in the vaccines used to protect against rabies infection. The results of these experiments will potentially give insights into the role of superantigens in vaccination.

Roles of superantigens in autoimmunity?

It is well known that normal subjects have some autoreactive T and B cells in their circulation. Polyclonal activation of B cells induces production of autoantibodies, and autoreactive CD4+ T cells can easily be isolated after stimulation *in vitro* with autoantigens.^{118–125} The overall tolerance mechanisms can maintain a healthy equilibrium between the presence of autoreactive cells and the prevention of their predominance. Several points in the above

discussion make it likely that superantigens may play a part in the induction of autoimmune diseases: (a) superantigens can induce polyclonal antibody responses; (b) superantigens can activate T cells with many different fine specificities, possibly including autoreactive T cells.

Mimicry by microbes has been proposed previously for a trigger of autoimmune reactions.¹²⁶ In this hypothesis a sequence with similarity to host sequences is expressed by the microbe. A strong immune reaction to the invading microorganism leads to the breaking of tolerance to an autoantigen. Superantigens do not even have to express such a cross reactive epitope. By activation of a substantial proportion of the host immune cells expressing millions of different antigen specificities, several autoreactive cells might be activated (fig 5). It is not clear at present what happens to these cells when they encounter an auto-antigen after superantigen activation. In experimental allergic encephalomyelitis, a model disease for human multiple sclerosis, few TCR molecules are found on the autoreactive T cells. Most share expression of TCR V β 8·2.^{150 151} The bacterial superantigen SEB reactivates all the V β 8 bearing T cells. Preliminary experiments indicate that SEB induces relapses of experimental allergic encephalomyelitis in mice, clearly suggesting the possibility of a role in classical antigen responses by potentially anergised cells (Steinman L A, personal communication). Most likely it depends on the cell type which presents the autoantigen after the superantigen encounter.

Another elegant experiment gives support for both the original mimicry as well as the superantigen hypothesis in the induction of autoimmunity.^{127 128} Mice were generated which bear two types of transgene. One

Figure 5 Potential role in autoimmunity of superantigens. APC=antigen presenting cell; TCR=T cell receptor.

transgene determines the expression of an epitope for cytotoxic T cells exclusively on the pancreatic β cells, the targets of autoimmune destruction in insulin dependent diabetes mellitus. The antigen is the glycoprotein of the leucocytic choriomeningitis virus. The other transgene controls the expression of the TCR α and β chains isolated from a cytotoxic T cell specific for a peptide of this glycoprotein. Mice expressing both transgenes are non-diabetic and remain normal. Infection with a control leucocytic choriomeningitis virus which does not bear the T cell epitope does not induce diabetes, whereas infection with the leucocytic choriomeningitis virus expressing the T cell epitope induces diabetes within a few days.

Several recent studies have begun to consider questions about the role of superantigens in autoimmunity. In rheumatoid arthritis, again using semiquantitative PCR, an overrepresentation in the rheumatoid lesions of a particular TCR V_{β} chain has been found, with reduced representation in the periphery.¹²⁹ These findings were made using semiquantitative PCR and have to be confirmed by more quantitative methods. As soon as the necessary monoclonal antibodies are available, such analyses can be repeated on a large number of patients. In another autoimmune disease, Kawasaki disease, a high percentage of T cells expressing $V_{\beta}2$ has been found in the peripheral blood of patients.⁸² Kawasaki disease is an acute multisystem vasculitis affecting young children. It is often found in Japanese patients. *Mycoplasma arthritidis* antigens, which have not been biochemically characterised, might yet be involved in the induction of polyarthritis in mice.^{81 130 148 149} Other candidates are yersinia superantigens and streptococcal antigens.^{55 83-85 145-147} Reiter's disease is a reactive arthritis, a term used for arthritis developing after infection. Candidate infectious agents causing reactive arthritis in humans are yersinia, shigella, salmonella, and others.¹³¹⁻¹³³ An involvement of superantigens

or classical mimicry in autoimmunity would be compatible as well with geographical variations in the prevalence of autoimmune diseases.

It is still too early clearly to correlate specific autoimmune diseases with specific bacterial or viral superantigens. Indeed, it is not yet clear whether superantigens are important for autoimmunity or not. So far all the correlations amount to little more than speculation. Even if microbes can be clearly linked to the induction of autoimmune diseases, mechanisms other than superantigens are likely to contribute. The near future will tell what their role is in the known autoimmune diseases. Whatever the outcome of these studies, pathogens compensate for the morbidity they inflict by providing us with highly valuable tools for dissecting the immune system and perhaps will give us insight into the cause of some forms of autoimmunity and its prevention.

HAO was supported by a START fellowship of the Swiss National Science Foundation.

- 1 Bjorkman P J, Parham P. Structure, function, and diversity of class I major histocompatibility complex molecules. *Annu Rev Biochem* 1990; 59: 253-88.
- 2 Nepom G T, Erlich H. MHC class II molecules and autoimmunity. *Annu Rev Immunol* 1991; 9: 493-526.
- 3 Rothbard J B, Geffer M L. Interactions between immunogenic peptides and MHC. *Annu Rev Immunol* 1991; 9: 527-66.
- 4 Davis M M, Bjorkman P J. T-cell antigen receptor genes and T-cell recognition. *Nature* 1988; 334: 395-402.
- 5 Jorgensen J L, Reay P A, Ehrlich E W, Davis M M. Molecular components of T-cell recognition. *Annu Rev Immunol* 1992; 10: 835-73.
- 6 Janeway C A Jr. The T cell receptor as a multicomponent signaling machine: CD4/CD8 coreceptors and CD45 in T cell activation. *Annu Rev Immunol* 1992; 10: 645-74.
- 7 White J, Herman A, Pullen A M, et al. The V_{β} -specific superantigen staphylococcal enterotoxin B: stimulation of mature T cells and clonal deletion in neonatal mice. *Cell* 1989; 56: 27-35.
- 8 Abe R, Hodes R. T cell recognition of minor lymphocyte stimulating (Mls) gene products. *Annu Rev Immunol* 1989; 7: 683-708.
- 9 Acha-Orbea H, Palmer E. Mls—a retrovirus exploits the immune system. *Immunol Today* 1991; 12: 356-61.
- 10 Janeway C A Jr. Self-superantigens? *Cell* 1990; 63: 659-61.
- 11 Marrack P, Kappler J. The staphylococcal enterotoxins and their relatives. *Science* 1990; 248: 705-11.
- 12 MacDonald H R, Glasebrook A L, Schneider R, et al. T-cell reactivity and tolerance to Mls^s encoded antigens. *Immunol Rev* 1989; 107: 89-108.

- 13 MacDonald H R, Schneider R, Lees R L, *et al.* T-cell receptor V β use predicts reactivity and tolerance to Mls^I-encoded antigens. *Nature* 1988; **332**: 40–5.
- 14 Kappler J W, Staerz U D, White J, Marrack P C. Self-tolerance eliminates T cells specific for Mls-modified products of the major histocompatibility complex. *Nature* 1988; **332**: 35–40.
- 15 Kappler J W, Roehm N, Marrack P. T cell tolerance by clonal elimination in the thymus. *Cell* 1987; **49**: 273–80.
- 16 Kappler J W, Wade T, White J, *et al.* A T cell receptor V β segment that imparts reactivity to a class II major histocompatibility complex product. *Cell* 1987; **49**: 263–71.
- 17 Abe R, Vacchio M S, Fox B, Hodes R. Preferential expression of the T-cell receptor V β 3 gene by Mls^c reactive T cells. *Nature* 1988; **335**: 827–30.
- 18 Fry A M, Matis L A. Self-tolerance alters T-cell receptor expression in an antigen-specific MHC-restricted immune response. *Nature* 1988; **335**: 830–2.
- 19 Mollick J A, Cook R G, Rich R R. Class II MHC molecules are specific receptors for staphylococcal enterotoxin A. *Science* 1989; **244**: 817–20.
- 20 Fischer H, Dohlstein M, Lindwall M, Sjögren H-O, Carlsson R. Binding of staphylococcal enterotoxin A to HLA-DR on B cell lines. *J Immunol* 1989; **142**: 3151–7.
- 21 Fraser J D. High-affinity binding of staphylococcal enterotoxins A and B to HLA-DR. *Nature* 1989; **339**: 221–3.
- 22 Dellabonna P, Peccaud J, Kappler J, *et al.* Superantigens interact with MHC class II molecules outside of the antigen groove. *Cell* 1990; **62**: 1115–21.
- 23 Herman A, Kappler J, Marrack P, Pullen A M. Superantigens: mechanism of T-cell stimulation and role in immune responses. *Annu Rev Immunol* 1991; **9**: 745–72.
- 24 Pullen A M, Wade T, Marrack P, Kappler J. Identification of the region of the T cell receptor β chain that interacts with the self-superantigen Mls-1^a. *Cell* 1990; **61**: 1365–74.
- 25 Pullen A M, Bill J, Kubo R, Marrack P, Kappler J W. Analysis of the interaction site for the self superantigen Mls-1^a on T cell receptor V β . *J Exp Med* 1991; **173**: 1183–92.
- 26 Cazenave P A, Marche P, Jouvin-Marche E, *et al.* V β 17 gene polymorphism in wild-derived mouse strains: two amino acid substitutions in the V β 17 region alter drastically T cell receptor specificity. *Cell* 1990; **63**: 717–28.
- 27 Choi Y, Herman A, DiGusto D, *et al.* Regions of the variable region of the T-cell receptor β -chain that interact with *S. aureus* toxin superantigens. *Nature* 1990; **346**: 471–3.
- 28 Karp D R, Long E O. Identification of HLA-DR1 β chain residues critical for binding staphylococcal enterotoxins A and E. *J Exp Med* 1992; **175**: 415–24.
- 29 Kappler J, Herman A, Clements J, Marrack P. Mutations defining functional regions of the superantigen enterotoxin B. *J Exp Med* 1992; **175**: 387–96.
- 30 Swaminathan S, Furey W, Pletcher J, Sax M. Crystal structure of staphylococcal enterotoxin B, a superantigen. *Nature* 1992; **359**: 801–6.
- 31 Kisielow P, Blüthmann H, Staerz U D, Steinmetz M, von Boehmer H. Tolerance in T-cell-receptor transgenic mice involves deletion of nonmature CD4⁺8⁺ thymocytes. *Nature* 1988; **333**: 742–6.
- 32 von Boehmer H, Kisielow P. Self-nonsel self discrimination by T cells. *Science* 1990; **248**: 1369–72.
- 33 Sha W C, Nelson C A, Newberry R D, *et al.* Positive and negative selection of an antigen receptor on T cells in transgenic mice. *Nature* 1988; **336**: 73–6.
- 34 Uematsu Y, Ryser S, Dembic Z, *et al.* In transgenic mice the introduced functional T cell receptor β chain gene prevents expression of endogenous β genes. *Cell* 1988; **52**: 831–41.
- 35 Pircher H P, Lang R, Hengartner H, Zinkernagel R M. Tolerance induction in double specific T cell receptor transgenic mice varies with antigen. *Nature* 1989; **342**: 559–61.
- 36 Berg L J, de St Groth B F, Pullen A M, Davis M M. Phenotypic differences between alpha beta versus beta T-cell receptor transgenic mice undergoing negative selection. *Nature* 1989; **340**: 559–62.
- 37 Webb S, Morris C, Sprent J. Extrathymic tolerance of mature T cells: clonal elimination as a consequence of immunity. *Cell* 1990; **63**: 1249–56.
- 38 Rammensee H-G, Kroschewsky R, Frangoulis B. Clonal anergy induced in mature V β 6 T lymphocytes on immunizing Mls-1^b mice with Mls-1^a expressing cells. *Nature* 1989; **339**: 541–4.
- 39 Kisielow P, Teh H S, Blüthmann H, von Boehmer H. Positive selection of antigen-specific T cells in thymus by restricting MHC molecules. *Nature* 1988; **335**: 730–3.
- 40 MacDonald H R, Lees R K, Schneider R, Zinkernagel R M, Hengartner H. Positive selection of CD4⁺ thymocytes controlled by MHC class II gene products. *Nature* 1988; **336**: 471–3.
- 41 Benoist C, Mathis D. Positive selection of the T cell repertoire: where does it occur? *Cell* 1989; **58**: 1027–33.
- 42 MacDonald H R, Pedrazzini T, Schneider R, *et al.* Intrathymic elimination of Mls^c-reactive (V β 6⁺) cells during neonatal tolerance induction to Mls-1^a encoded antigens. *J Exp Med* 1988; **167**: 2005–10.
- 43 Festenstein H. Immunogenic and biological aspects of in vitro allotransformation (MLR) in the mouse. *Transplant Rev* 1973; **15**: 62–88.
- 44 Janeway C A Jr, Yagi J, Conrad P J, *et al.* T-cell responses to Mls and to bacterial proteins that mimic its behavior. *Immunol Rev* 1989; **107**: 61–88.
- 45 Jones L A, Chin L T, Longo D L, Kruisbeek A M. Peripheral clonal elimination of functional T cells. *Science* 1990; **250**: 1726–9.
- 46 Fowlkes B J, Schwartz P H, Pardoll D M. Deletion of self-reactive thymocytes occurs at a CD4⁺8⁺ precursor stage. *Nature* 1988; **334**: 620–3.
- 47 Blackman M R, Gerhardt-Burgert H, Woodland D L, *et al.* A role for clonal inactivation in T cell tolerance to Mls-1^a. *Nature* 1990; **345**: 540–2.
- 48 Kawabe Y, Ochi A. Programmed cell death and extrathymic reduction of V β 8⁺ CD4⁺ T cells in mice tolerant to Staphylococcus aureus enterotoxin B. *Nature* 1991; **349**: 245–8.
- 49 MacDonald H R, Baschieri S, Lees R K. Clonal expansion precedes anergy and death of V β 8⁺ peripheral T cells responding to staphylococcal enterotoxin B in vivo. *Eur J Immunol* 1991; **21**: 1963–6.
- 50 Fleischer B, Schrezenmeier H. T cell stimulation by staphylococcal enterotoxins. Clonally variable response and requirements for major histocompatibility complex class II molecules on accessory or target cells. *J Exp Med* 1988; **167**: 1697–707.
- 51 Kappler J, Kotzin B, Herron L, *et al.* V β -specific stimulation of human T cells by staphylococcal toxins. *Science* 1989; **244**: 811–13.
- 52 Choi Y, Kotzin B, Herron L, *et al.* Interaction of Staphylococcus aureus toxin superantigens with human T cells. *Proc Natl Acad Sci USA* 1989; **86**: 8941–5.
- 53 Tomai M, Kolb M, Majumdar G, Beachey E H. Superantigenicity of streptococcal M protein. *J Exp Med* 1990; **172**: 359–62.
- 54 Tumang J R, Posnett D N, Cole B C, Crow M K, Friedman S M. Helper T cell-dependent human B cell differentiation mediated by a mycoplasma superantigen bridge. *J Exp Med* 1990; **171**: 2153–8.
- 55 Stuart P M, Woodward J G. Yersinia enterocolitica produces superantigen activity. *J Immunol* 1992; **148**: 225–33.
- 56 Woodland D, Happ M P, Bill J, Palmer E. Requirement for cotolerogenic gene products in the clonal deletion of I-E reactive T cells. *Science* 1990; **247**: 964–7.
- 57 Woodland D L, Happ M P, Gollub K J, Palmer E. An endogenous retrovirus mediating deletion of $\alpha\beta$ T cells? *Nature* 1991; **349**: 529–30.
- 58 Woodland D L, Lund F E, Happ M P, *et al.* Endogenous superantigen expression is controlled by mouse mammary tumor proviral loci. *J Exp Med* 1991; **174**: 1255–8.
- 59 Acha-Orbea H, Shakhov A N, Scarpellino L, *et al.* Clonal deletion of V β 14 positive T cells in mammary tumor virus transgenic mice. *Nature* 1991; **350**: 207–11.
- 60 Choi Y, Kappler J W, Marrack P. A superantigen encoded in the open reading frame of the 3' long terminal repeat of mouse mammary tumor virus. *Nature* 1991; **350**: 203–7.
- 61 Choi Y, Marrack P, Kappler J. Structural analysis of a mouse mammary tumor virus superantigen. *J Exp Med* 1992; **175**: 847–52.
- 62 Dyson P J, Knight A M, Fairchild S, Simpson E, Tomonari K. Genes encoding ligands for deletion of V β 11 T cells cosegregate with mammary tumor virus genomes. *Nature* 1991; **349**: 531–2.
- 63 Frankel W N, Rudy C, Coffin J M, Huber B T. Linkage of Mls genes to endogenous mammary tumor viruses of inbred mice. *Nature* 1991; **349**: 526–8.
- 64 Marrack P, Kushnir E, Kappler J. A maternally inherited superantigen encoded by a mammary tumor virus. *Nature* 1991; **349**: 524–6.
- 65 Six A, Jouvin-Marche E, Loh D Y, Cazanave P A, Marche P N. Identification of a T cell receptor β chain variable region, V β 20, that is differentially expressed in various strains of mice. *J Exp Med* 1991; **174**: 1263–6.
- 66 Pullen A M, Choi Y, Kushnir E, Kappler J, Marrack P. The open reading frames in the 3' long terminal repeats of several mouse mammary tumor virus integrants encode V β 3-specific superantigens. *J Exp Med* 1992; **175**: 41–7.
- 67 Imberti L, Sottini A, Bettinardi A, Puoti M, Primi D. Selective depletion in HIV infection of T cells that bear specific T cell receptor V β sequences. *Science* 1992; **254**: 860–2.
- 68 Laurence J, Hodsre A S, Posnett D N. Superantigen implicated in dependence of HIV-1 replication in T cells on TCR V β expression. *Nature* 1992; **358**: 255–9.
- 69 Hügin A W, Vacchio M S, Morse H C III. *Science* 1991; **252**: 424–7.
- 70 Lafon M, Lafage M, Martinez-Arends A, *et al.* Evidence for a human superantigen in humans. *Nature* 1992; **358**: 507–10.
- 71 Mourad W, Scholl P, Diaz A, Geha R, Chatila T. The staphylococcal toxic shock syndrome toxin 1 triggers B cell proliferation and differentiation via major histocompatibility complex-unrestricted cognate T/B cell interaction. *J Exp Med* 1989; **170**: 2011–22.
- 72 Mithke T, Wahl C, Heeg K, Echtenacher B, Krammer P H, Wagner H. T cell-mediated lethal shock triggered in mice by the superantigen staphylococcal enterotoxin B: critical role of tumor necrosis factor. *J Exp Med* 1992; **175**: 91–8.
- 73 Herrmann T, Accolla R S, MacDonald H R. Different staphylococcal enterotoxins bind preferentially to distinct major histocompatibility complex class II isotypes. *Eur J Immunol* 1989; **19**: 2171–4.
- 74 Herrmann T, Mariansky J L, Romero P, Fleischer B, MacDonald H R. Activation of MHC class I-restricted CD8⁺ CTL by microbial T cell mitogens. Dependence

- upon MHC class II expression of the target cells and V_β usage of the responder cells. *J Immunol* 1990; **144**: 1181–6.
- 75 O'Hehir R E, Lamb J R. Induction of specific clonal anergy in human T lymphocytes by *Staphylococcus aureus* enterotoxins. *Proc Natl Acad Sci USA* 1990; **87**: 8884–8.
 - 76 Bergdoll M S. Enterotoxins. In: Moutie T C, Kadis J, Aji S J, eds. *Microbial toxins. III. Bacterial protein toxins*. New York: Academic Press, 1970: 265–326.
 - 77 Bergdoll M S. Staphylococcal intoxications. In: Riemann H, Bryan F L, eds. *Food borne infections and intoxications*. New York: Academic Press, 1979: 443–94.
 - 78 Todd J K, Fishaut M, Kapral F, Welch T. Toxic shock syndrome associated with phage-group I staphylococci. *Lancet* 1978; **i**: 1116–8.
 - 79 Bergdoll M S, Crass B A, Reiser R F, Robbins R N, Davis J P. A new staphylococcal enterotoxin F, associated with toxic shock syndrome isolates. *Lancet* 1981; **i**: 1017–21.
 - 80 Schlievert P M, Shands K N, Dan B B, Schmid G P, Nishimura R D. Identification and characterization of an exotoxin from *Staphylococcus aureus* associated with toxic shock syndrome. *J Infect Dis* 1981; **143**: 509–16.
 - 81 Cole B C, Washburn L R, Taylor-Robinson D. Mycoplasma-induced arthritis. In: Razin S, Barile M F, eds. *The mycoplasmas* Vol IV. New York: Academic Press, 1985: 105–60.
 - 82 Abe J, Kotzin B L, Jujo K, et al. Selective expansion of T cells expressing T-cell receptor variable regions V_β2 and V_β8 in Kawasaki disease. *Proc Natl Acad Sci USA* 1992; **89**: 4066–70.
 - 83 Bartter T, Dascal A, Carroll K, Curley F J. Toxic streptococcal syndrome. *Arch Intern Med* 1988; **148**: 1421–4.
 - 84 Quinn DeJoy S, Ferguson K M, Sapp T M, et al. Streptococcal cell wall arthritis. Passive transfer with a T cell line and crossreactivity of streptococcal cell wall antigens with *Mycobacterium tuberculosis*. *J Exp Med* 1989; **170**: 369–82.
 - 85 Bisno A L. Group A streptococcal infections and acute rheumatic fever. *N Engl J Med* 1991; **325**: 783–93.
 - 86 Held W, Shakhov A N, Waanders G A, et al. An exogenous mouse mammary tumor virus with properties of Mls-1^a (Mtv-7). *J Exp Med* 1992; **175**: 1623–33.
 - 87 Beutner U, Frankel W N, Cote M S, Coffin J M, Huber B T. Mls-1 is encoded by the long terminal repeat open reading frame of the mouse mammary tumor virus Mtv-7. *Proc Natl Acad Sci USA* 1992; **89**: 5432–6.
 - 88 Festenstein H, Berumen L. BALB.D2-Mls^a new congenic mouse strain. *Transplantation* 1983; **37**: 322–4.
 - 89 Abe R, Ryan J J, Hodes R J. Mls is not a single gene, allelic system. Different stimulatory Mls determinants are the products of at least two nonallelic, unlinked genes. *J Exp Med* 1987; **166**: 1150–5.
 - 90 Abe R, Ryan J J, Hodes R J. Clonal analysis of the Mls system. A reappraisal of polymorphism and allelism among Mls^a, Mls^b, and Mls^d. *J Exp Med* 1987; **165**: 1113–29.
 - 91 Pullen A M, Marrack P, Kappler J W. Evidence that Mls-2 antigens which delete V_β3⁺ T cells are controlled by multiple genes. *J Immunol* 1989; **142**: 3033–7.
 - 92 Pullen A M, Marrack P, Kappler J W. The T-cell repertoire is heavily influenced by tolerance to polymorphic self-antigens. *Nature* 1988; **335**: 796–801.
 - 93 Bill J, Appel V, Palmer E. An analysis of T cell receptor variable region gene expression in major histocompatibility complex disparate mice. *Proc Natl Acad Sci USA* 1988; **85**: 9184–8.
 - 94 Bill J, Kanagawa O, Woodland D, Palmer E. The MHC molecule I-E is necessary but not sufficient for the clonal deletion of V_β11-bearing T cells. *J Exp Med* 1989; **169**: 1405–19.
 - 95 Happ M P, Woodland D C, Palmer E. A third T cell receptor V_β gene encodes reactivity to Mls-1^a gene products. *Proc Natl Acad Sci USA* 1989; **86**: 6293–6.
 - 96 Okada C Y, Holzmann B, Guidos S, Palmer E, Weissman I L. Characterization of a rat antibody specific for a determinant encoded by the V_β7 gene segment. *J Immunol* 1990; **144**: 3473–7.
 - 97 Vacchio M S, Hodes R. Selective decrease in T cell receptor V_β expression: decreased expression of specific V_β families is associated with expression of multiple MHC and non-MHC gene products. *J Exp Med* 1989; **170**: 1335–46.
 - 98 Vacchio M S, Ryan J J, Hodes R J. Characterization of the ligand(s) responsible for negative selection of V_β11- and V_β12-expressing T cells: effects of a new Mls-determinant. *J Exp Med* 1990; **172**: 807–13.
 - 99 Singer P A, Balderas R S, Theofilopoulos A N. Thymic selection defines multiple T cell receptor V_β "repertoire phenotypes" at the CD4/CD8 subtype level. *EMBO J* 1990; **9**: 3641–8.
 - 100 Donehower L A, Huang A L, Hagler G L. Regulatory and coding potential of the mouse mammary tumor virus long terminal repeat. *J Virol* 1981; **37**: 226–37.
 - 101 Fasel N, Pearson K, Buetti E, Diggelmann H. The region of mouse mammary tumor virus DNA containing the long terminal repeat includes a long coding sequence and signals for hormonally regulated transcription. *EMBO J* 1982; **1**: 3–7.
 - 102 Dickson C. Molecular aspects of mouse mammary tumor virus biology. *Int Rev Cytol* 1985; **108**: 119–47.
 - 103 Salmons B, Erfle V, Brem G, Günzburg W H. naf, a trans-regulating negative-acting factor encoded within the mouse mammary tumor virus open reading frame region. *J Virol* 1990; **64**: 6355–9.
 - 104 van Klaveren P, Bentvelzen P. Transactivating potential of the 3' open reading frame of murine mammary tumor virus. *J Virol* 1988; **62**: 4410–3.
 - 105 Kozak C, Peters G, Pauley R, et al. A standardized nomenclature for endogenous mouse mammary tumor viruses. *J Virol* 1987; **61**: 1651–4.
 - 106 Acha-Orbea H, Held W, Waanders G A, et al. Exogenous and endogenous mouse mammary tumor superantigens. *Immunol Rev*. In press.
 - 107 Nandi S, McGrath C M. Mammary neoplasia in mice. *Adv Cancer Res* 1973; **17**: 353–414.
 - 108 Hageman P C, Calafat J, Hilgers J. The biology of the mouse mammary tumor virus. In: Hilgers J, Sluysen M, eds. *Mammary tumors of the mouse*. Amsterdam, New York, Oxford: Elsevier, North Holland Biomedical Press, 1981: 391–463.
 - 109 Bentvelzen P, Hilgers J. Murine mammary tumor virus. In: Klein G, ed. *Viral oncology*. New York: Raven Press, 1980: 311–55.
 - 110 Dickson C, Peters G. Proteins encoded by mouse mammary tumor virus. *Curr Top Microbiol Immunol* 1983; **106**: 1–34.
 - 111 Held W, Shakhov A N, Izui S, et al. Superantigen-reactive CD4⁺ T cells are required to stimulate B cells after infection with mouse mammary tumor virus. *J Exp Med*. In press.
 - 112 Tsubura A, Inaba M, Imai S, et al. Intervention of T-cells in transportation of mouse mammary tumor virus (milk factor) to mammary gland cells in vivo. *Cancer Res* 1988; **48**: 6555–8.
 - 113 Nusse R, Varmus H E. Many tumors induced by the mouse mammary tumor virus contain a provirus integrated in the same region of the host genome. *Cell* 1982; **31**: 99–109.
 - 114 Peters G, Brookes S, Smith R, Dickson C. Tumorigenesis by mouse mammary tumor virus: evidence for a common region for provirus integration in mammary tumors. *Cell* 1983; **33**: 369–77.
 - 115 Ignatowicz L, Kappler J, Marrack P. The effects of chronic infection with a superantigen-producing virus. *J Exp Med* 1992; **175**: 917–23.
 - 116 He X, Goronzy J, Weyand C. Selective induction of rheumatoid factors by superantigens and human helper T cells. *J Clin Invest* 1992; **89**: 673–80.
 - 117 Tumang J R, Cherniack P, Gietl D M, et al. T helper cell dependent, microbial superantigen-induced murine B cell activation: microclonal and antigen-specific antibody responses. *J Immunol* 1991; **147**: 432.
 - 118 Hohlfield R, Toyka K V, Heininger K, et al. Autoimmune human T lymphocytes specific for acetylcholine receptor. *Nature* 1984; **310**: 244–6.
 - 119 Martin R, Jaraquemada D, Flerlage M, et al. Fine specificity and HLA restriction of myelin basic protein-specific cytotoxic T cell lines from multiple sclerosis patients and healthy individuals. *J Immunol* 1990; **145**: 540–8.
 - 120 Pette M, Fujita K, Wilkinson D, et al. Myelin autoreactivity in multiple sclerosis: recognition of myelin basic protein in the context of HLA-DR2 products by T lymphocytes of multiple-sclerosis patients and healthy donors. *Proc Natl Acad Sci USA* 1990; **87**: 7968–72.
 - 121 Wucherpfennig K W, Ota K, Endo N, et al. Shared human T cell receptor V_β usage to immunodominant regions of myelin basic protein. *Science* 1990; **248**: 1016–9.
 - 122 Sun J-B, Olsson T, Wang W-Z, et al. Autoreactive T and B cells responding to myelin proteolipid protein in multiple sclerosis and controls. *Eur J Immunol* 1991; **21**: 1461–8.
 - 123 Ben-Nun A, Liblau R S, Cohen L, et al. Restricted T-cell receptor V_β usage by myelin basic protein-specific T-cell clones in multiple sclerosis: predominant genes vary in individuals. *Proc Natl Acad Sci USA* 1991; **88**: 2466–70.
 - 124 Liblau R, Tourmieu-Lasserre E, Maciazek J, et al. T cell response to myelin basic protein epitopes in multiple sclerosis patients and healthy subjects. *Eur J Immunol* 1991; **21**: 1391–5.
 - 125 Shirai A, Cosentino M, Leitman-Klinman S F, Klinman D M. Human immunodeficiency virus infection induces both polyclonal and virus-specific B cell activation. *J Clin Invest* 1992; **89**: 561–6.
 - 126 Oldstone M B A. Molecular mimicry and autoimmune disease. *Cell* 1987; **50**: 819–20.
 - 127 Ohashi P S, Oehen S, Bürki K, et al. Ablation of "tolerance" and induction of diabetes by virus infection in viral antigen transgenic mice. *Cell* 1991; **65**: 305–17.
 - 128 Oldstone M B A, Nerenberg M, Lewicki H. Virus infection triggers insulin-dependent diabetes mellitus in a transgenic model: role of anti-self (virus) immune responses. *Cell* 1991; **65**: 319–31.
 - 129 Paliard X, West S G, Lafferty J A, et al. Evidence for the effects of a superantigen in rheumatoid arthritis. *Science* 1991; **253**: 325–9.
 - 130 Cole B C, Karchner D R, Wells D J. Stimulation of mouse lymphocytes by a mitogen derived from mycoplasma arthritis (MAM). VIII. Selective activation of T cells expressing distinct V_β T cell receptors from various strains of mice by the superantigen MAM. *J Immunol* 1990; **144**: 425–31.
 - 131 Ahvonen P, Sievers K, Aho K. Arthritis associated with *Yersinia enterocolitica* infection. *Acta Rheum Scand* 1969; **15**: 232–53.
 - 132 Keat A. Reiter's syndrome and reactive arthritis in perspective. *N Engl J Med* 1983; **309**: 1606–15.

- 133 Inman R D. The interplay of microbe and MHC in the pathogenesis of Reiter's syndrome. *Clin Exp Rheumatol* 1986; **4**: 75-82.
- 134 Takimoto H, Yoshikai Y, Kishihara K, *et al*. Stimulation of all T cells bearing V β 1, V β 3, V β 11 and V β 12 by staphylococcal enterotoxin A. *Eur J Immunol* 1990; **20**: 617-21.
- 135 Callahan J E, Herman A, Kappler J, Marrack P. Stimulation of B10.Br T cells with superantigenic staphylococcal enterotoxins. *J Immunol* 1989; **144**: 2473-9.
- 136 Yagi J, Baron J, Buxser S, Janeway J A Jr. Bacterial proteins that mediate the association of a defined subset of T cell receptor: CD4 complexes with class II MHC. *J Immunol* 1990; **144**: 892-901.
- 137 Wilson R K, Lai E, Concannon P, Barth R K, Hood L E. Structure, organization and polymorphism of murine and human T-cell receptor α and β chain gene families. *Immunol Rev* 1988; **101**: 149-72.
- 138 Fairchild S, Knight A M, Dyson P J, Tomonari K. Co-segregation of a gene encoding a deletion ligand for Tcrb-V3* T cells with Mtv-3. *Immunogenetics* 1991; **34**: 227-30.
- 139 Fairchild S, Rosenwasser O A, Dyson P J, Tomonari K. Tcrb-V3* T-cell deletion and a new mouse mammary tumor provirus, Mtv-44. *Immunogenetics* 1992; **36**: 189-94.
- 140 Jouvin-Marche E, Cazenave P-A, Voegtli D, Marche P. V β 17 T-cell deletion by endogenous mammary tumor virus in wild-type-derived mouse strains. *Proc Natl Acad Sci USA* 1992; **89**: 3232-5.
- 141 Rudy C K, Kraus E, Palmer E, Huber B T. Mls-1-like superantigen in the MA/MyJ mouse is encoded by a new mammary tumor provirus. *J Exp Med* 1992; **175**: 1613-22.
- 142 Tomonari K, Fairchild S. Positive and negative selection of Tcrb-V6* T cells. *Immunogenetics* 1992; **36**: 230-7.
- 143 Vacchio M S, Kanagawa O, Tomonari K, Hodes R J. Influence of T cell receptor V α expression on Mls^s superantigen-specific T cells. *J Exp Med* 1992; **175**: 1405-8.
- 144 Ferrick D A, Cho K, Gemmell-Hori L, Morris D W. Genetic analysis of the effects of Mtv-2 on the T cell repertoire in the WXG-2 mouse strain. *International Immunology* 1992; **4**: 805-10.
- 145 Tomai M A, Aelion J A, Dockter M E, *et al*. T cell receptor V gene usage by human T cells stimulated with the superantigen streptococcal M protein. *J Exp Med* 1991; **174**: 285-8.
- 146 Imanishi K, Igarashi H, Uchiyama T. Activation of murine T cells by streptococcal pyrogenic exotoxin type A. *J Immunol* 1990; **145**: 3170-6.
- 147 Nelson K, Schlievert P M, Selander R K, Musser J M. Characterization and clonal distribution of speA gene encoding pyrogenic exotoxin A (scarlet fever toxin) in *Streptococcus pyogenes*. *J Exp Med* 1991; **174**: 1271-4.
- 148 Cole B C, Kartchner D R, Wells D J. Stimulation of mouse lymphocytes by a mitogen derived from *Mycoplasma arthritidis*. VII. Responsiveness is associated with expression of a product(s) of the V β 8 gene family present on the T cell receptor $\alpha\beta$ for antigen. *J Immunol* 1989; **142**: 4131-7.
- 149 Cole B C, Kartchner D R, Wells D J. Stimulation of mouse lymphocytes by a mitogen derived from *Mycoplasma arthritidis* (MAM). VIII. Selective activation of T cells expressing distinct V β T cell receptors from various strains of mice by the "superantigen" MAM. *J Immunol* 1990; **144**: 425-31.
- 150 Acha-Orbea H, Mitchell D J, Timmerman L, *et al*. Limited heterogeneity of T cell receptors in experimental allergic encephalomyelitis. *Cell* 1988; **54**: 263-73.
- 151 Acha-Orbea H, Steinman L, McDevitt H O. T cell receptors in murine autoimmune diseases. *Annu Rev Immunol* 1989; **7**: 371-405.
- 152 Acha-Orbea H, Scarpellino L, Shakhov A N, *et al*. Inhibition of mammary tumor virus-induced T cell responses in vivo by monoclonal antibodies to an open reading frame protein. *J Exp Med* 1992; **176**: 1769-72.
- 153 Winslow G M, Scherer M T, Kappler J W, Marrack P. Detection and characterization of the mouse mammary tumor virus 7 superantigen (Mls-1^s). *Cell* 1992; **71**: 719-30.
- 154 McDuffie M, Schweiger D, Reits B, *et al*. I-E-independent deletion of V β 17a* T cells by Mtv-3 from the nonobese diabetic mouse. *J Immunol* 1992; **148**: 2097-102.