MATTERS ARISING

Enthesitis of the ligamentum teres during ankylosing spondylitis: histopathological report

We were interested to read the article by Inoue *et al*¹ on ossification of the medial acetabular floor, including the ligamentum teres as a possible mechanism of lateral subluxation in coxopathy associated with ankylosing spinal hyperostosis. A few years ago we saw a patient suffering from ankylosing spondylitis (AS) with right coxitis including inflammatory enthesopathy of the ligamentum teres (ligamentum capitis femoris).

A 57 year old man with a five year history of a painful right hip was first seen at the rheumatology clinic in 1976, when the diagnosis of AS was made (HLA B27 was present). Anteroposterior radiography of the pelvis demonstrated joint space narrowing in the right hip, sclerosis and new bone formation on the right acetabular rim, enthesophytes of the femoral head, and an ankylosis in the sacro-iliac joints. Due to persistent pain and major disability, an arthroplasty was performed on the right hip.

On opening the capsule, the synovium was inflamed around the ligamentum capitis femoris although there was no obvious pannus the articular cartilage was considerably ulcerated at the femoral head. Histological examination showed an intense vascular proliferation in the central area of attachment of the femoral head ligament to the fovea capitis, especially in the deep subchondral perivascular area (figure). Lymphoplasmocytic cells had spread along the proliferating vessels which dissociate the

vertical collagen fibres at the ligamentum teres attachment. In the narrow spaces surrounding the damaged area there was evidence of chondrocyte proliferation. Ossifying enthesitis was also present on the attachment of the capsule in the acetabular labrum.

Published reports are scant concerning the histopathology in the early stage of AS. Nevertheless, several papers have underlined the role of enthesitis in the pathogenesis of AS and related spondylarthropathies.²⁻⁴ Ossifying enthesopathies on the great trochanter and capsular enthesiophytes are common features in AS⁵ and Forestier's disease⁶ with the resulting tendency to ankylosis of the hip.

Enthesopathies include all changes, whether traumatic, degenerative, metabolic or inflammatory of entheses.^{7,8} As enthesopathies usually produce ossification, they are very useful diagnostically and nosologically, especially when they are diffuse or multiple.⁵ The most extreme example of metabolic enthesopathy is diffuse idiopathic skeletal hyperostosis (DISH, more commonly known as Forestier's disease). Conversely, AS, which is the archetype of spondylarthropathies, is the more classic inflammatory enthesopathy.

The most distinctive feature of the natural history of ossifying enthesopathies, suggests that during AS and DISH, enthesopathy of the ligamentum capitis femoris, may initiate intra-articular involvement, as observed in the knee with cruciate ligaments.³ The presence of an intra-ligamentary artery probably enhances its proclivity to promote coxitis during AS.

P GILLET
P PÉRÉ
J Y JOUZEAU
J FLOQUET
A GAUCHER

Departments of Rheumatology and Pathology,
URA CNRS 1288, Faculté de Médecine, BP 184,
54505 Nancy,
France

1 Inoue K, Schichikawa K, Takenaka I, Yamauchi E, Nischioka N, Huda S. Ossification of the ligamentum teres as a possible mechanism of lateral subluxation in coxopathy associated

with ankylosing spinal hyperostosis. *Ann Rheum Dis* 1993; 52: 306-7.

2 Ball J. Enthesopathy of rheumatoid and ankylosing spondylitis. *Ann Rheum Dis* 1971; 30: 213-23.

3 Jacobs C J. Spondylarthritides and Enthesopathy. *Arch Int Med* 1983; 143: 103-7.

4 Schischikawa K, Tsujimoto M, Nischioka J, Nishibayashi Y, Matsumoto K. Histopathology of early sacroiliitis and enthesitis in ankylosing spondylitis. In: M Ziff, M B Cohen, eds. *Advances in inflammation research, vol 9: the spondylarthropathies*. New York: Raven Press, 1985.

5 Gaucher A, Péré P, Régent D, Grandhay P, Aussedat R, Vivard T. Spondylarthropathies ou polyarthrites ossifiantes. *Rev Rhum Mal Osteoart* 1987; 54: 243-8.

6 Valouch P, Pazderka V, Gatterova J. Enthesopathy of the hip. *Clin Rheumatol* 1985; 4: 150-4.

7 Littlejohn G O. More emphasis on the entheses (editorial). *J Rheumatol* 1989; 16: 1020-2.

8 Gaucher A, Péré P, Gillet P. From ankylosing spondylitis to Forestier's disease: ossifying enthesopathy, an unifying concept. *J Rheumatol* 1990; 17: 854-6.

Plasma viscosity in giant cell arteritis

We read with great interest the paper by Gudmundsson *et al*.¹

It was interesting that at follow up there was evidence that plasma viscosity and the erythrocyte sedimentation rate (ESR) paralleled clinical findings and predicted flare ups better than other variables.

In previous studies, we have tried to establish a definition of biological parameters for monitoring giant cell arteritis (GCA) and polymyalgia rheumatica (PMR).^{2,3} Our studies have shown that in patients with GCA or PMR there is a positive correlation between ESR, haptoglobin and orosomucoid during the acute phase of the disease before treatment.

Under corticosteroid treatment, a correlation persists between clinical symptoms and ESR. But a discrepancy exists between ESR and acute phase proteins: haptoglobin and orosomucoid. These two proteins remain elevated while clinical symptoms and ESR have returned to normal. In individual cases this elevation is linked to a persistence of the disease and could lead to a possible flare up when attempting to taper corticosteroid treatment. Moreover, an elevation of orosomucoid and haptoglobin can signal a clinical flare up.

Our data suggest that acute phase protein elevation that is present in spite of improvement in clinical symptoms and ESR is a contraindication for tapering corticosteroid doses or for withdrawing treatment.

J B PAOLAGGI
Service of Rheumatology,
Hôpital A. Paré, 9 avenue Charles de Gaulle, 92100
Boulogne, France

D CHAOUAT
Service of Internal Medicine, Fondation A de
Rothschild, 25-29 rue Manin, 75019 Paris, France

1 Gudmundsson M, Nordborg E, Bengtsson B-Å, Bjelle A. Plasma viscosity in giant cell arteritis as a predictor of disease activity. *Ann Rheum Dis* 1993; 52: 104-9.

2 Paolaggi JB, Chaouat D, Barres D, Hoffman H, Auquier L. Etude des variations comparées de la vitesse de sédimentation, de l'haptoglobine et de l'orosomucoïde au cours de l'évolution des pseudopolyarthrites rhizoméliques et arthérites temporales. Tentative de définition des paramètres biologiques de ces affections. *Rev Rhum* 1982; 49: 413-19.

Histological specimen of femoral head showing enthesitis of the ligamentum teres with subchondral osteitis (haematoxylin and eosin stain).

- 3 Paolaggi J B, Chaouat D, Auquier L. An additional test for the diagnosis and monitoring of giant cell arteritis and polymyalgia rheumatica. *Arth Rheum* 1985; **28**: 837-8.

Rheumatoid nodules

The recent Leader on rheumatoid nodules by Veys and De Keyser¹ gave a very interesting and useful review of the subject. When I was working as a medical registrar with Dr Bernard Schlesinger at the Royal Northern Hospital in pre-war London, one of my tasks was to carry out routine nodule counts on children with rheumatic fever (a not uncommon disorder in those days). Dr Schlesinger had found that these nodules, particularly in younger children with more serious disease, carried a worse cardiac prognosis.² At that time there was considerable interest in the possible relation of rheumatic fever to rheumatoid arthritis. A recent report by Fink³ considers post-streptococcal reactive arthritis to be a form of

rheumatic fever, though arthritis or arthralgia is of a more prolonged nature appearing early after infection.

Massell *et al*⁴ reported in 1937 that nodules closely resembling those of rheumatic fever could be produced by injecting 3 cc of a patient's blood subcutaneously into the soft tissue of an elbow followed by rubbing the injected area six times daily; nodules appeared in 37 of 82 of their patients (45%). I was unable to confirm these findings⁵ in 40 children with rheumatic fever in acute or convalescent stages, five children with chronic juvenile arthritis (Still's disease), and 10 adults with rheumatoid arthritis, three of whom had already developed rheumatoid nodules over one elbow. Veys and De Keyser report that rheumatoid nodules are found most commonly at sites of mechanical irritation and friction and we found the same to be true of nodules in rheumatic fever, but could not produce them experimentally.

Considerable interest was shown during the 1930s and 1940s in rheumatoid nodule formation. Collins,⁶ Parkes Weber⁷ in the United Kingdom, and Horowitz⁸ in South Africa had worked and written on the subject,

but my efforts to reproduce them experimentally failed miserably!

F DUDLEY HART
24 Harcourt House,
20 Harley Street,
London W1N 1AN,
United Kingdom

- 1 Veys E M, De Keyser F. Rheumatoid nodules: differential diagnosis and immunological findings. *Ann Rheum Dis* 1993; **52**: 625-6.
- 2 Schlesinger B E. The public health aspects of heart disease in children. *Lancet* 1938; **1**: 593-9, 649-54.
- 3 Fink C W. The role of the streptococcus in post-streptococcal reactive arthritis in childhood polyarteritis nodosa. perspectives in paediatric rheumatology. *J Rheumatol* 1990; **18**: (Suppl 29), 14-20.
- 4 Massell B F, Mote J R, Jones T D. *Jour Clin Invest* 1937; **16**: 124-30.
- 5 Hart F D. Rheumatic subcutaneous nodule formation. *Ann Rheum Dis* 1939; **1**: 196-200.
- 6 Collins D H. Observations on the pathology of acute rheumatism and rheumatoid arthritis. *Ann Rheum Dis* 1939; **1**: 38-45.
- 7 Parkes Weber F. The nodules and lymph gland enlargement in rheumatoid arthritis *Ann Rheum Dis* 1944-5; **4**: 3-10.
- 8 Horowitz M. The subcutaneous nodules of chronic rheumatoid arthritis: their clinical and pathological features. *Clinical Proceedings* 1949; **8**: 73-116.