Circulating concentrations of soluble L-selectin (CD62L) in patients with primary Sjögren's syndrome

Mario García-Carrasco, Pilar Pizcueta, Ricard Cervera, Manuel Ramos-Casals, Antoni Sisó, Gloria de la Red, Miguel Ingelmo, Josep Font, Pablo Engel

Abstract

Objective—Serum concentrations of soluble (s) L-selectin (CD62L) were measured in patients with primary Sjögren's syndrome (SS) to relate these concentrations to clinical and immunological features of SS.

Methods—The study included 40 consecutive patients (38 women and two men) with a mean age of 61 years (range 24–78) who fulfilled four or more of the preliminary diagnostic criteria for SS proposed by the European Community Study Group in 1993, and 33 healthy blood donors from the hospital blood bank. A sandwich enzyme linked immunosorbent assay (ELISA) was used to detect the soluble form of human sL-selectin (CD62L).

Results—The mean (SEM) values of sL-selectin (CD62L) were 861 (66) µg/ml for patients with SS and 986 (180) µg/ml for healthy blood donors, but there was no significant difference. In patients with primary SS, serum sL-selectin (CD62L) concentrations were significantly higher in patients with Raynaud's phenomenon (1275 (112) µg/ml versus 789 (69) µg/ml, $p=0.007$), autoimmune thyroiditis (1162 (113) µg/ml versus 787 (69) µg/ml, $p=0.02$) and rheumatoid factor (993 (95) µg/ml versus 684 (70) µg/ml, $p=0.01$) when compared with patients without these features.

Conclusion—The presence of Raynaud's phenomenon, autoimmune thyroiditis and rheumatoid factor is associated with higher concentrations of circulating sL-selectin (CD62L) in the sera of patients with primary SS.

(Ann Rheum Dis 2000;59:297–299)

Sjögren's syndrome (SS) is an autoimmune disease that mainly affects exocrine glands and that usually presents as a persistent dryness of the mouth and the eyes resulting from functional impairment of the salivary and lacrimal glands.¹ The spectrum of the disease extends from an organ specific autoimmune disease (autoimmune exocrinopathy)² to a systemic process (musculoskeletal, pulmonary, gastric, haematological, vascular, dermatological, renal and nervous system involvement).³ In the absence of an associated systemic autoimmune disease, patients with this condition are classified as having "primary" SS. Because of this heterogeneity, attempts have been made to

identify subsets of patients that would permit more reliable prediction of the course of primary SS in the affected people.^{4,5}

Adhesion molecules regulate cell-cell and cell-matrix interactions and determine lymphocyte homing to different lymphoid organs and sites of inflammation.⁶ Such molecules have been classified into several families, of which the selectins are one.⁷ L-selectin (CD62L) mediates leucocyte rolling on vascular endothelium at sites of inflammation and the recirculation of lymphocytes through the binding to high endothelial venules of peripheral lymph nodes.^{7,8} Indeed, leucocytes from mice lacking L-selectin (CD62L) have deficiencies in both functions.⁹ L-selectin (CD62L) is constitutively expressed on the surface of most leucocytes, including lymphocytes, neutrophils, monocytes, eosinophils, haematopoietic progenitor cells, and immature thymocytes. L-selectin (CD62L) is cleaved enzymatically from the leucocyte surface membrane after cell activation and released into the bloodstream.¹⁰ Soluble (s) L-selectin (CD62L) is functionally active and is present at high concentrations in human serum.¹¹

Primary SS is associated with lymphocytic infiltration in salivary, lacrimal and other exocrine glands. The mechanism of lymphocytic infiltration is not known. The interaction between adhesion molecules, expressed on the surface of lymphocytes, and their specific ligands, expressed on endothelial cells and subsequently other cells in exocrine glands, may be the main mechanism for lymphocyte infiltration into sites of inflammation in this disease, contributing to the retention of lymphocytes in target organs of the autoimmune attack. No information is yet available on the physiological and clinical significance of the circulating shed forms of adhesion molecules in primary SS. This study was undertaken to determine sL-selectin (CD62L) concentrations in patients with primary SS and to evaluate the relation with the clinical and immunological features of SS.

Methods

PATIENTS

We included 40 consecutive patients (38 women and two men) with a mean age at onset of SS of 51 years (range 24–78). All patients fulfilled four or more of the preliminary diagnostic criteria for SS proposed by the European Community Study Group in 1993.¹² All patients underwent a complete history and

Systemic Autoimmune Diseases Unit, Department of Medicine, Institut d'Investigacions Biomèdiques August Pi i Sunyer, Hospital Clínic, School of Medicine, University of Barcelona, Barcelona, Catalonia, Spain

M García-Carrasco
R Cervera
M Ramos-Casals
A Sisó
G de la Red
J Font
M Ingelmo

Hepatic Haemodynamic Laboratory, Liver Unit
P Pizcueta

Immunology Unit, Department of Cellular Biology and Pathology
P Engel

Department of Rheumatology, Denervita Universidad Autonoma de Puebla, Puebla, Mexico
M García-Carrasco

Correspondence to:
Dr Font, Unitat de Malalties Autoimmunes Sistèmiques, Hospital Clínic, C/Villarroel, 170, 08036-Barcelona, Catalonia, Spain

Accepted for publication
22 November 1999

Figure 1 Serum concentrations of soluble L-selectin in patients with Sjögren's syndrome and controls. Box plots indicate the overall range (error bars), 25–75% range (boxes) and median value (horizontal lines).

physical examination, and diagnostic tests for SS were applied according to the recommendations of the European Community Study Group.¹² Ten (25%) patients had disease limited to the exocrine glands, and the remaining 30 (75%) had one or more of the following extraglandular manifestations: arthralgia and/or arthritis, Raynaud's phenomenon, interstitial pneumonitis (evidence of interstitial fibrosis on chest radiography and/or in the high resolution computed tomography), cutaneous vasculitis (demonstrated by cutaneous purpura and skin biopsy), autoimmune thyroiditis (positive autoantibodies and/or altered thyroid function), peripheral neuropathy (demonstrated by the clinical picture and nerve conduction tests) and liver involvement (altered liver function tests and/or liver biopsy).

The serum concentrations of sL-selectin (CD62L) of these patients were compared with those of the 33 healthy blood donors from our hospital blood bank.

LABORATORY STUDIES

A sandwich enzyme linked immunosorbent assay (ELISA) to detect the soluble form of human L-selectin (CD62L) was performed as previously described, with slight modifications.¹¹ Anti-L-selectin (CD62L) mAb LAM1-101 diluted to 3 µg/ml was used as the capture antibody. The presence of sL-selectin was shown with biotinylated anti-LAM1-116 monoclonal antibody at 3 µg/ml and avidin-horseradish peroxidase, with O-phenylene-diamine (0.125% wt/vol, Sigma, St Louis, MO) as substrate. Human serum samples were run in triplicate and diluted at 1/40 to obtain a measurement in the linear range of our assay. Absorbances at 450 nm were measured using an MRX Microplate ELISA reader (Dynatech, Denkendorf, Germany). The concentration of sL-selectin was determined by using a standard curve constructed for each ELISA plate with recombinant L-selectin (CD62L) and results are expressed as µg/ml.

Immunological tests included determination of antinuclear antibodies (ANA) by indirect immunofluorescence using mouse liver as substrate, antibodies to double stranded DNA by

Table 1 Mean serum sL-selectin (CD62L) in SS patients grouped according to the presence or absence of extraglandular and immunological features

	<i>sL-selectin (mean (SEM))</i>		<i>p</i>
	<i>Present</i>	<i>Absent</i>	
<i>Extraglandular manifestation</i>			
Articular involvement	766 (89)	958 (96)	NS
Raynaud's phenomenon	1275 (112)	789 (69)	0.007
Cutaneous vasculitis	936 (198)	843 (68)	NS
Liver involvement	839 (162)	868 (73)	NS
Autoimmune thyroiditis	1162 (113)	787 (72)	0.02
Interstitial pneumonitis	1286 (247)	839 (67)	NS
Peripheral neuropathy	863 (204)	861 (67)	NS
<i>Immunological manifestation</i>			
ANA	817 (76)	936 (124)	NS
Anti-Ro/SS-A	891 (126)	838 (81)	NS
Anti-La/SS-B	952 (158)	841 (75)	NS
RF	993 (95)	684 (70)	0.01
Cryoglobulins	1016 (236)	848 (75)	NS
Hypocomplementaemia	1055 (286)	768 (71)	NS

Farr's technique, precipitating antibodies to the extractable nuclear antigens Ro/SS-A and La/SS-B by counterimmunoelectrophoresis and rheumatoid factor (RF) by latex fixation and Waaler-Rose tests. Complement factors (C3 and C4) were estimated by nephelometry (Behring BNA nephelometer). Serum cryoglobulinaemia was measured after centrifugation. Centrifuged serum was incubated at 4°C for seven days and examined for cryoprecipitation.

STATISTICAL ANALYSIS

We used conventional χ^2 and Fisher's exact test to analyse qualitative differences, Student's test for the comparison of means in large samples of similar variance, and the non-parametric Mann-Whitney U test for small samples. A value of $p < 0.05$ was taken to indicate statistical significance. This statistical analysis was performed by means of the SPSS program using the information stored in the database program.

Results

We measured the serum concentrations of sL-selectin (CD62L) in 40 patients with primary SS and in 33 healthy blood donors. The mean (SEM) values of sL-selectin (CD62L) were 861 (66) µg/ml for patients with SS and 986 (180) µg/ml for healthy blood donors, but there was no significant difference (fig 1).

The most common extraglandular manifestations in our patients with primary SS were articular involvement in 20 (50%) patients, liver involvement in nine (23%), peripheral neuropathy in eight (20%), cutaneous vasculitis in eight (20%), autoimmune thyroiditis in eight (20%), Raynaud's phenomenon in six (15%) and interstitial pneumonitis in two (5%). Serum sL-selectin (CD62L) concentrations were significantly higher in patients with Raynaud's phenomenon (1275 (112) µg/ml *v* 789 (69) µg/ml, $p=0.007$) and autoimmune thyroiditis (1162 (113) µg/ml *v* 787 + 69 µg/ml, $p=0.02$) when compared with patients without these manifestations (table 1).

The most frequent immunological features were: ANA in 25 (63%) patients, RF in 23 (58%), anti-Ro/SS-A antibodies in 13 (33%),

cryoglobulinaemia in 6 of 34 (18%), hypocomplementaemia in 5 of 30 (17%) and anti-La/SS-B antibodies in five (13%) patients. No patient had a positive titre of anti-dsDNA or anti-Sm antibodies. Serum sL-selectin (CD62L) concentrations were significantly higher in patients with RF compared with those without RF (993 (95) $\mu\text{g/ml}$ compared with 684 (70) $\mu\text{g/ml}$, $p=0.01$) (table 1).

Discussion

Investigation into adhesion molecule pathophysiology in primary SS is scarce and has focused on characterising adhesion molecule expression in salivary gland biopsy specimens.¹³ L-selectin was found to be expressed by 18% of the infiltrating cells in labial salivary glands from patients with primary SS,¹³ which is considerably less than in the peripheral blood, and the comparatively reduced number of CD62 L positive lymphocytes might be attributable to shedding of the L-selectin (CD62L) after cell activation.

In our study, we found increased concentrations of circulating sL-selectin (CD62L) in patients with primary SS that present with some extraglandular manifestations. Firstly, patients with primary SS and Raynaud's phenomenon showed higher circulating concentrations of sL-selectin (CD62L). In primary SS, Raynaud's phenomenon is not constantly observed, and it could result from a specific immunological aggression. The molecule analysed in our study promotes the attachment of diverse leucocytes to endothelium, and thereby together regulates the local traffic of neutrophils, monocytes, and lymphocytes. L-selectin is shed after cell activation, and raised sL-selectin (CD62L) concentrations may reflect a higher grade of activation of the leucocytes occurring in this subset of patients with primary SS and associated Raynaud's phenomenon.

A second subset of patients with primary SS, those with autoimmune thyroiditis, showed higher values of circulating sL-selectin (CD62L) in their sera. Autoimmune thyroiditis is an organ specific, autoimmune destruction of the thyroid gland mediated by T lymphocytes, with anti-thyroglobulin and anti-microsomal antibody formation, frequently associated to SS.¹⁴ Stimulation of thyroid epithelial cells in vitro with tumour necrosis factor α or interferon γ induced expression of ICAM-1.¹⁵ ICAM-1 and LFA-3 expression has been identified on thyroid cells in autoimmune

thyroiditis, and it has been suggested that these molecules have functional importance in leading to the adherence of T cells to this target organ.¹⁶ Adhesion molecules might participate in the pathophysiology of autoimmune thyroiditis in primary SS patients.

The presence of Raynaud's phenomenon, autoimmune thyroiditis and rheumatoid factor is associated with higher concentrations of circulating sL-selectin (CD62L) in the sera of patients with primary SS. Measurement of sL-selectin (CD62L) in primary SS may also provide guidance in the selection of clinical subsets of patients in which pharmacological blockade of adhesion molecules should be attempted, to interfere with leucocyte migration and function.

Funding: supported by grants FIS 97/0669 and FIS 99/0280 from the Fondo de Investigaciones Sanitarias. M Ramos-Casals is a Research Fellow sponsored by a grant from the Hospital Clinic.

- 1 Moutsopoulos HM, Chused TM, Mann DL, Klippel JH, Fauci AS, Frank MM, *et al.* Sjögren's syndrome (sicca syndrome): Current issues. *Ann Intern Med* 1980;92:212-26.
- 2 Talal N. Clinical and pathogenetic aspects of Sjögren's syndrome. *Semin Clin Immunol* 1993;6:11-20.
- 3 Anaya JM, Gutierrez M, Espinosa LR. Síndrome de Sjögren primario. Manifestaciones clínicas extraglandulares. *Rev Esp Reumatol* 1994;21:337-42.
- 4 Ramos M, Cervera R, García-Carrasco M, Miret C, Muñoz FJ, Espinosa G, *et al.* Síndrome de Sjögren primario: características clínicas e inmunológicas en una serie de 80 pacientes. *Med Clin (Barc)* 1997;108:652-7.
- 5 Ramos-Casals M, Cervera R, Font J, García-Carrasco M, Espinosa G, Reino S, *et al.* Young onset of primary Sjögren's syndrome: clinical and immunological characteristics. *Lupus* 1998;7:202-6.
- 6 Springer TA. Adhesion receptors of the immune system. *Nature* 1990;346:425-34.
- 7 Springer TA. Traffic signals for lymphocyte recirculation and leukocyte emigration: the multistep paradigm. *Cell* 1994;76:301-14.
- 8 Tedder TF, Steeber DA, Chen A, Engel P. The selectins vascular adhesion molecules. *FASEB J* 1995;9:866-73.
- 9 Tedder TF, Steeber DA, Pizcueta P. L-selectin-deficient mice have impaired leukocyte recruitment into inflammatory sites. *J Exp Med* 1995;181:2259-64.
- 10 Chen A, Engel P, Tedder TF. Structural requirements regulate endoproteolytic release of the L-selectin (CD62L) adhesion receptor from the cell surface of leukocytes. *J Exp Med* 1995;182:519-30.
- 11 Spertini O, Schleiffenbaum B, White-Owen C, Ruiz Jr P, Tedder TF. ELISA for quantitation of L-selectin shed from leukocytes in vivo. *J Immunol Methods* 1992;156:115-23.
- 12 Vitali C, Bombardieri S, Moutsopoulos HM, Balestrieri G, Bencivelli W, Bernstein RM, *et al.* Preliminary criteria for the classification of Sjögren's syndrome. *Arthritis Rheum* 1993;36:340-7.
- 13 Aziz KE, McCluskey PJ, Wakefield D. Expression of selectins (CD62 E,L,P) and cellular adhesion molecules in primary Sjögren's syndrome: questions to immunoregulation. *Clin Immunol Immunopathol* 1996;80:55-66.
- 14 Cervera R, Ramos M, García-Carrasco M, Font J, Gaya J, Ingelmo M. Prevalence of thyroid disease in a series of 92 patients with primary Sjögren's syndrome. *Br J Rheumatol* 1997;36 (suppl 1):46.
- 15 Weetman AP, Cohen S, Makgoba MW, Borysiewicz LK. Expression of an intercellular adhesion molecule, ICAM-1, by human thyroid cells. *J Endocrinol* 1989;22:185-91.
- 16 Tandon N, Makgoba MW, Gahmberg CG, Weetman AP. The expression and role in T cell adhesion of LFA-3 and ICAM-2 on human thyroid cells. *Clin Immunol Immunopathol* 1992;64:30-5.