Characterisation of *Eubacterium* cell wall: peptidoglycan structure determines arthritogenicity

X Zhang, M Rimpiläinen, E Šimelyte, P Toivanen

Abstract

Objective—To elucidate factors involved in the arthritogenicity of bacterial cell walls.

Methods—For characterisation of an arthritogenic *Eubacterium aerofaciens* cell wall, peptidoglycan-polysaccharide (PG-PS) polymers were isolated by removing cell wall associated proteins (CWPs), PG and PS moieties were separated, and an attempt was made to de-*O*-acetylate PG-PS. The cell wall of *E. limosum* was used as a non-arthritogenic control. The chemical composition of these cell wall preparations was analysed by gas chromatography-mass spectrometry. Also, their ability to resist lysozyme degradation and to sustain experimental chronic arthritis was tested.

Results—The observations made with the cell wall of *E. aerofaciens*, an anaerobic habitant of the human intestine, were compared with those reported from a pathogenic *Streptococcus*, showing that in both strains a complex consisting of PG-PS is required for the induction of chronic arthritis. The PS moiety most probably protects PG from enzyme degradation, allowing prolonged tissue persistence and leading to the chronic synovial inflammation. CWPs attached to PG-PS are not necessary for this function. *O*-Acetylation of PG, which is required for arthritogenicity of the streptococcal cell

wall, seems not to be present in the arthritogenic *E. aerofaciens* PG or only occurs to a small degree; attempts to de-*O*-acetylate the *E. aerofaciens* cell wall did not affect its arthritogenicity or lysozyme resistance.

Conclusion—The results obtained indicate that the source of bacterial cell wall plays no part in the chemical or structural requirements for PG to induce chronic cell wall arthritis in the rats; the chemical structure of the PG moiety is decisive.

(Ann Rheum Dis 2001;60:269–274)

Turku Immunology
Centre, Department of
Medical Microbiology,
Turku University,
Turku, Finland
X Zhang
M Rimpiläinen
E Šimelyte
P Toivanen

Correspondence to:
Dr X Zhang, Department of
Medical Microbiology, Turku
University, Kiinamyllynkatu
13, FIN-20520 Turku,
Finland
xzhang@utu.fi

Accepted 1 August 2000

Figure 1 Gram positive bacterial cell wall contains mainly peptidoglycan (PG), polysaccharide (PS), and cell wall associated proteins (CWPs). The CWPs outside the PS and PG layers can be removed by proteolytic enzymes, whereas the proteins inside the PG layer are difficult to remove without breaking the PG structure. PS binds covalently to the PG layer, and protects PG polymers from the degrading enzymes. PG consists of several layers (up to 70) of sugar chains composed of *N*-acetylglucosamine and *N*-acetylmuramic acid alternately linked each other. Short peptides bound to the *N*-acetylmuramic acid (muramyl peptides) connect the sugar chains. PG and the muramyl peptides possess multiple biological activities.

The intestinal flora of patients with newly developed rheumatoid arthritis (RA) is significantly different from that of non-RA controls.^{1,2} This difference is primarily due to anaerobic bacterial species—namely, to *Eubacterium aerofaciens*.¹ On the other hand, a single intraperitoneal (IP) injection of cell walls isolated from this Gram positive bacterium, a normal habitant of the human gut, induces a chronic arthritis in Lewis rats, whereas cell walls from a related *E. limosum* are non-arthritogenic.^{3,4}

The Gram positive bacterial cell wall skeleton mainly contains peptidoglycan (PG), polysaccharide (PS), and the cell wall associated proteins (CWPs) (fig 1). The CWPs in the Gram positive cell wall extend from the plasma membrane through the PG and PS layers to the cell surface and are covalently attached to the PG.^{5,6} They interact with the extracellular environment and contribute to the development of bacterial infections. CWPs of *Staphylococcus aureus* interact with specific host molecules either by promoting attachment to the host tissues or by preventing the phagocytic clearance.⁷ CWPs of *Mycobacterium leprae* are the major stimulants of the cell mediated immunity,⁸ and CWPs of *M. tuberculosis* are known to play a part in establishing protective immunity.⁹ The M protein of the group A streptococcal CWPs is a major virulence factor with superantigenic activity.¹⁰ Group A streptococcal M protein has shared epitopes with the articular cartilage and synovium,¹¹ and systemic administration of streptococcal cell wall induces a chronic, destructive arthritis in susceptible rat strains.¹²

The bacterial cell wall PG-PS complex can initiate and sustain a chronic destructive inflammation.¹³ PG-PS isolated from human intestinal flora bacteria can induce chronic arthritis in rats.^{14,15} The arthritogenic ability of cell wall constituents is dependent on resistance to biodegradation and consequent persistence in the tissues.¹³ The PS moiety

covalently bound to PG is essential for the PG-PS complexes to persist and induce chronic inflammation. Streptococcal PG-PS did not sustain chronic arthritis when the PS moiety was removed,¹⁶ and PS alone induced only mild oedema in the joints.¹⁷ Another structural basis for streptococcal arthritogenicity is the presence of an acetyl moiety at the C-6 hydroxyl group of *N*-acetylmuramyl residues of PG, producing the corresponding 2,6-diacetylmuramyl PG (*O*-acetylated PG).¹⁸ Gonococcal *O*-acetylated PG is more resistant to enzyme degradation than non-*O*-acetylated PG,¹⁹ and subsequently persists in the tissues longer, leading to chronic arthritis.²⁰ De-*O*-acetylated group A streptococcal PG-PS induced less severe chronic arthritis than native PG-PS.²¹ In this study we aimed at defining the arthritogenic component(s) of the *E. aerofaciens* cell wall to determine whether the same requirements are valid for a representative of the normal human gut flora and a pathogenic *Streptococcus*.

Methods

BACTERIA

Eubacterium aerofaciens ATCC 25986 was obtained from the Culture Collection, University of Gothenburg, Sweden, with a nomination CCUG 28087. *E. limosum* ATCC 8486 was purchased from the American Type Culture Collection, Rockville, MD, USA. Both strains were grown overnight under strictly anaerobic conditions at 37°C in BBL Schaedler Broth (Becton Dickinson Microbiology Systems, Cockeysville, MD, USA) to the late logarithmic phase. They were characterised by 16S rDNA sequence analysis as described by Jalava *et al.*²²

CRUDE CELL WALL PREPARATION

Bacterial crude cell walls were isolated as described previously.²³ Briefly, the cells were broken with glass beads (diameter 0.45–0.50 mm) in an MSK cell homogeniser (B Braun Melsungen AG, Melsungen, Germany). The cell walls were collected by centrifugation, treated with deoxyribonuclease I (1 mg/g wet weight), ribonuclease A (10 mg/g wet weight), and trypsin (20 mg/g wet weight), washed, and sonicated in an ice bath for 75 minutes (Branson Sonifier, Smith Kline Co, Danbury, Conn., USA). The sonicated cell wall suspension was centrifuged at $10\,000 \times g$, 4°C, for 20 minutes. The supernatant was centrifuged by ultracentrifugation (Sorvall Ultracentrifuge OTD65B, rotor 60 Ti, Du Pont Co, Wilmington, USA) at $100\,000 \times g$, 4°C, for 60 minutes. The pellet containing the cell wall was suspended in water. For chemical analysis the cell walls were dialysed in membrane (MWCO 3500, Spectrum, CA, USA) against distilled water. For IP injection into rats the cell wall suspension was sterilised by heating at 90°C for 30 minutes.²⁴ The sterility was checked by cultures on agar plates at 37°C and at room temperature under aerobic and anaerobic conditions; no bacterial growth was detected after two days of culture. The endotoxin tests by E-TOXATE (Sigma

Chemical Co, MO, USA) were also found to be negative. These procedures were applied for all cell wall preparations used.

PG-PS POLYMERS

To obtain PG-PS polymers, the CWP were removed; the crude cell walls were treated with proteolytic enzymes, protease (Subtilisin Carlsberg, 5 mg/g wet weight), proteinase K (2 mg/g wet weight), pepsin (20 mg/g wet weight), and papain (5 mg/g wet weight). All enzymes were purchased from Sigma Chemical Co, MO, USA. After lyophilisation the cell walls were extracted with chloroform-methanol (2:1 vol/vol, 40 mg/ml) at room temperature for four hours to remove lipoteichoic acid or glycosyl glycerides.¹⁶ After washing once with water and dialysis against distilled water, the cell walls were treated by 8 M lithium chloride at room temperature for one hour to remove any non-covalently bound CWPs.²⁵ The cell walls were collected by ultracentrifugation and washed twice with distilled water. They were extracted with 0.1 M EDTA at room temperature for one hour to remove any divalent cation stabilised oligomeric proteins.²⁶ After ultracentrifugation and washing twice with distilled water the cell walls were suspended in acetone at room temperature for one hour and further mixed at 4°C for 30 hours to remove possible contaminating proteins and endotoxin.^{27,28} The cell walls were collected by ultracentrifugation, washed three times with distilled water, and dialysed against distilled water. The preparations thus obtained are PG-PS complexes as shown by gas chromatography-mass spectrometry (GC-MS) analysis.

PS AND SOLUBLE PG POLYMERS

PG and PS polymers were isolated from *E. aerofaciens* ATCC 25986 as described previously for *Lactobacillus*.²⁹ Briefly, the lyophilised cell walls were extracted with 10% trichloroacetic acid at 60°C for four hours. The suspension was centrifuged at $38\,800 \times g$ at 4°C for 30 minutes and washed once with distilled water. The supernatant contained the PS fraction, and the pellet the PG fraction. The soluble PG was obtained as described previously.²⁴ The PG fraction was lyophilised and suspended in water at 10 mg/ml and sonicated (Branson Sonifier) for 180 minutes (18×10 -minute cycles with breaks for cooling between the cycles) in an ice bath. The remaining insoluble PG fraction was removed by centrifugation at $10\,000 \times g$ at 4°C for 30 minutes. The supernatant (soluble PG fraction) was collected and dialysed against distilled water for 48 hours.

DE-O-ACYLATION OF CELL WALL

The treatment of the cell walls for de-*O*-acylation was done as described previously.²¹ Briefly, the cell walls were lyophilised, suspended in 0.05 M sodium hydroxide at 37°C for four hours. After washing three times with distilled water, the cell walls were fragmented by sonication and subsequently separated by differential centrifugation as described above for the cell wall preparation.

Table 1 Gas chromatography-mass spectrometry analysis of the cell wall preparations

Component	<i>E. aerofaciens</i>				<i>E. limosum</i>	
	Crude cell wall	PG-PS*	PG	PS	Crude cell wall	PG-PS
Carbohydrates						
Sugars						
Rhamnose	18.5	18.0	—	30.3	20.4	29.0
Glucose + galactose	30.9	27.0	1.0	50.1	14.9	3.8
Amino sugars						
<i>N</i> -Acetylmuramic acid	1.9	9.2	11.5	0.6	2.9	13.1
<i>N</i> -Acetylglucosamine	7.6	14.3	9.1	11.1	2.1	7.2
<i>N</i> -Acetylgalactosamine	—	—	—	—	3.9	9.0
Amino acids						
Alanine	5.0	4.9	20.4	0.3	3.6	3.5
Glutamic acid/glutamine	6.0	8.1	32.8	1.0	8.0	10.1
Lysine	9.7	6.4	24.2	0.3	9.0	4.6
Aspartic acid/asparagine	4.8	6.9	17.0	0.4	2.7	0.3
Ornithine	<0.1	<0.1	0.3	0.1	12.1	18.7
Serine	<0.1	—	0.1	<0.1	3.4	9.6
Leucine	0.4	<0.1	0.4	<0.1	2.5	<0.1
Isoleucine	0.4	<0.1	0.7	—	2.0	0.2
Phenylalanine	0.2	<0.1	0.7	<0.1	1.4	<0.1
Proline	0.3	<0.1	0.5	0.2	1.4	<0.1
Tyrosine	0.3	<0.1	0.3	<0.1	1.0	<0.1
Valine	0.4	—	—	—	1.6	<0.1

Each component is expressed as a percentage of the dry weight of cell wall. Figures in bold indicate PG amino sugars and amino acids. Norleucine, methionine, tryptophan were applied as internal standards for amino acid analysis. Cystine, glycine, threonine, diaminopimelic acid, cysteine, arginine, and histidine were not found.

*PG-PS = peptidoglycan-polysaccharide.

Figure 2 (A) A single intraperitoneal (IP) injection of *E. aerofaciens* peptidoglycan-polysaccharide (PG-PS; 150 µg/g body weight of rat) induced a more severe chronic arthritis than that induced with a crude cell wall (140 µg/g body weight of rat) ($p < 0.02$ by non-parametric Mann-Whitney U test). Each black symbol represents a mean (SEM) of six rats. Each white symbol represents a mean (SEM) of 11 rats. (B) Rats injected with *E. limosum* PG-PS (200 µg/g body weight of rat) IP or with the crude cell wall (200 µg/g body weight of rat) developed only mild acute arthritis. Each black symbol represents a mean (SEM) of eight rats. Each white symbol represents a mean (SEM) of six to 12 rats.

GAS CHROMATOGRAPHY-MASS SPECTROMETRY (GC-MS)

The chemical composition of the cell wall preparations was analysed by GC-MS as described previously.³⁰ The derivatised molecules were ionised by the electron impact method and analysed in the selected ion monitoring mode using single positive ions at a mass to charge ratio (m/z). Sugars were analysed as alditol acetate derivatives with the same ions as monitored by Gilbert *et al.*³⁰ except m/z 289 for rhamnose. Fucose was used as internal standard for rhamnose, allose for glucose/galactose, and *N*-methyl-D-glucamine for muramic acid. Amino acids were analysed as butyl heptafluorobutyl derivatives with the same ions as used by Gilbert *et al.*³⁰ L-Norleucine was used as internal standard for glycine, serine, alanine, leucine, cysteine, threonine, isoleucine, and valine; L-methionine for phenylalanine, glutamic acid/glutamine, proline, ornithine, lysine, aspartic acid/asparagine, and tyrosine; L-tryptophan for cystine, diaminopimelic acid, histidine, and arginine. One microlitre of the derivative was injected in the pulsed splitless mode and analysed by a gas chromatograph (model HP 5890A; Hewlett-Packard, Wilmington, DE) equipped with a fused silica capillary column (SE-54; Nordion Instruments, Helsinki) and coupled directly with a TRIO-1 mass spectrometer (VG Instruments, Manchester, UK). The oven temperature for sugars was started at 50°C, programmed to 270°C at a rate of 10°C/min, and held for one minute; finally, the oven was heated to 290°C for five minutes. For amino acids the oven temperature was started at 85°C, programmed to 280°C at a rate of 10°C/min, and held for one minute; finally, the oven was heated to 290°C for five minutes.

ANIMALS AND INDUCTION OF ARTHRITIS

All the animal experiments were done in concordance with the national and international

Figure 3 (A) Rats injected with *E. aerofaciens* peptidoglycan (PG; 33 μ g/g body weight of rat) intraperitoneally (IP) deprived of polysaccharide (PS), developed only acute arthritis. Rats injected with *E. aerofaciens* PS alone (122 μ g/g body weight of rat) did not develop any arthritis. Each symbol represents a mean (SEM) of five rats. (B) Rats injected with *E. aerofaciens* de-O-acylated cell wall (140 μ g/g body weight of rat) IP developed arthritis similar to that seen after injection of the crude cell wall (140 μ g/g body weight of rat). Each black symbol represents a mean (SEM) of six rats. Each white symbol represents a mean (SEM) of 11 rats.

laws and policies (Order No 1076/85, Finland; EEC Council Directive 86/609, OJL 358, Dec 1987). Inbred pathogen-free female Lewis rats weighing approximately 150 g were purchased from Harlan Sprague Dawley, Inc, Indianapolis, IN, USA. The animals were kept in Macrolon III cages with disposable filter tops (Scanbur, Denmark) and given autoclaved standard diet and water; all handling was performed in a laminar-flow hood. Arthritis was induced by a single IP injection of the cell wall preparations suspended in sterile phosphate buffered saline (PBS) on day zero. Control rats were injected with an equal volume of sterile PBS. To monitor development of arthritis, each limb was assigned a score of 0–4, based on the degree of erythema, oedema, painfulness, and functional disorder of the ankle and metatarsal joints (wrist and metacarpal joints) and judged by two independent observers as described previously.³¹ The arthritis score used in the “Results” is the sum of the score of each limb. The animals were observed for at least four weeks. Rats were killed under anaesthesia by Metofane (Pitman-Moore, Inc, Washington Crossing, NJ, USA).

SENSITIVITY TO LYSOZYME

The sensitivity of different cell wall preparations to lysozyme was tested as described previously.²¹ Suspensions of cell wall preparations (4 mg/ml) in 0.1 M sodium acetate buffer (pH 5.0) were incubated with lysozyme (400 μ g/ml) at 37°C. The optical density of the suspension was measured at wavelength 560 nm after five and 24 hours’ incubation.

STATISTICAL ANALYSIS

Differences of arthritis development between the experimental groups were analysed with the non-parametric Mann-Whitney U test. Differences were considered significant at $p < 0.05$.

Results

IDENTIFICATION OF BACTERIA

The gene coding for 16S ribosomal RNA is generally used to identify bacterial strains. To confirm the identity of the bacterial strains used in this study the genes coding for this region were partially sequenced (320–452 base pairs). Both for *E. aerofaciens* and for *E. limosum*, a 100% identity with respective sequences in the GenBank data base was confirmed. In addition, a 72.6% identity between the strains of *E. aerofaciens* and *E. limosum* was obtained.

ANALYSIS OF CELL WALL PREPARATION BY GC-MS

All bacterial cell wall preparations were analysed by GC-MS before injection into the rats (table 1). The total amount of protein amino acids is 2.2% in the *E. aerofaciens* crude cell wall, and less than 0.6% in its PG-PS polymer, in which the amount of individual protein amino acids is less than 0.1%. The amount of protein amino acids is 12.6% in the *E. limosum* crude cell wall, and less than 0.7% in its PG-PS polymer. Both bacterial PSs contain rhamnose and glucose/galactose. *E. aerofaciens* PS also contains *N*-acetylglucosamine. *N*-Acetylgalactosamine is present in the *E. limosum* cell wall and missing in the *E. aerofaciens* cell wall. PG of both cell walls contains the amino sugars *N*-acetylmuramic acid and *N*-acetylglucosamine. Arthritogenic *E. aerofaciens* has four PG amino acids: alanine, glutamic acid, lysine, and aspartic acid. Non-arthritogenic *E. limosum* cell wall has five PG amino acids: alanine, glutamic acid, lysine, ornithine, and serine (table 1). These findings are in concert with the previously reported PG types—A4a for *E. aerofaciens* and B2a for *E. limosum*.²³

INDUCTION OF ARTHRITIS

For induction of arthritis a single IP injection of the cell wall components was applied. *E. limosum* PG-PS appeared non-arthritogenic, similar to the crude cell wall of *E. limosum* (fig 2B). A high dose (200 μ g/g body weight) of PG-PS isolated from the arthritogenic *E. aerofaciens* crude cell wall was toxic, leading to death within three days (data not shown here), probably owing to the high PG content. When a lower dose (150 μ g/g body weight of rat) was used, *E. aerofaciens* PG-PS caused a significantly more severe chronic arthritis than the equivalent dose of the crude cell wall (fig 2A), suggesting that the PG-PS complex possesses

Table 2 Degradation of bacterial cell wall preparations by lysozyme

Substrate	Lysozyme 5 h		Lysozyme 24 h	
	-	+	-	+
<i>E. aerofaciens</i>				
Crude cell wall	2.0	15.6	1.9	20.7
De-O-acetylated	5.3	31.0	7.5	30.7
PG-PS	1.9	15.8	3.9	23.5
PG	0	75.9	0	79.3
<i>E. limosum</i>				
Crude cell wall	0.7	24.4	4.7	58.5

Results are given as the percentage decrease of the OD₅₆₀ after five hours and 24 hours' incubation with or without lysozyme, compared with the OD₅₆₀ at time 0. Mean values from triplicate experiments are given.

the structure decisive for arthritogenicity. However, after the PS moiety was removed, the PG alone induced only acute arthritis. Likewise, PS alone did not induce any arthritis (fig 3A). Further, attempts to de-O-acetylate the *E. aerofaciens* cell wall did not affect its arthritogenicity (fig 3B). None of the rats injected with PBS alone developed arthritis.

DEGRADATION OF BACTERIAL CELL WALL PREPARATIONS BY LYSOZYME

E. aerofaciens crude cell wall was found to be resistant to degradation by lysozyme; only 20.7% was degraded after 24 hours' treatment with this enzyme (table 2). In contrast, *E. limosum* crude cell wall was susceptible to lysozyme, with 58.5% degradation after 24 hours. De-O-acetylation of *E. aerofaciens* cell wall did not affect its susceptibility to lysozyme. *E. aerofaciens* PG-PS showed the same degree of lysozyme resistance as the crude cell wall, whereas PG alone (after removal of PS) was rapidly degraded by lysozyme; 75.9% was already degraded after five hours (table 2).

Discussion

It has been shown that synovial tissue of patients with RA may harbour traces of bacterial PG.³² The results presented in this paper, obtained with a normal bacterial habitant (*E. aerofaciens*) of the human intestine, indicate that the PG moiety of the bacterial cell wall contains a structure decisive for induction of chronic arthritis in rats. They also show that the PG-PS complex is necessary for the arthritis induction; the PS moiety most probably protects PG from enzyme digestion through steric hindrance, allowing prolonged tissue persistence and leading to the chronic synovial inflammation. Similar conclusions have previously been reached in studies on streptococcal cell wall arthritis.¹⁶ Up to this point, our observations indicate that the same principles apply for the chemical structure, determining arthritogenicity here also, independently of the bacterial species used. However, in contrast with the streptococcal cell wall, in which the O-acetylation is known to be associated with the arthritogenicity and resistance against biodegradation,²¹ O-acetylation does not seem to be a prerequisite for the arthritogenicity or biodegradation of the *E. aerofaciens* cell wall. Our findings suggest that O-acetylation does not occur in *E. aerofaciens* cell wall or only

occurs to a small degree. The gonococcal cell wall seems to resemble the streptococcal cell wall in this respect because O-acetylation is required for its arthritogenicity.^{19 20}

Lysozyme susceptibility is usually thought to be associated with arthritogenicity of the cell wall. Almost all bacterial strains studied for their ability to induce cell wall arthritis have been found to be lysozyme resistant; this includes species of *Lactobacillus*, *Streptococcus*, *Bifidobacterium*, and *Eubacterium*.³³ The only exception is the *E. limosum* cell wall, which is non-arthritogenic and which has been reported to be lysozyme resistant.³ However, our results obtained by using the same strain of *E. limosum* indicate both non-arthritogenicity and lysozyme sensitivity (table 2). Furthermore, because PG is degraded in vivo by other enzymes also (for example, N-acetylmuramyl-L-alanine amidase, N-acetylglucosaminidase, and others),^{19 34 35} it is unlikely that lysozyme resistance alone would be a decisive requirement for the ability of a cell wall to induce chronic arthritis.

It has so far remained undetermined whether CWP's attached to PG-PS contribute or are required for induction of chronic arthritis. Our results indicate that their presence is not necessary for arthritis induction. Removal of CWP's by proteolytic enzymes resulted in PG-PS which induced an even more severe chronic arthritis than that seen after injection of an equivalent amount of the crude cell wall (fig 2A). These findings parallel those showing that genetic mutation of the streptococcal M protein attached to PG does not change arthritogenicity of the cell wall.³⁶ Otherwise, results from GS-MS analyses (table 1) show very little, if any, further information about chemical or structural requirements for arthritogenicity. For instance, the presence of rhamnose does not correlate with this ability, even though such a correlation has previously been suggested.³⁷

In a previous study on four *Eubacterium* strains it was shown that the chemical structure of PG present in the bacterial cell walls is decisive in determining arthritogenicity/non-arthritogenicity.²³ The present results, based on characterisation of PG-PS complexes, further strengthen that conclusion by indicating that the source of a cell wall plays no part in the chemical or structural requirements of PG arthritogenicity; it does not matter whether the cell wall is isolated from a normal habitant of the human intestine or from a pathogenic strain of *Streptococcus*. The structure of PG is decisive. Studies with other bacterial species—for example, with *Clostridia* and *Bifidobacteria*, are needed to elucidate further the detailed structural determinants of the PG arthritogenicity.

We thank Leena Kivistö, Marja-Riitta Teräsjarvi, Karolina Karlson, and Heli Niittymäki for excellent technical assistance, Jari Jalava for 16S rDNA sequence analysis, and Yong Zhang for help in evaluating the arthritis.

This work was supported by EVO of Turku University Central Hospital.

1 Eerola E, Möttönen T, Hannonen P, Luukkainen R, Kantola I, Vuori K, et al. Intestinal flora in early rheumatoid arthritis. *Br J Rheumatol* 1994;33:1030-8.

- 2 Peltonen R, Nenonen M, Helve T, Hanninen O, Toivanen P, Eerola E. Faecal microbial flora and disease activity in rheumatoid arthritis during a vegan diet. *Br J Rheumatol* 1997;36:64–8.
- 3 Severijnen AJ, van Kleef R, Hazenberg MP, van de Merwe JP. Chronic arthritis induced in rats by cell wall fragments of *Eubacterium* species from the human intestinal flora. *Infect Immun* 1990;58:523–8.
- 4 Šimelyte E, Rimpiläinen M, Rantakokko K, Lehtonen L, Zhang X, Aho H, et al. Tissue distribution and persistence of arthritogenic and nonarthritogenic *Eubacterium* cell walls. *Clin Exp Rheumatol* 1999;17:281–8.
- 5 Pancholi V, Fischetti VA. Isolation and characterization of the cell-associated region of group A streptococcal M6 protein. *J Bacteriol* 1988;170:2618–24.
- 6 Ton That H, Faull KF, Schneewind O. Anchor structure of staphylococcal surface proteins. A branched peptide that links the carboxyl terminus of proteins to the cell wall. *J Biol Chem* 1997;272:22285–92.
- 7 Foster TJ, McDevitt D. Surface-associated proteins of *Staphylococcus aureus*: their possible roles in virulence. *FEMS Microbiol Lett* 1994;118:199–205.
- 8 Melancon Kaplan J, Hunter SW, McNeil M, Stewart C, Modlin RL, Rea TH, et al. Immunological significance of *Mycobacterium leprae* cell walls. *Proc Natl Acad Sci USA* 1988;85:1917–21.
- 9 Barnes PF, Mehra V, Hirschfield GR, Fong SJ, Abou Zeid C, Rook GA, et al. Characterization of T cell antigens associated with the cell wall protein-peptidoglycan complex of *Mycobacterium tuberculosis*. *J Immunol* 1989;143:2656–62.
- 10 Tomai M, Kotb M, Majumdar G, Beachey EH. Superantigenicity of streptococcal M protein. *J Exp Med* 1990;172:359–62.
- 11 Baird RW, Bronze MS, Kraus W, Hill HR, Veasey LG, Dale JB. Epitopes of group A streptococcal M protein shared with antigens of articular cartilage and synovium. *J Immunol* 1991;146:3132–7.
- 12 Cromartie WJ, Craddock JG, Schwab JH, Anderle SK, Yang CH. Arthritis in rats after systemic injection of streptococcal cells or cell walls. *J Exp Med* 1977;146:1585–602.
- 13 Schwab JH. Phlogistic properties of peptidoglycan-polysaccharide polymers from cell walls of pathogenic and normal-flora bacteria which colonize humans. *Infect Immun* 1993;61:4535–9.
- 14 Stimpson SA, Brown RR, Anderle SK, Klapper DG, Clark RL, Cromartie WJ, et al. Arthropathic properties of cell wall polymers from normal flora bacteria. *Infect Immun* 1986;51:240–9.
- 15 Severijnen AJ, van Kleef R, Hazenberg MP, van de Merwe JP. Cell wall fragments from major residents of the human intestinal flora induce chronic arthritis in rats. *J Rheumatol* 1989;16:1061–8.
- 16 Fox A, Brown RR, Anderle SK, Chetty C, Cromartie WJ, Gooder H, et al. Arthropathic properties related to the molecular weight of peptidoglycan-polysaccharide polymers of streptococcal cell walls. *Infect Immun* 1982;35:1003–10.
- 17 Chetty C, Brown RR, Schwab JH. Edema-producing activity of group A streptococcal polysaccharide and its possible role in the pathogenesis of cell wall-induced polyarthritis. *J Exp Med* 1983;157:1089–100.
- 18 Clarke AJ, Dupont C. O-Acetylated peptidoglycan: its occurrence, pathobiological significance, and biosynthesis. *Can J Microbiol* 1991;38:85–91.
- 19 Rosenthal RS, Folkening WJ, Miller DR, Swim SC. Resistance of O-acetylated gonococcal peptidoglycan to human peptidoglycan-degrading enzymes. *Infect Immun* 1983;40:903–11.
- 20 Fleming TJ, Wallsmith DE, Rosenthal RS. Arthropathic properties of gonococcal peptidoglycan fragments: implications for the pathogenesis of disseminated gonococcal disease. *Infect Immun* 1986;52:600–8.
- 21 Stimpson SA, Lerch RA, Cleland DR, Yarnall DP, Clark RL, Cromartie WJ, et al. Effect of acetylation on arthropathic activity of group A streptococcal peptidoglycan-polysaccharide fragments. *Infect Immun* 1987;55:16–23.
- 22 Jalava J, Mäntymaa ML, Ekblad U, Toivanen P, Skurnik M, Lassila O, et al. Bacterial 16S rDNA polymerase chain reaction in the detection of intra-amniotic infection. *Br J Obstet Gynaecol* 1996;103:664–9.
- 23 Zhang X, Rimpiläinen M, Šimelyte E, Toivanen P. What determines arthritogenicity of bacterial cell wall? A study on *Eubacterium* cell wall-induced arthritis. *Rheumatology* 2000;39:274–82.
- 24 Rosenthal RS, Dziarski R. Isolation of peptidoglycan and soluble peptidoglycan fragments. *Methods Enzymol* 1994;235:253–85.
- 25 Doyle RJ, Streips UN, Fan VS, Brown WC, Mobley H, Mansfield JM. Cell wall protein in *Bacillus subtilis*. *J Bacteriol* 1977;129:547–9.
- 26 de Maagd RA, Wientjes FB, Lugtenberg BJ. Evidence for divalent cation (Ca²⁺)-stabilized oligomeric proteins and covalently bound protein-peptidoglycan complexes in the outer membrane of *Rhizobium leguminosarum*. *J Bacteriol* 1989;171:3989–95.
- 27 Hill SA, Judd RC. Identification and characterization of peptidoglycan-associated proteins in *Neisseria gonorrhoeae*. *Infect Immun* 1989;57:3612–18.
- 28 Majcherzyk PA, Langen H, Heumann D, Fountoulakis M, Glauser MP, Moreillon P. Digestion of *Streptococcus pneumoniae* cell walls with its major peptidoglycan hydrolase releases branched stem peptides carrying proinflammatory activity. *J Biol Chem* 1999;274:12537–43.
- 29 Sato K, Saito H, Tomioka H, Yokokura T. Enhancement of host resistance against listeria infection by *Lactobacillus casei*: efficacy of cell wall preparation of *Lactobacillus casei*. *Microbiol Immunol* 1988;32:1189–200.
- 30 Gilbert J, Harrison J, Parks C, Fox A. Analysis of the amino acid and sugar composition of streptococcal cell walls by gas chromatography-mass spectrometry. *J Chromatogr* 1988;441:323–33.
- 31 Zhang Y, Gripenberg Lerche C, Soderstrom KO, Toivanen A, Toivanen P. Antibiotic prophylaxis and treatment of reactive arthritis. Lessons from an animal model. *Arthritis Rheum* 1996;39:1238–43.
- 32 van der Heijden IM, Wilbrink B, Tchetverikov I, Schrijver IA, Schouls LM, Hazenberg MP, et al. Presence of bacterial DNA and bacterial peptidoglycans in joints of patients with rheumatoid arthritis and other arthritides. *Arthritis Rheum* 2000;43:593–8.
- 33 Šimelyte E, Rimpiläinen M, Lehtonen L, Zhang X, Toivanen P. Bacterial cell wall-induced arthritis: chemical composition and tissue distribution of four *Lactobacillus* strains. *Infect Immun* 2000;68:3535–40.
- 34 Hoijer MA, Melief MJ, Calafat J, Roos D, van den Beemd RW, van Dongen JJ, et al. Expression and intracellular localization of the human N-acetylmuramyl-L-alanine amidase, a bacterial cell wall-degrading enzyme. *Blood* 1997;90:1246–54.
- 35 Striker R, Kline ME, Haak RA, Rest RF, Rosenthal RS. Degradation of gonococcal peptidoglycan by granule extract from human neutrophils: demonstration of N-acetylglucosaminidase activity that utilizes peptidoglycan substrates. *Infect Immun* 1987;55:2579–84.
- 36 DeJoy SQ, Ferguson Chanowitz KM, Sapp TM, Oronsky AL, Lapierre LA, Zabriskie JB, et al. M protein deficient streptococcal cell walls can induce acute and chronic arthritis rats. *Cell Immunol* 1990;125:526–34.
- 37 Lehman TJ, Allen JB, Plotz PH, Wilder RL. Bacterial cell wall composition, lysozyme resistance, and the induction of chronic arthritis in rats. *Rheumatol Int* 1985;5:163–7.