Unusual and memorable.....

Case Number 22: An interesting case of herpes zoster in rheumatoid arthritis

Series editor: Gary D Wright

An 82 year old woman with seronegative rheumatoid arthritis was admitted during a flare in her disease. Since diagnosis five years previously her arthritis had been managed with low dose oral corticosteroids at a dose of 10 mg daily.

Shortly after admission she complained of severe pain on the sole of the

left foot. Examination at the time showed a discrete area of vesicular eruption (fig 1). Oral antiviral treatment (acyclovir) was started for herpes zoster. Within 24 hours she had developed a more extensive zoster rash distributed over the posterior aspect of the left leg from ankle to buttock in the S1/S2

dermatome (fig 2). Symptomatically the neuralgic pain completely settled within two days while the rash persisted for a further few weeks.

Herpes zoster infection is well documented in patients with rheumatoid arthritis, particularly those receiving oral corticosteroids and methotrexate.¹⁻⁴ Although most cases are limited to the typical cutaneous zoster eruption, more complicated infections can occur such as disseminated cutaneous herpes zoster complicated by necrotising fasciitis,⁵ herpes zoster encephalomyelitis,⁷ and lower motor neurone paresis.⁸ Sciatic nerve involvement is unusual in herpes zoster reactivation.

E Campalani, G K Meenagh, M B Finch,
Department of Rheumatology, Royal Victoria
Hospital, Grosvenor Road Belfast BT12 6BA, UK

Correspondence to: Dr Meenagh;
garymeenagh@yahoo.co.uk

Figure 1 Vesicular eruption on the sole of the foot.

Figure 2 Herpes zoster rash.

REFERENCES

- 1 **Antonelli MA**, Moreland LW, Brick JE. Herpes zoster in patients with rheumatoid arthritis treated with weekly, low-dose methotrexate. *Am J Med* 1991;90:295-8.
- 2 **McCarty DJ**, Harman JG, Grassanovich JL, Qian C, Klein JP. Combination drug therapy of seropositive rheumatoid arthritis. *J Rheumatol* 1995;22:1636-45.
- 3 **Hernandez-Cruz B**, Cardiel MH, Villa AR, Alcocer-Varela J. Development, recurrence and severity of infections in Mexican patients with rheumatoid arthritis. A nested case-control study. *J Rheumatol* 1998;25:1990-7.
- 4 **Van der Veen MJ**, Van der Heide A, Kruize AA, Bijlsma JW. Infection rate and use of antibiotics in patients with rheumatoid arthritis treated with methotrexate. *Ann Rheum Dis* 1994;53:224-8.
- 5 **Ching DW**. Severe, disseminated, life threatening herpes zoster infection in a patient with rheumatoid arthritis treated with methotrexate. *Ann Rheum Dis* 1995;54:155.
- 6 **Jarrett P**, Ha T, Oliver F. Necrotizing fasciitis complicating disseminated cutaneous herpes zoster. *Clin Exp Dermatol* 1998;23:87-8.
- 7 **Lyon CC**, Thompson D. Herpes zoster encephalomyelitis associated with low dose methotrexate for rheumatoid arthritis. *J Rheumatol* 1997;24:589-91.
- 8 **Molloy MG**, Goodwill CJ. Herpes zoster and lower motor neurone paresis. *Rheumatol Rehabil* 1979;18:170-3.