PostScript

MATTERS ARISING

If you have a burning desire to respond to a paper published in the *Annals of the Rheumatic Diseases*, why not make use of our "rapid response" option?

Log on to our website (www.annrheumdis.com), find the paper that interests you, and send your response via email by clicking on the "eLetters" option in the box at the top right hand corner.

Providing it isn't libellous or obscene, it will be posted within seven days. You can retrieve it by clicking on "read eLetters" on our homepage.

The editors will decide as before whether also to publish it in a future paper issue.

Power Doppler sonography: synovial tissue assessment in RA

We read with interest the paper by Carotti *et al* that appeared in the October issue of the *Annals of the Rheumatic Diseases*.¹ The authors investigated the usefulness of contrast enhanced power Doppler ultrasonography (PDS) in determining the vascularisation rate of the synovial pannus of patients with rheumatoid arthritis (RA). After intravenous infusion of an echo contrast (Levovist, Schering, Germany) PDS was performed on the knee joints of 42 patients with RA, who had previously been classified as having active ($n=15$), moderately active ($n=14$), and inactive knee synovitis ($n=13$). The results were expressed as the area underlying time-intensity curves. Highest vascularisation scores were found in patients with active synovitis (216.2 (33.4), mean (SD)), whereas lower values were found in those with moderately active (186.8 (25.8)) and inactive synovitis (169.6 (20.6)). Significant difference ($p<0.01$) was found between the active and the inactive group only. Also, significant correlations were found between the vascularisation scores and some indices of the disease activity. The authors concluded that PDS may be a valuable method of distinguishing between inflammatory and non-inflammatory pannus of knee joints in RA and thus may have a clinical potential for both diagnostic and therapeutic purposes.

Unfortunately, Carotti *et al* did not provide any data on the sensitivity and specificity of the method used. Moreover, no figure showing distribution of the vascularisation scores in individual patients and/or confidence limits of the mean was available. This is a major problem and, in our opinion, the data presentation does not allow for any conclusions about how the PDS really performs in clinical settings. It should be emphasised that statistical significance alone is not sufficient as it could easily be achieved if there were some lower and some higher vascularisation scores in patients with inactive and active synovitis, respectively.

We are aware that the study was designed as a preliminary one and a relatively small

group of patients was included. Nevertheless, we believe that all *Annals* readers would benefit much from seeing the sensitivity and specificity figures/estimates or the presentation of the individual vascularisation scores. We wonder if these could be provided by the authors.

The statement that ultrasonography offers a simple, non-invasive, reproducible, non-radiating, and inexpensive imaging technique was made several times in the paper. It holds true for ultrasonography as a general imaging technique. However, it seems somewhat misleading in the context of the methodology used by the authors. PDS is not a simple technique as it requires a team of well trained and experienced radiologists and technicians who are aware of possible caveats of the method. The procedure requires venepuncture and an intravenous contrast infusion, and thus can at most be regarded as a low invasive one. It should be emphasised that Levovist, although generally well tolerated, may rarely lead to some adverse reactions (for example, local tissue irritations, dyspnoea, changes in pulse/blood pressure, nausea, vomiting, headaches, dizziness, skin reactions—data taken from the compound data sheet). To our knowledge, no data exist on the reproducibility of PDS for the knee pannus examination. A high-end equipment has to be used to achieve good results which is relatively expensive. It seems that from the long list of adjectives used by the authors, "non-radiating" is the only valid one for the PDS method described.

P Hrycaj, J K Lacki

Department of Rheumatology and Clinical Immunology, University School of Medical Sciences, Winogrady 144, 61-626 Poznan, Poland

Correspondence to: Dr P Hrycaj; phrycaj@icpnet.pl

Reference

- 1 Carotti M, Salaffi F, Manganelli P, Salera D, Simonetti B, Grassi W. Power Doppler sonography in the assessment of synovial tissue of the knee joint in rheumatoid arthritis: a preliminary experience. *Ann Rheum Dis* 2002;61:877-82.

Authors' reply

We thank Drs Hrycaj and Lacki very much for their comments because they give us the opportunity to illustrate further details of our study.¹ Recent papers have shown a very close relationship between the results of power Doppler sonography (PDS) and those of dynamic gadolinium (Gd) enhanced magnetic resonance imaging (MRI; a sensitivity of 88.8% and a specificity of 97.9%),² and between PDS findings and histopathological findings³ of synovial membrane vascularity. It has also been shown that contrast enhanced PDS significantly improves the detection of intra-articular vascularity, compared with the unenhanced technique.⁴ Contrast enhanced PDS demonstrated significant difference in intra-articular colour flow signals between joints of patients with inactive RA and those with active RA, between joints of patients with inactive RA and those with moderately active RA, and between joints of patients with moderately active RA and those with active RA.⁵

With regard to the sensitivity and specificity of the PDS technique, as far as we know no data have been reported. Our study has all the features of a preliminary investigation and, obviously, cannot answer all the questions on the topic. However, the receiver operating characteristic (ROC) curve was employed to describe how well various values of the area underlying the enhancement curves of the PDS can detect patients with RA with high and low synovitis activity (fig 1).⁶ This analysis involves plotting the true positive rate (sensitivity) against the false positive rate ($100 - \text{specificity}$) for possible cut off scores. Sensitivity and specificity (which constitute the discriminating ability of a test) were used in agreement with the current definitions of these terms.⁷ Moreover, we calculated 95% confidence intervals (CI) of sensitivity and specificity. Each point on the ROC plot represents a sensitivity/specificity pair corresponding to a particular decision threshold.^{8,9} The area under the ROC curve is used to evaluate the screening scale's performance. For PDS, the mean (SEM) area under the curve is 0.859 (0.072); 95% CI 0.675 to 0.960. According to Swets *et al*, areas from 0.50 to about 0.70 represent poor accuracy, those from 0.70 to 0.90 are "useful for some purposes", and higher values represent high accuracy.⁷ From the ROC curve in fig 1 we computed the optimal cut off point, corresponding to the maximum sum of sensitivity and specificity. For our method an optimal point of 191 comes close to maximising both sensitivity and specificity. With this cut off point for the area under the time-intensity curve, the sensitivity is 80.0% and specificity is 84.6%. These findings have implications for both clinical practice and research purposes in assessment of disease activity in patients with RA.

Figure 2 shows the distribution of PDS scores in individual patients with active and inactive synovitis and the confidence limits of the mean. The mean (SD) values of the areas under the curves showed a significant difference ($p<0.01$) between the patients with active synovitis (216.2 (33.4); 95% CI 193.1 to

Figure 1 The ROC curve illustrating the relation between sensitivity and complement of specificity ($100 - \text{specificity}$) in rheumatoid patients for the PDS, with the index of synovitis activity as the external criterion (inactive synovitis v active synovitis). The triangle on the curve shows the optimal cut off point, corresponding to the highest sensitivity/specificity combination.

Figure 2 Box plot of the values of area under the time-intensity curve from patients with inactive and those with active synovitis. Boxes encompass the 25th, median (50th), and 75th centile values; vertical bars represent the 10th and 90th centile values. Statistical comparison was performed with the Mann-Whitney U-test.

239.4) and those with inactive synovitis (169.6 (20.6); 95% CI 150.8 to 188.4) (fig 2).

The characteristics of PDS are no more complex than those of many other clinical or instrumental procedures commonly used in daily clinical practice. It obviously requires adequate training and high quality equipment. In our study all the investigations were performed by one of the authors (MC), who is specialist both in rheumatology and radiology. As with all drugs, echo contrast agents may have side effects, although most of the side effects have been reported to be mild and not prolonged. In our patients no clinical side effects from the agents were detected. We excluded from the study patients with galactosaemia, with suspected pregnancy and critical illness and patients who had taken any contrast agent within 24 hours.

We agree with Drs Hrycaj and Lacki that the use of echo contrast agents is, at this time, relatively expensive, and they are not yet routinely available, but it may be a cost effective alternative to dynamic MRI. Moreover, the implementation of microinjection devices that are currently being studied may decrease the required dose of contrast medium, and this should reduce overall examination time and the cost of ultrasound contrast agent for each patient. Consequently, the cost of contrast enhanced PDS should be comparable with that of other techniques. The assessment of reproducibility is a problem of all diagnostic procedures based on the use of contrast agents for obvious ethical reasons. This and other complex aspects of a new methodological approach to the study of rheumatoid synovitis cannot be extensively examined in a single preliminary report.

Although PDS with contrast agents has some well known limitations, it allows access to otherwise undetectable information that may improve our basic knowledge of rheumatoid synovitis and, in the near future, might also be useful in the assessment of disease activity in RA.⁵

M Carotti, B Simonetti, D Salera

Department of Radiology, University of Ancona, Italy

F Salaffi, W Grassi

Department of Rheumatology, University of Ancona, Italy

P Manganelli

Rheumatic Disease Unit, Hospital of Parma, Italy

Correspondence to Dr F Salaffi, Department of Rheumatology, University of Ancona, Ospedale A Murri, via dei Colli 52, 60035, Jesi (Ancona), Italy; fsalaffi@tin.it

References

- 1 **Carotti M**, Salaffi F, Manganelli P, Salera D, Simonetti B, Grassi W. Power Doppler sonography in the assessment of synovial tissue of the knee joint in rheumatoid arthritis: a preliminary experience. *Ann Rheum Dis* 2002;61:877-82.
- 2 **Szkudlarek M**, Court-Payen M, Strandberg C, Klarlund M, Klausen J, Østergaard M. Power Doppler ultrasonography for assessment of synovitis in the metacarpophalangeal joints of patients with rheumatoid arthritis. A comparison with dynamic magnetic resonance imaging. *Arthritis Rheum* 2001;44:2018-23.
- 3 **Walther M**, Harms H, Krenn V, Stephan R, Faehndrich T-P, Göhlke F. Correlation of power Doppler sonography with vascularity of the synovial tissue of the knee joint in patients with osteoarthritis and rheumatoid arthritis. *Arthritis Rheum* 2001;44:331-8.
- 4 **Magarelli N**, Guglielmi G, Di Matteo L, Tartaro A, Mattei PA, Bonomo L. Diagnostic utility of an echo-contrast agent in patients with synovitis using power doppler ultrasound: a preliminary study with comparison to contrast-enhanced MRI. *Eur Radiol* 2001;11:1039-46.
- 5 **Klauser A**, Frauscher F, Schirmer M, Halpern E, Pallwein L, Herold M, et al. The value of contrast-enhanced color Doppler ultrasound in the detection of vascularization of finger joints in patients with rheumatoid arthritis. *Arthritis Rheum* 2002;46:647-53.
- 6 **Swets JA**. Measuring the accuracy of diagnostic systems. *Science* 1988;240:1285-93.
- 7 **Swets JA**. ROC analysis applied to the evaluation of medical imaging techniques. *Invest Radiol* 1979;14:109-21.
- 8 **Zweig MH**, Campbell G. Receiver-operating characteristic (ROC) plots: a fundamental evaluation tool in clinical medicine. *Clin Chem* 1993;39:561-77.
- 9 **Hauley JA**, McNeil BJ. The meaning and use of the area under a receiver operating characteristic (ROC) curve. *Radiology* 1982;143:29-36.

A case of orbital myositis associated with rheumatoid arthritis

In the October 2002 issue of the *Annals* Nabili and colleagues described a case of orbital myositis (OM) associated with rheumatoid arthritis and suggested that the major differential diagnosis was thyroid ophthalmopathy.¹

The mechanical component of orbital involvement in Graves' disease (GD) can produce diplopia (5-10%), proptosis, chemosis, scleral injections, and retrobulbar pressure or pain, whereas eyelid lag or retraction and stare are caused by the spastic component (in some cases lid retraction can be obscured by periorbital oedema). The extraocular muscles most commonly affected are the medial and inferior rectus muscles.

The authors suggested characteristics which would distinguish the two conditions, and among these they included the systemic manifestations of GD.

GD should be seen as a systemic disease with various combinations of hyperthyroid goitre, ophthalmopathy, and dermopathy. Current pathogenetic theory shows that ophthalmopathy itself is a direct, independent manifestation of autoimmunity, caused by autoreactivity to the thyrotropin receptor expressed by the preadipocyte subpopulation

of orbital fibroblasts, mediated by type 2 helper T cells and thyroid stimulating antibodies. Moreover, it is well established that ophthalmopathy may precede hyperthyroidism or occur after its resolution (particularly in smokers after treatment with radioactive iodine). GD can be found in association with other autoimmune disorders such as type 1 diabetes mellitus, Addison's disease, vitiligo, pernicious anaemia, alopecia areata, myasthenia gravis, coeliac disease, systemic lupus erythematosus, Sjögren's syndrome, and other HLA-DR3 associated diseases. The association with rheumatoid arthritis is rarer, but recognised, even in multiple autoimmune syndromes. So where there is autoimmunity with orbital involvement the absence of clinical hyperthyroidism should not rule out the diagnosis of GD.

Finally, the diagnosis of thyroid ophthalmopathy was unlikely because only one eye was affected. As suggested by the authors, magnetic resonance imaging is indicated to establish the correct diagnosis. This would show oedematous extraocular muscles with enlargement of the tendons in OM, and normal tendons in GD. Early response to systemic glucocorticoids confirms the diagnosis of OM.

Another disease that should be considered in the differential diagnosis is arteriovenous malformation, such as indirect carotid-cavernous fistula, which can develop spontaneously in older women. In this case the presence of a diffuse head murmur would be an important observation.²⁻⁵

G P Fadini

University of Padua, Italy

Correspondence to: Dr G P Fadini, v. Turazza 48/A, 35100 Padova, Italy; Gianpaolofadini@hotmail.com

References

- 1 **Nabili S**, McCarey DW, Browne B, Capell HA. A case of orbital myositis associated with rheumatoid arthritis. *Ann Rheum Dis* 2002;61:938-9.
- 2 **Wartofsky L**. Graves' ophthalmopathy: current concepts regarding pathogenesis and management. *Endocr Rev* 1993;14:747.
- 3 **Bahn RS**, Dutton CM, Natt N, Joba W, Spitzweg C, Heufelder AE. Thyrotropin receptor expression in Graves' orbital adipose/connective tissue: potential autoantigen in Graves' ophthalmopathy. *J Clin Endocrinol Metab* 1998;83:998-1002.
- 4 **Mourits MP**, Prummel MF, Wiersinga WM, Koornneef L. Clinical activity score as a guide in the management of patients with Graves' ophthalmopathy. *Clin Endocrinol (Oxf)* 1997;47:9-14.
- 5 **Weetman PA**. Graves' disease. *N Engl J Med* 2000;343:1236-48.

Authors' reply

We thank Dr Fadini for his interest in our case report of orbital myositis in association with rheumatoid arthritis.¹

He rightly discusses in some detail the possibility of thyroid ophthalmopathy occurring in the absence of a systemic hyperthyroid diathesis. As we made clear in our article, there is a considerable variety of presenting clinical features in this condition. We sought to present these as concisely as possible, with particular reference to the most common presentation.

The differential diagnosis of an arteriovenous malformation was not likely in the clinical setting outlined. We agree that, ultimately, the diagnosis was made on the

basis of typical magnetic resonance imaging findings and the prompt clinical response to corticosteroid treatment.

S Nabili, D W McCarey, B Browne, H A Capell

Glasgow Royal Infirmary, Centre for Rheumatic Diseases, 84 Castle Street, Glasgow G4 0SF, UK

Correspondence to: Dr DW McCarey; gcl376@clinmed.gla.ac.uk

Reference

- 1 Nabili S, McCarey DW, Browne B, Capell HA. A case of orbital myositis associated with rheumatoid arthritis. *Ann Rheum Dis* 2002;61:938-9.

NOTICES

MSc Programme in Clinical Rheumatology

Applications are invited for places on this MSc programme, starting September 2003, which provides an excellent academic basis for those aiming at a career in rheumatology or a related subject. Applicants should be medically qualified and should have had at least two years of general medical experience after qualification. Previous experience in rheumatology is desirable, but not essential. The programme is undertaken part time over two years and is now well established, entering its ninth year.

Topics covered will include: basic science, clinical skills, peripheral joint problems, spinal problems, connective tissue disease and vasculitis, and the epidemiology of musculoskeletal diseases. A supervised project, which may be either clinical or laboratory based, is an integral part of the programme. The closing date for applications is 30 May 2003.

Further details can be obtained from Miss Lisa McClair, ARC Epidemiology Unit, Stopford Building, University of Manchester, Oxford Road, Manchester, M13 9PT, UK. Tel: (0) 161 275 5993. Fax: (0) 161 275 5043. Email: Lisa.Mcclair@man.ac.uk

Ten topics in rheumatology

A two day postgraduate course in advanced rheumatology will be held at St Thomas' Hospital, London on 5 and 6 June 2003. The registration fee is £120, and 60 subsidised registrations are offered at £20 to current SpRs to the first 60 applications received. CME accreditation

Topics include:

- What's new in lupus?
- What's new in osteoporosis?
- What's new in scleroderma?
- What's new in RA?
- Hughes syndrome: 20 years on

- Pregnancy and infertility
- Pulmonary hypertension
- Accelerated atheroma
- "Hot topics"
- Nightmare on Lambeth Palace Road

Further details: Dr Graham R V Hughes, Lupus Research Unit, The Rayne Institute, St Thomas' Hospital, London, SE1 7EH. Tel: 020 7928 9292 ext. 2888/3357. Fax: 020 7633 9422. Email: sandy.hampson@kcl.ac.uk

FORTHCOMING EVENTS

30th Annual Meeting of the International Society for the Study of the Lumbar Spine

13-17 May; Vancouver, Canada

Contact: Dr S Boden, Sunnybrook and Women's Health Science Center, Room MG 323, 2075 Bayview Avenue, Toronto, Canada M4N 3M5

Tel: 416 480 4833

Fax: 416 480 6055

Email: shirley.fitzgerald@swchsc.on.ca

4th Annual European Congress of Rheumatology

18-21 June 2003; Lisbon, Portugal

Contact: Fred Wyss, Executive Secretary EULAR, Witikonstrasse 15, CH-8032, Zurich, Switzerland

Tel: +41 1 383 9690

Fax: +41 1 383 9810

Email: eular@bluewin.ch

Website: www.eular.org

25th Annual Meeting of the American Society for Bone and Mineral Research (ASBMR)

19-23 September 2003; Minneapolis, Minnesota, USA

Tel: +1 202 367 1161

Fax: +1 202 367 2161

Email: asbmr@dc.sba.com

Website: www.asbmr.org

7th EULAR Sonography Course

9-12 October, 2003; Rome, Italy

An introductory and practical course on musculoskeletal ultrasonography

Scientific secretariat: Professor Guido Valesini

Email: annamaria.iagrocco@uniroma1.it

Contact: Organising secretariat: Michela Civelli, EDRA Spa, Medical Publishing and News Media, Viale Monza, 133 - 20125, Milan, Italy

Tel: +39 (0)2 281 72300

Fax: +39 (0)2 281 72399

Email: edracongressi@dsmedigroup.com

OARSI World Congress on Osteoarthritis

12-15 October 2003; Berlin, Germany

Tel: +1 856 439 1385

Fax: +1 856 439 0525

Email: oarsi@oarsi.org

Website: www.oarsi.org

Fourth International Symposium on Clinical and Economic Aspects of Osteoporosis and Osteoarthritis

14-17 November 2003; Nice, France

Contact: Organisation Secretariat, YP Communication, 108 boulevard G Kleyer, 4000 Liège, Belgium

Tel: +32 (4) 254 12 25

Fax: +32 (4) 254 12 90

Email: yoland@piettecommunication.com

Website: http://nice.piettecommunication.com

XIth International Conference on Behçet's Disease

24-27 October 2004; Antalya, Turkey

Contact: Congress Secretariat, Figur Congress and Organization Services Ltd. STI, Ayazmad-eresi Cad. Karadut Sok. No: 7 80888 Dikilitas, Istanbul

Tel: +90 (212) 258 60 20

Fax: +90 (212) 258 60 78

Email: behcet2004@figur.net

Website: www.behcet2004.org

Future EULAR congresses

9-12 June 2004; EULAR 2004; Berlin, Germany

8-11 June 2005; EULAR 2005; Vienna, Austria

21-24 June 2006; EULAR 2006; Amsterdam, The Netherlands

Future ACR meetings

24-28 October 2003; 67th Annual Scientific Meeting; Orlando, Florida

16-21 October 2004; 68th Annual Scientific Meeting; San Antonio, Texas