The melatonin-cytokine connection in rheumatoid arthritis

M Cutolo, G J M Maestroni

Melatonin up regulates cytokine production and immune function

A substance from pineal gland extracts, which lightens the skin melanocytes of amphibians and fishes, was isolated in 1958 and called melatonin (MLT, *N*-acetyl-5-methoxytryptamine).¹ In both diurnal and nocturnal species, the absence of light at night stimulates MLT biosynthesis. Electrical signals originating from the retina reach the suprachiasmatic nuclei which, in turn, send inputs via the paraventricular nuclei to the spinal cord and then to the superior cervical ganglia. The fibres terminate at the pinealocytes.² Absence of light results in increased norepinephrine release and activation of $\alpha 1$ and β -adrenergic receptors on the pinealocytes. This triggers a series of intracellular responses, resulting in activation of the enzymes *N*-acetyltransferase (EC 2.3.1.87) and hydroxyindole-*O*-methyl transferase (EC 2.1.1.4), which convert serotonin into MLT.² The circadian nocturnal release of MLT has a profound influence on the internal environment of the organism, with diverse physiological effects. The main function of MLT seems to be that of synchronising the organism in the photoperiod and it may have a role in reproduction, metabolism, seasonality, thermoregulation, and immunity.

MELATONIN AND THE HUMAN IMMUNE SYSTEM

Animal studies have shown that binding of MLT to specific receptors in antigen activated T helper (Th) cells results in an up regulation of cytokine production and immune function.³ In general, the immunoenhancing action of MLT seems restricted to T dependent antigens and to be most pronounced in immunodepressed situations. For example, MLT may completely counteract thymus involution and the immunological depression induced by stress or glucocorticoid treatment,⁴ or restore depressed immunological functions after soft tissue trauma and haemorrhagic shock.⁵ MLT may also rescue haematopoiesis in mice treated with cancer chemotherapeutic compounds.⁶

This effect apparently involves the endogenous release of granulocyte/macrophage colony stimulating factor and MLT-induced opioid cytokines.^{3 6 7}

The immunoenhancing action of MLT has been confirmed and extended in a variety of animal species and in humans and in birds.⁸⁻¹² In human peripheral blood mononuclear cells, at physiological concentrations MLT has been reported to stimulate the production of interleukin 1 (IL1), IL2, interferon γ (IFN γ), IL6, and IL12 but not IL4.¹³⁻¹⁵ Physiologically, the nocturnal MLT peak has been associated with a high IFN γ /IL10 ratio—that is, the MLT rhythm positively correlates with the rhythmicity of the Th1/Th2 cell ratio.¹⁶ In patients with ischaemic stroke an impaired nocturnal MLT excretion has been associated with impaired cell mediated immunity and changes of lymphocyte subsets.¹⁷ Most interestingly, reduction of MLT secretion has been reported to parallel disease progression and to correlate with serum IL12 levels in HIV-1 infected patients.¹⁸ These human studies confirmed that MLT possesses important immuno-enhancing properties and suggest that MLT may favour a Th1 cell response.

“MLT possesses important immuno-enhancing properties”

Most recently, the connection between IL2 and MLT in human lymphocytes has been strengthened by the observation that MLT may be synthesised and released in large quantity by human lymphocytes.^{19 20} The lymphocyte MLT production seems to be strictly linked to the release of IL2 as inhibition of MLT synthesis resulted in a decrease of IL2 production and addition of exogenous MLT resulted in an increase of IL2 production. These findings indicate that in addition to pineal gland, human lymphoid cells are an important physiological source of melatonin and that this melatonin might play a part in the regulation of the human immune system, possibly by acting as an intracrine, autocrine, and/or paracrine

substance.^{13 14 19 20} The MLT/IL2 connection seems particularly relevant because IL2 has a central role in immune homeostasis. Recent studies indicate that a failure in the production of CD4+CD25+ regulatory T cells is the underlying cause of autoimmunity in the absence of IL2.²¹ Yet, IL2 has been used clinically to enhance T cell immunity in patients with AIDS or cancer, and blocking antibodies to the IL2 receptor are used to inhibit T cell responses against transplanted tissues.²² In analogy to IL2, MLT has also been used to enhance T cell immunity in patients with cancer.²³ Perhaps, the ability of MLT to enhance production of inflammatory cytokines from human monocytes/macrophages, including IL12,²⁴⁻²⁶ turns the MLT/IL2 connection towards the enhancement of T cell immunity rather than to formation of T regulatory cells.

MLT, IL2, AND GLUCOCORTICOID RESISTANCE

Endogenous glucocorticoids, the end-effectors of the hypothalamus-pituitary-adrenal (HPA) axis, also have an important role in regulating homeostasis of the immune system.²⁷ It is generally recognised that glucocorticoids suppress peripheral Th cell responses.²⁸ However, the effect of glucocorticoids on T cell functions may depend on the activation and differentiation state of T cells, as well as on the conditions or agents used for stimulation.²⁸ Thus, resting T cells appear to differ from activated T cells in their response to glucocorticoids, while infections, cytokines, and proinflammatory factors can alter the sensitivity of T cells.²⁹

In clinical practice, reduced T cell sensitivity to the suppressive action of glucocorticoids is a common problem that complicates the management of chronic inflammatory and autoimmune diseases. Patients with glucocorticoid resistance usually have severe inflammation, associated with significantly high levels of cytokines, growth factors, and costimulatory factors in the inflamed tissues.³⁰⁻³² In particular, it has been reported that IL2 may induce glucocorticoid resistance in Th cells.³³ As mentioned above, MLT may counteract the immunosuppressive effect of endogenous or exogenous glucocorticoids.³⁴ The mechanism of this effect was related to stimulation of opioid cytokines in T lymphocytes and/or to a direct inhibition of glucocorticoid receptor expression.^{35 36} In addition to these effects, the MLT/IL2 connection suggests that excessive MLT production might be involved in glucocorticoid resistance. This hypothesis should be tested.

MLT, CYTOKINES, CIRCADIAN RHYTHMS, AND RA

In 2002 our first study evaluated MLT levels in patients who had rheumatoid arthritis (RA), with a focus upon analyses of circadian variations.³⁷ MLT serum levels at 8 pm and 8 am were found to be significantly higher in patients who had RA than in healthy controls ($p < 0.05$). The differences were more evident in patients who were older than 60 years.

"MLT serum levels are significantly higher in patients with RA than in controls"

In patients who had RA and in healthy subjects, MLT levels increased progressively from 8 pm to the early morning hours; however, they reached peak levels at midnight in patients who had RA, which was at least 2 hours earlier than in controls. Subsequently, MLT concentrations in patients who had RA reached a plateau that lasted for 2–3 hours; this was not observed in controls. After 2 am, MLT levels decreased similarly in patients who had RA and in healthy subjects. The study confirmed that the nocturnal rhythm of MLT occurs also in patients who have RA, but with an earlier peak and a longer duration in the early morning.³⁷ IFN γ , IL1, IL6, tumour necrosis factor α (TNF α), IL2, and IL12 production (Th1 cytokines/promoting cytokines) reach their peak during the night and early morning, at the same time that MLT serum levels are highest and plasma cortisol is lowest.

Accordingly, among the signs of joint inflammation in patients who have RA, the intensity of pain varies as a function of the hours of the day; pain is greater after awakening in the morning than in the afternoon or evening.³⁸ Circadian changes also are observed in joint swelling and finger size in the early morning in patients affected by RA.³⁸ Therefore, MLT may activate the inflammatory response during the night, at least in RA, which is considered to be a Th1 cytokine driven immune disease.³⁹

We found that MLT was detectable in high concentration in synovial fluids from patients who had RA, and binding sites for MLT were present in synovial macrophages.^{40–41} In addition, cultured RA synovial macrophages respond to MLT stimulation with an increased proinflammatory cytokine production.²⁴

Interestingly, recent studies and reviews examined the epidemiological evidence which suggests that ultraviolet radiation may have a protective role in RA. A gradient of increasing incidence of RA with latitude and seasonal variation has also been reported^{42–44}

Therefore, we evaluated serum MLT, cortisol, TNF α , and IL6 circadian rhythm in patients who had RA from a northern European country (Estonia).

Furthermore, we compared the MLT and cortisol levels in that group with patients from a southern European country (Italy), to detect a possible influence of different daily winter photoperiods.^{45–46} Blood samples were obtained during the months of January and February. The study showed that, a significantly higher MLT concentration ($p < 0.01$) and an earlier peak was seen in Estonian patients who had RA than in controls matched for age and sex. In addition, MLT serum concentrations were significantly higher in Estonian patients who had RA than in the Italian patients, at midnight, and the difference was even greater over the study duration. No significant difference was seen in serum cortisol levels between Estonian patients who had RA and their healthy controls.^{45–46}

Significantly higher serum IL6 and TNF α concentrations were also observed at 4 pm and midnight in Estonian patients who had RA than in the Italian patients who had RA.^{45–46}

This important study shows, for the first time, that in a northern European country, the circadian serum concentrations of MLT and TNF α are significantly higher than in matched patients who had RA from a southern European country. In addition, MLT and TNF α concentrations were found to be increased in patients who had RA.

"Circadian serum concentrations of MLT and TNF α are significantly higher in patients with RA from northern Europe than in those from further south"

The reduced daily light exposure in northern Europe, at least during the winter, might explain the higher and more prolonged MLT serum concentrations that were found in northern patients who had RA.

Interestingly, several studies in patients with RA showed that overall activity of the HPA axis, particularly the glucocorticoid synthesis, remained inappropriately normal (or relatively low) and apparently was found to be insufficient to inhibit continuing inflammation.⁴⁷ Therefore, an imbalance between the anti-inflammatory effects of cortisol and the proinflammatory effects of MLT seems evident in patients affected by RA and supports the increased Th1 cytokine production during the night.⁴⁸

Finally, the increased prevalence of autoimmune diseases, such as RA, which is seen in northern Europe also may be related to the increased

immunostimulatory effects that are exerted during the night by MLT and to a reduced neuroendocrine modulation during the light phase of the photoperiod (cortisol). The prevalence of RA is, in fact, much higher in northern Europe than in the Mediterranean countries, with rates of 1.96% in Finland, 1.1% in England, 0.9% in Sweden, Denmark, and the Netherlands compared with 0.2% in Greece and 0.3% in Italy and Israel.^{49–50}

Ann Rheum Dis 2005;**64**:1109–1111.

doi: 10.1136/ard.2005.038588

Authors' affiliations

M Cutolo, Research Laboratory and Division of Rheumatology, Department of Internal Medicine, University of Genova, Viale Benedetto XV, 6, 16132 Genova, Italy
G J M Maestroni, Centre for Experimental Pathology, Cantonal Institute of Pathology, Via In Selva 24, PO Box, 6601 Locarno, Switzerland

Correspondence to: Professor M Cutolo, mcutolo@unige.it

REFERENCES

- 1 Lerner AB, Case JD, Takahashi Y, Lee TH, Mori W. Isolation of melatonin, the pineal gland factor that lightens melanocytes. *J Am Chem Soc* 1958;**80**:2587–92.
- 2 Yu H-S, Tsien ATC, Reiter RJ. Melatonin: history, biosynthesis, and assay methodology. In: Yu H-S, Reiter RJ, eds. *Melatonin. Biosynthesis, physiological effects, and clinical applications*. Boca Raton: CRC Press, 1993:1–17.
- 3 Maestroni GJM. Therapeutic potential of melatonin in immunodeficiency states, viral diseases, and cancer. *Adv Exp Med Biol* 1999;**467**:217–26.
- 4 Maestroni GJM, Conti A. The pineal neurohormone melatonin stimulates activated CD4+, Thy-1+ cells to release opioid agonist(s) with immuno-enhancing and anti-stress properties. *J Neuroimmunol* 1990;**28**:167–76.
- 5 Wichmann MW, Zellweger R, DeMaso A, Ayala A, Chaudry IH. Melatonin administration attenuates depressed immune functions trauma-hemorrhage. *J Surg Res* 1996;**63**:256–62.
- 6 Maestroni GJM, Covacci V, Conti A. Hematopoietic rescue via T-cell-dependent, endogenous GM-CSF by the pineal neurohormone melatonin in tumor bearing mice. *Cancer Res* 1994;**54**:2429–32.
- 7 Maestroni GJM, Conti A, Lissoni P. Colony-stimulating activity and hematopoietic rescue from cancer chemotherapy compounds are induced by melatonin via endogenous interleukin 4. *Cancer Res* 1994;**54**:4740–3.
- 8 Nelson RJ, Demas GE. Role of melatonin in mediating seasonal energetic and immunologic adaptations. *Brain Res Bull* 1997;**44**:423–30.
- 9 Rafii-El-Idrissi M, Calvo JR, Harmouch A, Garcia-Maurino S, Guerrero JM. Specific binding of melatonin by purified cell nuclei from spleen and thymus of the rat. *J Neuroimmunol* 1998;**86**:190–7.
- 10 Yellon SM, Teasley LA, Fagoaga OR, Nguyen HC, Truong HN, Nehlsen-Cannarella L. Role of photoperiod and the pineal gland in T cell-dependent humoral immune reactivity in the Siberian hamster. *J Pineal Res* 1999;**27**:243–8.
- 11 Kliger CA, Gehad AE, Hulet RM, Roush WB, Lillehoj HS, Mashaly MM. Effects of photoperiod and melatonin on lymphocyte activities in male broiler chickens. *Poult Sci* 2000;**79**:18–25.
- 12 Moore CB, Siopes TD. Effects of lighting conditions and melatonin supplementation of the

- cellular and humoral immune responses in Japanese quail *Coturnix coturnix japonica*. *Gen Comp Endocrinol* 2000;**119**:95–104.
- 13 **Garcia-Maurino S**, Gonzalez-Haba MG, Calvo JR, Rafii-El-Idrissi M, Sanchez-Margalet V, Goberna R, *et al*. Melatonin enhances IL-2, IL-6, and IFN- γ production by human circulating CD4+ cells. *J Immunol* 1997;**159**:574–81.
 - 14 **Morrey MK**, McLachlan JA, Serkin CD, Bakouche O. Activation of human monocytes by the pineal neurohormone melatonin. *J Immunol* 1994;**153**:2671–80.
 - 15 **Garcia-Maurino S**, Pozo D, Carrillo-Vico A, Calvo JR, Guerrero JM. Melatonin activates Th1 lymphocytes by increasing IL-12 production. *Life Sci* 1999;**65**:2143–50.
 - 16 **Petrovsky N**, Harrison L. The chronobiology of human cytokine production. *Int Rev Immunol* 1998;**16**:635–49.
 - 17 **Fiorina P**, Lattuada G, Silvestrini C, Ponari O, Dall'Aglio P. Disruption of nocturnal melatonin rhythm and immunological involvement in ischaemic stroke patients. *Scand J Immunol* 1999;**50**:228–31.
 - 18 **Nunnari G**, Nigro L, Palermo F, Leto D, Pomerantz RJ, Cacopardo B. Reduction of serum melatonin levels in HIV-1-infected individuals' parallel disease progression: correlation with serum interleukin-12 levels. *Infection* 2003;**31**:379–82.
 - 19 **Carrillo-Vico A**, Calvo JR, Abreu P, Lardone PJ, Garcia-Maurino S, Reiter RJ, *et al*. Evidence of melatonin synthesis by human lymphocytes and its physiological significance: possible role as intracrine, autocrine, and/or paracrine substance. *FASEB J* 2004;**18**:537–9.
 - 20 **Carrillo-Vico A**, Lardone PJ, Fernandez-Santos JM, Martin-Lacave I, Calvo JR, Karasek M, *et al*. Human lymphocyte-synthesized melatonin is involved in the regulation of the IL-2/IL-2 receptor system. *J Clin Endocrinol Metab* 2005;**90**:992–1000.
 - 21 **Malek TR**, Bayer AL. Tolerance, not immunity, crucially depends on IL-2. *Nat Rev Immunol* 2004;**4**:665–74.
 - 22 **Nelson BH**. IL-2, regulatory T cells, and tolerance. *J Immunol* 2004;**172**:3983–8.
 - 23 **Maestroni GJM**. The immunotherapeutic potential of melatonin. *Exp Opin Invest Drugs* 2001;**10**:467–76.
 - 24 **Cutolo M**, Villaggio B, Candido F, Valenti S, Giusti M, Felli L, *et al*. Melatonin influences interleukin-12 and nitric oxide production by primary cultures of rheumatoid synovial macrophages and THP-1 cells. *Ann N Y Acad Sci* 1999;**876**:246–54.
 - 25 **Garcia-Maurino S**, Pozo D, Carrillo-Vico A, Calvo JR, Guerrero JM. Melatonin activates Th1 lymphocytes by increasing IL-12 production. *Life Sci* 1999;**65**:2143–50.
 - 26 **Lissoni P**. Modulation of anticancer cytokines IL-2 and IL-12 by melatonin and the other pineal indoles 5-methoxytryptamine and 5-methoxytryptophol in the treatment of human neoplasms. *Ann N Y Acad Sci* 2000;**917**:560–7.
 - 27 **Webster JJ**, Tonelli L, Sternberg EM. Neuroendocrine regulation of immunity. *Annu Rev Immunol* 2002;**20**:125–63.
 - 28 **Ashwell JD**, Lu FW, Vacchio MS. Glucocorticoids in T cell development and function. *Annu Rev Immunol* 2000;**18**:309–45.
 - 29 **Kino T**, Chrousos GP. Tissue-specific glucocorticoid resistance-hypersensitivity syndromes: multifactorial states of clinical importance. *J Allergy Clin Immunol* 2002;**109**:609–13.
 - 30 **Corrigan CJ**, Brown PH, Barnes NC, Tsai JJ, Frew AJ, Kay AB. Glucocorticoid resistance in chronic asthma. Peripheral blood T lymphocyte activation and comparison of the T lymphocyte inhibitory effects of glucocorticoids and cyclosporin A. *Am Rev Respir Dis* 1991;**144**:1026–32.
 - 31 **Sher ER**, Leung DY, Surs W, Kam JC, Zieg G, Kamada AK, *et al*. Steroid-resistant asthma. Cellular mechanisms contributing to inadequate response to glucocorticoid therapy. *J Clin Invest* 1994;**93**:33–9.
 - 32 **Leung DY**, de Castro M, Szeffler SJ, Chrousos GP. Mechanisms of glucocorticoid-resistant asthma. *Ann N Y Acad Sci* 1998;**840**:735–46.
 - 33 **Walker KB**, Potter JM, House AK. Interleukin 2 synthesis in the presence of steroids: a model of steroid resistance. *Clin Exp Immunol* 1987;**68**:162–7.
 - 34 **Maestroni GJM**, Conti A, Pierpaoli W. Role of the pineal gland in immunity. Circadian synthesis and release of melatonin modulates the antibody response and antagonize the immunosuppressive effect of corticosterone. *J Neuroimmunol* 1986;**13**:19–30.
 - 35 **Maestroni GJM**, Conti A, Pierpaoli W. Role of the pineal gland in immunity. III. Melatonin antagonizes the immunosuppressive effect of acute stress via an opiate mechanism. *Immunology* 1988;**63**:465–9.
 - 36 **Sainz RM**, Mayo JC, Reiter RJ, Antolin I, Esteban MM, Rodriguez C. Melatonin regulates glucocorticoid receptor: an answer to its antiapoptotic action in thymus. *FASEB J* 1999;**13**:1547–56.
 - 37 **Sulli A**, Maestroni GJM, Villaggio B, Hertens E, Cravioito C, Pizzorni C, *et al*. Melatonin serum levels in rheumatoid arthritis. *Ann N Y Acad Sci* 2002;**966**:276–83.
 - 38 **Cutolo M**, Masi AT. Circadian rhythms and arthritis. *Rheum Dis Clin North Am* 2005;**31**:115–29.
 - 39 **Schulze-Koops H**, Lipsky PE, Kavanaugh AF, Davis LS. Elevated Th1 or Th2-like cytokine mRNA in peripheral circulation of patients with rheumatoid arthritis. *J Immunol* 1995;**155**:5029–37.
 - 40 **Maestroni GJM**, Sulli A, Cutolo M, Villaggio B, Cutolo M. Melatonin in rheumatoid arthritis: synovial macrophages show melatonin receptors. *Ann N Y Acad Sci* 2002;**966**:271–5.
 - 41 **Maestroni GJM**, Sulli A, Cutolo M, Villaggio B, Cutolo M. Melatonin in rheumatoid arthritis: a disease promoting and modulating hormone? *Clin Exp Rheumatol* 2002;**20**:872–3.
 - 42 **Cantorna MT**. Vitamin D and autoimmunity: is vitamin D status an environmental factor affecting autoimmune disease prevalence? *Proc Soc Exp Biol Med* 2000;**223**:230–3.
 - 43 **Rozin A**, Balbir-Gurman A, Chapira D. Seasonal distribution of relapse onset in rheumatoid arthritis and spondyloarthritis: the possible effect of solar factor. *Clin Exp Rheumatol* 2003;**21**:161–9.
 - 44 **Cutolo M**. Solar light effects on onset/releases and circannual/circadian symptomatology in rheumatoid arthritis. *Clin Exp Rheumatol* 2003;**21**:148–50.
 - 45 **Otsa K**, Peets T, Maestroni GJM, Aakre O, Veldi T, Hertens E. Circadian rhythms of melatonin in Estonian rheumatoid arthritis patients [abstract]. *Ann Rheum Dis* 2004;**63**(suppl):S139.
 - 46 **Cutolo M**, Maestroni GJM, Otsa K, Aakre O, Villaggio B, Capellino S, *et al*. Circadian melatonin and cortisol levels in rheumatoid arthritis patients in winter time: a north and south Europe comparison. *Ann Rheum Dis* 2005;**64**:212–16.
 - 47 **Straub RH**, Cutolo M. Involvement of the hypothalamic-pituitary-adrenal/gonadal axis and the peripheral nervous system in rheumatoid arthritis: viewpoint based on a systemic pathogenetic role. *Arthritis Rheum* 2001;**44**:493–507.
 - 48 **Cutolo M**, Villaggio B, Otsa K, Aakre O, Sulli A, Seriola B. Altered circadian rhythms in rheumatoid arthritis play a role in disease's symptoms. *Autoimmunity Rev*, (in press).
 - 49 **Abdel-Nasser AM**, Rasker JJ, Valkenburg HA. Epidemiological and clinical aspects relating to the variability of rheumatoid arthritis. *Semin Arthritis Rheum* 1997;**27**:123–40.
 - 50 **Spector TD**. Rheumatoid arthritis. *Rheum Dis Clin North Am* 1990;**16**:513–37.