4. **Jackson DW**, Evans NA, Thomas BD. Accuracy of needle placement into intra-articular space of the knee. *J Bone Joint Surg Am* 2002;**84**:1522–7.
5. **Balint PV**, Kane D, Hunter J, McInnes IB, Field M, Sturrock RD. Ultrasound guided versus conventional joint and soft tissue fluid aspiration in rheumatology practice: a pilot study. *J Rheumatol* 2002;**29**:2209–13.
6. **Raza K**, Lee CY, Pilling D, Heaton S, Situnayake RD, Carruthers DM, *et al*. Ultrasound guidance allows accurate needle placement from small joints in patients with early arthritis. *Rheumatology (Oxford)* 2003;**42**:976–9.
7. **Kane D**, Greaney T, Shanahan M, Duffy G, Bresnihan B, Gibney R, *et al*. The role of ultrasonography in the diagnosis and management of idiopathic plantar fasciitis. *Rheumatology (Oxford)* 2001;**40**:1002–8.
8. **Naredo E**, Cabero F, Beneyto P, Cruz A, Mondejar B, Uson J, *et al*. A randomized comparative study of short term response to blind injection versus sonographic-guided injection of local corticosteroids in patients with painful shoulder. *J Rheumatol* 2004;**31**:308–14.
9. **Furtado RN**, Oliveira LM, Natour J. Polyarticular corticosteroid injection versus systemic administration in treatment of rheumatoid arthritis patients: a randomized controlled study. *J Rheumatol* 2005;**32**:1691–8.
10. **van Vugt RM**, van Dalen A, Bijlma JWW. The current role of high-resolution ultrasonography of the hand and wrist in rheumatic diseases. *Clin Exp Rheumatol* 1998;**16**:454–8.

Correction

There was an error in the list of authors of a paper published in the May issue of the journal (Increased exposure to statins in patients developing chronic muscle diseases: a 2-year retrospective study. *Ann Rheum Dis* 2008;**67**:614–19). The correct list of authors is as follows: L Sailler, C Pereira, A Bagheri, E Uro-Coste, B Roussel, D Adoue, B Fournie, M Laroche, L Zabraniecki, P Cintas, P Arlet, M Lapeyre-Mestre, J L Montastruc.

Ann Rheum Dis 2008;**67**:1200. doi:10.1136/ard.2007.075523.corr1