M Mosca,¹ T Giuliano,¹ M Curcio,² M Doveri,¹ F De Feo,¹ C Tani,¹ L Bazzichi,¹ S Bombardieri¹

¹ Rheumatology Unit, Department of Internal Medicine, University of Pisa, Italy;

² Immunohematology and Transfusional Medicine Unit, Azienda Ospedaliera Universitaria Pisana, Pisa, Italy

Correspondence to: M Mosca, Rheumatology Unit, Department of Internal Medicine, University of Pisa, Via Roma 67, 56126 Pisa, Italy; marta.mosca@int.med.unipi.it

Competing interests: None declared.

Ethics approval: Ethics approval was obtained.

Accepted 24 March 2008

Ann Rheum Dis 2009;**68**:155–156. doi:10.1136/ard.2008.089821

REFERENCES

1. **Lambert NC**, Pang JM, Yan TD, Stevens AM, Furst DE, Nelson JL. Male microchimerism in women with systemic sclerosis and in healthy women who have never given birth to a son. *Ann Rheum Dis* 2005;**64**:845–8.
2. **Lambert NC**, Erickson TD, Yan Z, Pang JM, Guthrie KA, Furst DE, *et al*. Quantification of maternal microchimerism by an HLA-specific real time polymerase chain reaction. *Arthritis Rheum* 2004;**50**:906–14.
3. **Zhong XY**, Burk MR, Troeger C, Kang A, Holzgreve W, Hahn S. Fluctuation of maternal and fetal free extra-cellular circulatory DNA in the maternal plasma. *Obstet Gynecol* 2000;**96**:991–6.
4. **Mosca M**, Curcio M, Lapi S, Valentini G, D'Angelo S, Rizzo G, Bombardieri S. Correlations of Y chromosome microchimerism with disease activity in patients with SLE: analysis of preliminary data. *Ann Rheum Dis* 2003;**62**:651–4.
5. **Siebert PD**, Larrick JW. PCR MIMICS: competitive DNA fragments for use as internal standards in quantitative PCR. *BioTechniques* 1993;**14**:244.
6. **Lo YM**, Tein MS, Lau TK, Haines CJ, Leung TN, Poon PM, *et al*. Quantitative analysis of fetal DNA in maternal plasma and serum: implications for non-invasive pre-natal diagnosis. *Am J Hum Genet* 1998;**64**:768–75.
7. **Knoop C**, Andrien M, Defleur V, Antonine M, de Francquen P, Goldman M, *et al*. Detection of blood chimerism after lung and heart-lung transplantation: The superiority of nested as compared with standard polymerase chain reaction amplification. *Transplantation* 1994;**58**:1335–8.
8. **Carter AS**, Bunce M, Cerundolo L, Welsh KI, Morris PJ, Fuggle SV. Detection of microchimerism after allogeneic blood transfusion using nested polymerase chain reaction amplification with sequence-specific primers: a cautionary tale. *Blood* 1998;**92**:683–9.
9. **Bein G**, Glaser R, Kirchner H. Rapid HLA-DRB1 genotyping by nested PCR amplification. *Tissue Antigens* 1992;**39**:68–73.
10. **Valentini G**, Della Rossa A, Bombardieri S, Bencivelli W, Silman AJ, D'Angelo S, *et al*. European multi-centre study to define disease activity criteria for systemic sclerosis. II. Identification of disease activity variables and development of preliminary activity indexes. *Ann Rheum Dis* 2001;**60**:592–8.

Correction

The following article was originally published with an incorrect digital object identifier (doi): Song IH, Althoff CE, Hermann KG, Scheel AK, Knetsch T, Schoenharting M, *et al*. Knee osteoarthritis. Efficacy of a new method of contrast-enhanced musculoskeletal ultrasonography in detection of synovitis in patients with knee osteoarthritis in comparison with magnetic resonance imaging. *Ann Rheum Dis* 2008;**67**:19–25. It has been updated with the correct doi: 10.1136/ard.2007.067462. We apologise for any inconvenience caused.

Ann Rheum Dis 2009;**68**:156. doi:10.1136/ard.2007.067462.corr2