Background and objectives CD55 (decay-accelerating factor) is a complement-regulatory protein, which is highly expressed by fibroblast-like synoviocytes (FLS). CD55 is also a ligand for CD97, an adhesion-type G protein-coupled receptor abundantly present on leucocytes. We recently showed that lack of either CD55 or CD97 ameliorates disease in murine collagen-induced and K/BxN serum transfer models of arthritis.¹ Little is known regarding the regulation of CD55 expression in FLS. We therefore investigated the effect of toll-like receptors ligation and pro-inflammatory cytokines on CD55 expression.

Materials and methods Synovial fibroblasts, obtained from biopsy samples of arthritis patients, were cultured and stimulated with cytokines (TNF, IFN γ , IL-1 β , IL-6, IFN α) or TLR ligands (LTA, poly (I:C), LPS, imiquimod, CpG). Expression of CD55 was measured by flow cytometry using domain-specific monoclonal antibodies and recombinant CD97-loaded fluorescent beads. Chloroquine was used to inhibit TLR3 activity. Upregulation and functionality of dsRNA sensors in response to poly (I:C) or 5'-triphosphate RNA was analysed by PCR. Apoptosis was measured by PI/annexinV staining and was blocked with the pan-caspase inhibitor Q-VD-OPH.

Results Cultured synovial fibroblasts of patients with rheumatoid arthritis (RA), osteoarthritis, psoriatic arthritis, and spondylarthritis express equal amount of CD55. Stimulation of RA-FLS with IL-1 β (p=0.02) and poly (I:C) (p=0.001) induced a significant upregulation of CD55. Engagement of TLR3 by the dsRNA analog poly (I:C) was confirmed using chloroquine, an inhibitor of endosomal acidification that impairs TLR3 signaling. Synovial fibroblasts also expressed the cytoplasmic dsRNA sensors melanoma differentiation-associated gene 5 (MDA5) and retinoic acid-inducible gene I (RIG-I). Stimulation of these receptors with either poly (I:C) or 5'-triphosphate RNA induced CD55 expression, but, in case of MDA5, also induced significant cell death (p<0.001) that was caspase-dependent. Upregulation of CD55 in response to dsRNA receptor activation increased the binding capacity of synovial fibroblasts for CD97-loaded beads.

Conclusions We identify dsRNA as a potent inducer of CD55 upregulation on synovial fibroblasts. Our findings suggest that CD55 induction by viral dsRNA or dsRNA may facilitate the accumulation of CD97-expressing inflammatory immune cells in the synovial tissue.

REFERENCE

1. **Hoek RM**, de LD, Kop EN, Yilmaz-Elis AS, Lin F, Reedquist KA, Verbeek JS, Medof ME, Tak PP, Hamann J. Deletion of either CD55 or CD97 ameliorates arthritis in mouse models. *Arthritis Rheum* 2010;**62**:1036–1042.

A61 TRIGGERING OF VIRAL RNA SENSORS INDUCES CD55 EXPRESSION ON SYNOVIAL FIBROBLASTS

Karpus O N,¹ Heutinck K M,^{1,2} Wijnker P J M,¹ Tak P P,³ Hamann J¹ ¹Department of Experimental Immunology, Academic Medical Center, Amsterdam, The Netherlands; ²Renal Transplant Unit, Academic Medical Center, Amsterdam, The Netherlands; ³Division of Clinical Immunology and Rheumatology, Academic Medical Center, Amsterdam, The Netherlands

10.1136/ard.2010.149104.5