A10

IRF8 ALLELE ASSOCIATED WITH SUSCEPTIBILITY TO MULTIPLE SCLEROSIS IS ASSOCIATED WITH SERUM INTERFERON α AND SEROLOGIC PROFILE IN SYSTEMIC LUPUS ERYTHEMATOSUS

Beverly S Franek,¹ Silvia N Kariuki,¹ Jasmine Arrington,¹ Rachel A Mikolaitis,² Meenakshi Jolly,² Tammy O Utset,¹ Mary Zervou,³ Dimitrios T Boumpas,³ George Goulielmos,³ Timothy B Niewold¹ ¹Section of Rheumatology and Gwen Knapp Center for Lupus and Immunology Research, University of Chicago, Chicago, Illinois, USA; ²Section of Rheumatology, Rush University Medical Center, Chicago, Illinois, USA; ³Internal Medicine and Rheumatology, Clinical Immunology and Allergy, University of Crete Medical School, Heraklion, Greece

10.1136/ard.2010.149096.10

Objective Alleles of IRF8 have been associated with susceptibility to both systemic lupus erythematosus (SLE) and multiple sclerosis (MS). While interferon α (IFN- α) is thought to be causal in SLE, recombinant human IFN- α is used as a therapy in MS. We investigated whether the IRF8 alleles associated with these two diseases were associated with differences in serum IFN- α or serologic profile in a multi-ancestral cohort of SLE patients.

Methods The rs12444486 and rs17445836 single nucleotide polymorphisms (SNPs) in IRF8 (associated with SLE and MS respectively) were genotyped with Taqman primer-probe sets. We studied 548 SLE patients (258 African-American, 147 European-American, and 143 Cretan) and 526 matched controls (298 African-American, 117 European-American, and 111 Cretan). All patients had serum IFN- α and serology data available, and had been previously genotyped at SLE-risk SNPs in the IRF5 and IRF7 loci. Data from each ancestral background

was analysed separately initially, and combined in meta-analysis when associations were not significantly heterogeneous between ancestral backgrounds. Principal component analysis was used to control for proportional ancestry at the individual level in logistic regressions.

Results In case-control meta-analysis, we observed a stronger and more consistent cross-ancestral background association between IRF8 and SLE at the MS-associated rs17445836 SNP than at the previously reported SLE-associated rs12444486 SNP (OR=0.66, $p=2.7 \times 10^{-3}$, Cochran's $Q=0.80$). The MS-associated rs17445836 G allele was associated with the presence of anti-dsDNA autoantibodies in SLE patients of both ancestral backgrounds (meta-analysis OR=2.01 (1.09–3.68), $p=0.024$). The same allele was also associated with increased serum IFN- α activity in both ancestral backgrounds (meta-analysis $p=0.017$). There was no evidence for statistical interaction between rs17445836 G and SNPs in IRF5 and IRF7 which have been previously associated with anti-dsDNA in SLE patients. No significant associations were observed between the SLE-associated rs12444486 SNP and serum IFN- α or serologic profile.

Conclusions The rs17445836 G allele associated with susceptibility to MS was associated with SLE, and with anti-dsDNA antibodies and serum IFN- α in SLE patients of both African-American and European ancestry. This is interesting, given the therapeutic effect of IFN- α in MS patients, and the pathogenic effect of this same cytokine in SLE. Further exploration of the impact of the IRF8 locus upon in vivo IFN- α levels should provide insight into both diseases.