A206 SKEWED X CHROMOSOME INACTIVATION IN RHEUMATOID ARTHRITIS WOMEN

Doua F Azzouz,¹ Onur Emre Onat,² Nathalie Balandraud,^{1,3} Sami B Kanaan,¹ Jean Roudier,^{1,3} Tayfun Ozcelik,² Nathalie C Lambert¹ ¹INSERM UMR639, Marseille, France; ²Department of Molecular Biology and Genetics, Bilkent University, Ankara, Turkey; ³Rheumatology department, Hôpital La Conception, Marseille, France

10.1136/ard.2010.149021.16

Background and objectives It has long been recognised that women have a greater prevalence of autoimmune diseases. Rheumatoid arthritis (RA) does not escape this rule with a women:men ratio of 3:1.

X chromosome inactivation (XCI) is a dosage compensation mechanism used by mammals to ensure that XX females and XY males equalise X chromosome gene expression. As a consequence, females are a mosaic of two cell lines, one expressing maternal X-linked and the other expressing paternal X-linked genes with a ratio close to 50:50 when XCI is random.

However skewing, defined as a deviation from the 50:50 ratio has been described in females with autoimmune thyroid diseases, scleroderma and juvenile idiopathic arthritis (for review¹). The aim of the current study is to test whether women with RA also have a skewed XCI.

Methods The highly polymorphic CAG repeat on the first exon of the androgen receptor gene was genotyped, as described elsewhere² to determine XCI bias in 84 women with RA and 100 healthy women.

Results A total of 54 patients and 69 controls were informative for androgen receptor polymorphism. Among them 31.5% of women with RA (17/54) had a skewed XCI ($\geq 80:20$) compared to only 17.4% of healthy women (12/69). Only extreme skewing was statistically significant with 18.5% of patients following this pattern and 2.9% of controls ($p=0.004$).

Conclusions Our preliminary data indicate that skewed XCI may be a risk factor for the occurrence of RA in women. Further studies need to be done to analyse whether women who have a skewed pattern have less genetic susceptibilities (shared epitope) or less specific autoantibodies (anti-CCP) as their risk factor is X chromosome linked.

REFERENCE

1. Lambert NC. The price of silence. *Arthritis Rheum* 2009;**60**:3164–7.
2. Ozbalkan Z, Bagislar S, Kiraz S, et al. Skewed X chromosome inactivation in blood cells of women with scleroderma. *Arthritis Rheum* 2005;**52**:1564–70.