EXTENDED REPORT

Variation in the *ICAM1–ICAM4–ICAM5* locus is associated with systemic lupus erythematosus susceptibility in multiple ancestries

Kwangwoo Kim,¹ Elizabeth E Brown,² Chan-Bum Choi,³ Marta E Alarcón-Riquelme on behalf of BIOLUPUS,^{4,5} Jennifer A Kelly,⁴ Stuart B Glenn,⁴ Joshua O Ojwang,⁴ Adam Adler,⁴ Hye-Soon Lee,³ Susan A Boackle,⁶ Lindsey A Criswell,⁷ Graciela S Alarcón,² Jeffrey C Edberg,² Anne M Stevens,⁸ Chaim O Jacob,⁹ Gary S Gilkeson,¹⁰ Diane L Kamen,¹⁰ Betty P Tsao,¹¹ Juan-Manuel Anaya,¹² Joel M Guthridge,⁴ Swapan K Nath,⁴ Bruce Richardson,¹³ Amr H Sawalha,^{4,14,15} Young Mo Kang,¹⁶ Seung Cheol Shim,¹⁷ Chang-Hee Suh,¹⁸ Soo-Kon Lee,¹⁹ Chang-sik Kim,²⁰ Joan T Merrill,⁴ Michelle Petri,²¹ Rosalind Ramsey-Goldman,²² Luis M Vil,²³ Timothy B Niewold,²⁴ Javier Martin,²⁵ Bernardo A Pons-Estel, on behalf of GENLES,²⁶ Timothy J Vyse,²⁷ Barry I Freedman,²⁸ Kathy L Moser,⁴ Patrick M Gaffney,⁴ Adrienne Williams,²⁸ Mary Comeau,²⁸ John D Reveille,²⁹ Judith A James,¹⁴ R Hal Scofield,¹⁴ Carl D Langefeld,²⁸ Kenneth M Kaufman,^{4,15} John B Harley,³⁰ Changwon Kang,¹ Robert P Kimberly,² Sang-Cheol Bae³

► Additional figures and tables are published online only. To view these files, please visit the journal online (<http://ard.bmj.com/content/early/2012/04/26/ard.2011.201110>).

For numbered affiliations see end of article.

Correspondence to

Changwon Kang, Korea Advanced Institute of Science and Technology, Daejeon 305-701, Republic of Korea; ckang@kaist.ac.kr or Robert P Kimberly, The University of Alabama at Birmingham, Birmingham, Alabama 35294, USA; Robert.Kimberly@ccc.uab.edu, or Sang-Cheol Bae, Hanyang University Hospital for Rheumatic Diseases, Seoul 133-792, Republic of Korea; scbae@hanyang.ac.kr

Received 30 November 2011

Accepted 24 February 2012

Published Online First
20 April 2012

KK and EEB contributed equally to this work and are joint first authors. CK, RPK and SCB contributed equally to this work and are joint corresponding authors.

ABSTRACT

Objective Systemic lupus erythematosus (SLE; OMIM 152700) is a chronic autoimmune disease for which the aetiology includes genetic and environmental factors. *ITGAM*, integrin α_M (complement component 3 receptor 3 subunit) encoding a ligand for intracellular adhesion molecule (ICAM) proteins, is an established SLE susceptibility locus. This study aimed to evaluate the independent and joint effects of genetic variations in the genes that encode *ITGAM* and *ICAM*.

Methods The authors examined several markers in the *ICAM1–ICAM4–ICAM5* locus on chromosome 19p13 and the single *ITGAM* polymorphism (rs1143679) using a large-scale case-control study of 17 481 unrelated participants from four ancestry populations. The single-marker association and gene-gene interaction were analysed for each ancestry, and a meta-analysis across the four ancestries was performed.

Results The A-allele of *ICAM1–ICAM4–ICAM5* rs3093030, associated with elevated plasma levels of soluble ICAM1, and the A-allele of *ITGAM* rs1143679 showed the strongest association with increased SLE susceptibility in each of the ancestry populations and the trans-ancestry meta-analysis ($OR_{meta} = 1.16$, 95% CI 1.11 to 1.22; $p = 4.88 \times 10^{-10}$ and $OR_{meta} = 1.67$, 95% CI 1.55 to 1.79; $p = 3.32 \times 10^{-46}$, respectively). The effect of the *ICAM* single-nucleotide polymorphisms (SNPs) was independent of the effect of the *ITGAM* SNP rs1143679, and carriers of both *ICAM* rs3093030-AA and *ITGAM* rs1143679-AA had an OR of 4.08 compared with those with no risk allele in either SNP (95% CI 2.09 to 7.98; $p = 3.91 \times 10^{-5}$).

Conclusion These findings are the first to suggest that an ICAM integrin-mediated pathway contributes to susceptibility to SLE.

Systemic lupus erythematosus (SLE; OMIM 152700) is a chronic autoimmune disease for which the aetiology includes diverse genetic and environmental factors. The formation and deposition of immune complexes, which consist of nuclear autoantigens and autoantibodies, in vital organs result in chronic inflammation, organ failure and severe morbidity and mortality in patients with SLE.

Previous genome-wide and candidate-gene association studies reported an association of SLE with a non-synonymous variant in *ITGAM*, encoding integrin α_M (complement component 3 receptor 3 subunit), among European and African descents.¹ *ITGAM* is a ligand for intercellular adhesion molecule (ICAM) proteins, suggesting the involvement of an integrin–ICAM-mediated adhesion pathway for SLE susceptibility.

In this candidate gene-association study, we examined the independent and combined effects of common variation in *ITGAM* and the *ICAM1–ICAM4–ICAM5* locus, encoding other components of the integrin–adhesion pathway, with SLE susceptibility using a large multi-ancestry population.

METHODS

Study population

As part of the first phase of this study, we included a total of 14 719 patients with SLE and controls from diverse ancestry backgrounds, recruited from multiple institutions worldwide as part of the Lupus Association Study (LLAS)-2, with approval from the appropriate institutional review boards (online supplementary table S1). The affected had a minimum of four of eleven 1997 American College of Rheumatology revised criteria for the classification of SLE.³ All participants in this study were filtered

by principal component analysis to identify population outliers as previously reported.⁴ The sample exceeding five SDs along any statistically significant principal component were defined as outliers and removed from the study.

For the second phase to demonstrate consistency of results for rs3093030 in this larger Korean population, we constituted an independent cohort consisting of 2762 SLE cases and controls of Korean ancestry. Hospital-based SLE cases and controls were recruited from six hospitals in Korea with approval from the respective institutional review boards (online supplementary table S1).

Genotyping

We genotyped the 12 *ICAM* variants in phase I (rs5030340, rs5030390, rs5030391, rs5030351, rs5491, rs1799969, rs5498, rs5030400, rs3093032, rs281437, rs3093030, rs2228615), which tagged most of the common single-nucleotide polymorphisms (SNPs) in the *ICAM1-ICAM4-ICAM5* locus, in addition to an *ITGAM* variant (rs1143679), using customised arrays based on the Illumina iSelect platform at the Lupus Genetics Studies Unit of the Oklahoma Medical Research Foundation. For the SLE-associated SNP in the *ICAM1-ICAM4-ICAM5* locus with the strongest effect (rs3093030), we genotyped this marker for the phase II Korean cohort using the Sequenom iPLEX platform at the Korea Advanced Institute of Science and Technology.

SNP association with SLE susceptibility

The statistical assessment for SNP association with SLE susceptibility was performed by the χ^2 test with allelic genetic model for each ancestry and meta-analysis across the four ancestries using the PLINK 1.07 (<http://pngu.mgh.harvard.edu/~purcell/plink/>).⁵ In the meta-analysis, homogeneity of effect sizes was evaluated using the I^2 heterogeneity index and Cochrane's Q statistic.⁶ For $p > 0.01$ in heterogeneity Q statistic, fixed-effects models were implemented; otherwise random-effects models were used. For meta-analysis, GWAMA 1.4 software (<http://www.well.ox.ac.uk/gwama/>) was also used to calculate OR and 95% CI.⁷ Hardy-Weinberg equilibrium (HWE) was examined by comparison between genotype distributions expected and observed in controls using the χ^2 test. Pair-wise D' and r^2 values between SNPs were calculated by the Haploview 4.2 program. Statistical power was estimated using the CaTS-Power calculator (<http://www.sph.umich.edu/csg/abecasis/CaTS/index.html>).

Imputation

Using the genotype data for the 12 *ICAM* SNPs, 58 HapMap SNPs located in a chromosome 19 region from position 10 200 kb to 10 320 kb (build 36.3) were imputed using the MACH V.1.0 program (<http://www.sph.umich.edu/csg/abecasis/MACH/index.html>) with the ancestry-matched template data provided by the International HapMap Project (<http://www.hapmap.org>) Phase 3 (Genome Browser release 2). Europeans were imputed using the CEU (Utah residents with Northern and Western European ancestry) data, Africans using the ASW (African ancestry in Southwest USA) data, Hispanics using the MEX (Mexican ancestry in Los Angeles, California) data, and Koreans using the CHB (Han Chinese in Beijing, China) data. Among the 58 imputed SNPs, 69.0% in Europeans, 77.6% in Africans, 58.6% in Hispanics and 63.8% in Koreans showed high (≥ 0.70) average posterior probability for the most likely genotype, although some SNPs were not confidently imputed because of their weak correlations with the 12 genotyped SNPs.

Gene-gene interaction

Epistatic interactions between the seven SLE-associated SNPs in the *ICAM1-ICAM4-ICAM5* locus and the non-synonymous SNP rs1143679 in *ITGAM* were examined using multiplicative logistic regression with the affected status as a dependent factor, two SNPs as main factors and the two-SNP interaction term. The analysis was performed using the PLINK '--epistasis' option.^{5,8} In addition, case-only analysis using the PLINK '--fast-epistasis' and '--case-only' options was used to examine the dependency or the correlation between the tested SNPs in the groups of SLE cases.^{5,8} To access combined effects of the risk alleles in rs3093030 and rs1143679, fixed-effects OR and 95% CI were estimated in stratification by the number of risk alleles (0 to 4) as compared with the group having no risk allele, separately among the European, African and Hispanic populations using the Case-control And Tdt Meta-Analysis Package (<http://cran.r-project.org/web/packages/catmap/index.html>). Two-copy carriers homozygous for *ICAM* or *ITGAM* risk allele were removed from the two-risk-allele carrier group in order to see the combined effect.

SNP association with gene expression

Using the Genevar 2.0.1 database, which includes SNP genotype and mRNA profiling data for fibroblasts, lymphoblastoid cell lines, and T cells from the 75 GenCord individuals,^{9,10} the 19 SNPs located in a chromosome 19 region from position 10 200 kb to 10 320 kb were assessed for their associations with mRNA levels of *ICAM1*, *ICAM4* and *ICAM5* using a t statistic for Spearman's rank correlation coefficient (ρ). To minimise the type I error caused by multiple testing, the significance level of the p value was set to $\alpha = 0.05/(19 \text{ SNPs})/(\text{five probes})/(\text{three cell lines}) = 1.75 \times 10^{-4}$.

RESULTS

ICAM-SLE association study

In phase I, we evaluated 12 common variants directly typed in the *ICAM1-ICAM4-ICAM5* locus spanning ~22 kb in a total of 14 719 unrelated SLE cases and controls of European ($n=7427$; SLE/controls=3936/3491), African ($n=3613$; 1679/1934), Hispanic ($n=2299$; 1492/807) and Korean ($n=1380$; 640/740) ancestries (online supplementary table S1). Proportions of genetic ancestry for the study subjects were filtered using principal component analysis as described in a previous study (online supplementary figure S1).⁴ All variants were typed with call rates of over 97.9%, and genotype proportions were consistent with expectations of the HWE ($p \geq 0.001$) among controls from each of the four ancestries (online supplementary table S2).

Of the 12 *ICAM* SNPs typed, five were significantly associated with SLE susceptibility under a statistical threshold of $p \leq 4.17 \times 10^{-3}$ in the trans-ancestry meta-analysis after Bonferroni correction for multiple comparisons (rs5030390, rs5498, rs281437, rs3093030 and rs2228615; $1.05 \times 10^{-8} \leq p_{\text{meta}} \leq 2.80 \times 10^{-4}$) (online supplementary table S2). Moreover, the directions of the effect sizes of each SNP were same among the ancestries except for rs5030390, the association of which was considered unreliable (online supplementary table S2).

The strongest association with SLE was found with rs3093030 located in the 0.4 kb intergenic region between *ICAM1* and *ICAM4* ($p=2.49 \times 10^{-8}$; OR=1.16, 95% CI 1.10 to 1.22). The effect size was similar in each of the four ancestries (online supplementary table S2), with OR=1.12 (95% CI 1.05 to 1.20) in European, OR=1.18 (95% CI 1.01 to 1.37) in African, OR=1.28 (95% CI 1.13 to 1.45) in Hispanic and OR=1.15 (95% CI 0.98

Table 1 Association of directly typed and imputed markers in the *ICAM* locus and *ITGAM* with susceptibility to systemic lupus erythematosus

		European (n=7427)		African (n=3613)		Hispanic (n=2299)		Korean (n=1380 or 4142) [†]		Meta-analysis (n=17 481) [‡]	
Gene	SNP*	MAF (SLE/controls)	OR	MAF (SLE/controls)	OR	MAF (SLE/controls)	OR	MAF (SLE/controls)	OR	OR	p Value
ICAM1–ICAM4–ICAM5											
	rs3093030 (G>A)	(0.471/0.442)	1.12	(0.107/0.092)	1.18	(0.580/0.518)	1.28	(0.353/0.318)	1.17	1.16	4.88×10 ^{−10}
	rs2569693 (C>T)	(0.427/0.403)	1.11	(0.097/0.085)	1.16 [§]	(0.550/0.492)	1.26	(0.363/0.332)	1.14 [§]	1.14	4.50×10 ^{−7}
	rs5498 (A>G)	(0.470/0.445)	1.11	(0.188/0.173)	1.11 [§]	(0.582/0.524)	1.26	(0.414/0.384)	1.13 [§]	1.13	6.08×10 ^{−7}
	rs2228615 (G>A)	(0.427/0.403)	1.11	(0.098/0.087)	1.14 [§]	(0.552/0.496)	1.25	(0.346/0.319)	1.13 [§]	1.14	1.05×10 ^{−6}
	rs2569702 (T>C)	(0.431/0.408)	1.10	(0.097/0.085)	1.16 [§]	(0.552/0.494)	1.26	(0.345/0.318)	1.13 [§]	1.14	1.27×10 ^{−6}
	rs892188 (C>T)	(0.431/0.408)	1.10	(0.097/0.085)	1.16 [§]	(0.552/0.494)	1.26	(0.345/0.318)	1.13 [§]	1.14	1.27×10 ^{−6}
	rs281437 (G>A)	(0.276/0.288)	0.94 [§]	(0.339/0.379)	0.84 [§]	(0.175/0.220)	0.75	(0.088/0.092)	0.95 [§]	0.89	8.28×10 ^{−6}
ITGAM											
	rs1143679 (G>A)	(0.194/0.126)	1.67	(0.157/0.104)	1.82	(0.168/0.100)	1.59	(0.002/0.000)	NC	1.67	3.32×10 ^{−46}

*Four SNPs (rs3093030, rs5498, rs2228615 and rs281437) were genotyped and the other three SNPs (rs2569693, rs2569702 and rs892188) were imputed. The major allele is more common than the minor allele in the whole control population but not necessarily in each subpopulation.

[†]A total of 4142 Korean participants from phase I and phase II were genotyped for rs3093030, and the separate analyses are shown in online supplementary table S2.

[‡]Fixed-effects p value and OR were calculated from the meta-analysis because the effect size of each SNP was not significantly different among the ancestry populations ($p > 0.01$ in Cochran's Q statistic). For *ITGAM* rs1143679, the meta-analysis was performed with European, African and Hispanic results.

[§]Allelic associations with SLE were not statistically significant at a threshold of $\alpha = 0.05$. The detailed information is shown in online supplementary table S2.

MAF, minor allele frequency; NC, not calculated; SLE, systemic lupus erythematosus; SNP, single-nucleotide polymorphism.

to 1.35) in Korean ancestries ($p = 0.307$ in Q statistic, $I^2 = 17.0$). In a conditional logistic regression analysis followed by a meta-analysis, we found that the SLE association of rs5498, rs281437 and rs2228615 accounted for that of rs3093030, and they presented a consensus signal of SLE susceptibility locus, possibly because they are in high linkage disequilibrium with rs3093030 ($D' \geq 0.84$; online supplementary table S3).

Although the effect magnitude of rs3093030 among Koreans was similar to those of the other ancestry populations, it did not achieve statistical significance, in contrast with the others ($p = 0.0841$ in Koreans versus $p = 5.95 \times 10^{-4}$ in Europeans, $p = 4.02 \times 10^{-2}$ in Africans and $p = 7.28 \times 10^{-5}$ in Hispanics), perhaps because of the small sample size of the Korean population relative to the other ancestry populations. With the minor allele frequency of 0.324 in the Korean controls, the statistical power to detect the effect size found in the meta-analysis (fixed-effects OR=1.16) was only 42% in the Korean population.

To increase the statistical power for Korean detection, in phase II we recruited an additional independent population of 2762 unrelated Korean participants (online supplementary table S1). Genotyping call rate was 96.9%, and the control subject genotypes were distributed under HWE ($p = 0.691$). Indeed, a similar effect size was detected in the validation population (OR=1.18), and by combining the phase I and phase II populations of Koreans ($n = 4142$), statistical significance ($p = 1.07 \times 10^{-3}$; OR=1.17, 95% CI 1.07 to 1.29) was achieved (online supplementary table S2). Therefore, with the additional Korean validation population, each of the four ancestry populations independently showed a statistically significant association between the *ICAM* locus SNP rs3093030 and SLE susceptibility, and the meta-analysis across the four ancestral populations revealed a more significant overall relationship ($p = 4.88 \times 10^{-10}$; OR=1.16, 95% CI 1.11 to 1.22; online supplementary table S1).

To further characterise this genomic region as an SLE susceptibility locus, we imputed HapMap SNPs around the *ICAM* gene cluster that included *EDG5*, *MRPL4*, *ICAM1*, *ICAM4*, *ICAM5*, *RAVER1* and *ICAM3*. Using the genotype data for the 12 *ICAM* SNPs, we imputed 58 HapMap SNPs located in a chromosome 19 region from position 10 200 kb to 10 320 kb (Genome build 36.3) using the MACHV1.0 program, with the ancestry-matched haplotype templates provided by the HapMap3 Genome Browser release 2 of the International HapMap Project.

Three imputed SNPs (rs2569693, rs2569702 and rs892188) with high imputation quality (average posterior probability for the most likely genotype ≥ 0.97) were significantly associated with SLE susceptibility (table 1) while demonstrating the same direction of SLE-risk effects in each of the SNPs. In addition, all SLE-associated SNPs showed larger effects in Hispanics than in the other ethnic groups. These imputed SLE-associated SNPs were localised to the *ICAM1-ICAM4-ICAM5* region and correlated highly with the genotyped SLE-associated SNP rs3093030 ($0.81 \leq r^2 \leq 0.96$). Trans-ancestry comparisons for the seven SLE-associated SNPs (ie, four genotyped and three imputed) supported that the *ICAM1-ICAM4-ICAM5* locus alone associated sufficiently with SLE susceptibility ($p \leq 8.28 \times 10^{-6}$; figure 1).

Pathway-based *ICAM* and *ITGAM* interaction study

Because *ICAM1* and *ICAM4* are binding partners of an $\alpha_M\beta_2$ integrin, in which the α subunit is encoded by an SLE-susceptibility gene, *ITGAM*, we examined the statistical interaction between SLE-risk alleles of the *ICAM1-ICAM4-ICAM5* region and the previously reported SLE-risk alleles of the *ITGAM* SNP rs1143679 to characterise the pathway-based effect on SLE susceptibility.² For this analysis, all participants of the phase I population were also genotyped for the non-synonymous SNP rs1143679 located in exon 3 of *ITGAM*. Call rates were $\geq 99.6\%$ within each ancestry population, and genotype distributions were consistent with expectations of HWE in each of the control groups (online supplementary table S2).

The overall fixed effect of the SLE-risk-associated A allele in the *ITGAM* SNP was OR=1.67 (95% CI 1.55 to 1.79, $p = 3.32 \times 10^{-46}$), showing similar effect sizes among European, African and Hispanic ancestries ($p = 0.541$ in Q-statistic, $I^2 = 0.00\%$; online supplementary table S2), although the risk-associated allele was too rare (0.07%) to evaluate the interaction among Koreans. The infrequency in Koreans was consistent with the previous report.¹¹ Statistical interaction between *ITGAM* rs1143679-A and each of the seven SLE-associated SNPs in the *ICAM1-ICAM4-ICAM5* locus ($p \leq 4.17 \times 10^{-3}$; table 1) were examined among cases and controls using the PLINK '--epistasis' analysis and among only the cases using the PLINK '--fast-epistasis' analysis.

The *ITGAM* and *ICAM* SNPs independently affected SLE susceptibility, showing no notable *ICAM-ITGAM* epistatic interactions ($p > 0.05$; online supplementary table S4). Using five

Figure 1 A systemic lupus erythematosus association map for the genotyped (squares) and imputed single-nucleotide polymorphisms (SNPs) (diamonds) within and around the *ICAM1-ICAM4-ICAM5* locus. Negative logarithms (base 10) of the p values calculated from the meta-analysis (y axis) are plotted against SNP positions on human chromosome 19 (x axis). The direction of effect size was the same in all four ancestries for some SNPs (filled symbols) but not for others (empty symbols). The arrows indicate transcription directions and gene sizes.

subgroups according to the number of risk-associated alleles (0 to 4) in the *ICAM* SNP, rs3093030, and the *ITGAM* SNP, rs1143679, we assessed the combined effects of the SLE risk-associated alleles in the *ICAM*-integrin pathway. Fixed-effects OR and 95% CI for SLE susceptibility in each subgroup were estimated with reference to the subgroup with no risk allele. The natural logarithm of OR correlated linearly ($r^2=0.9803$) with the number of risk alleles (0 to 4), and four-copy carriers had OR of 4.08 for risk of SLE compared with carriers with 0 copies (figure 2).

Gene expression

To determine whether the genetic variation has a role in gene expression, we evaluated expression quantitative trait loci (eQTL) associations within the *ICAM* locus using the Genevar (GENe Expression VARIation) database (<http://www.sanger.ac.uk/resources/software/genevar/>), which publicly provides data for 75 GenCord individuals on both SNP genotypes and expression profiles from primary fibroblasts, lymphoblastoid cell lines and T cells.^{9 10} The 19 Genevar SNPs included rs2569693, rs5498, rs2228615 and rs281437 among the SLE-associated SNPs. However, none of the evaluated markers were associated with aberrant mRNA levels of *ICAM1*, *ICAM4* or *ICAM5* in any cell type (online supplementary figure S2).

DISCUSSION

To our knowledge, this is the first report of a comprehensive approach used to evaluate the effect of the *ICAM1-ICAM4-ICAM5* locus on SLE risk. We observed that several SNPs within this locus are associated with susceptibility to SLE in multiple ancestry populations.

ICAM1 is mainly expressed in the vascular endothelium, macrophages and lymphocytes, and plays a role in immunological events including extravasation and T-cell-mediated responses.¹² However, a plausible role for *ICAM4* and *ICAM5* in SLE pathogenesis is less clear because they are preferentially expressed in red blood cells and brain, respectively.^{13 14}

The *ICAM* polymorphisms were not associated with altered mRNA levels of *ICAM1*, *ICAM4* or *ICAM5* in the Genevar database, although the sample size ($n=75$) was too small to make this null association conclusive. However, soluble *ICAM1* levels

Figure 2 Combined effects of *ICAM1-ICAM4-ICAM5* rs3093030-A and *ITGAM* rs1143679-A alleles on systemic lupus erythematosus (SLE) susceptibility. Natural logarithms of the relative odds of SLE (y axis) are plotted against the number of at-risk alleles (x axis) in the two putative SLE susceptibility single-nucleotide polymorphisms (SNPs), rs3093030 in *ICAM1-ICAM4-ICAM5* and rs1143679 in *ITGAM*. OR values are provided next to the data points, and error bars represent corresponding 95% CI. A linear trend line (dashed) and r^2 are shown.

in plasma have been associated with two SNPs (rs3093030 and rs5498) in the *ICAM1-ICAM4-ICAM5* locus by protein QTL analysis in two independent studies involving 6578 European women ($p=5.9 \times 10^{-23}$ and $p=4.8 \times 10^{-25}$) and 9813 European individuals ($p=3.5 \times 10^{-23}$ and $p=2.5 \times 10^{-21}$).^{15 16} The alleles that were associated with increased soluble *ICAM1* in those studies (A in rs3093030 and G in rs5498) were also associated with increased susceptibility to SLE in this study, providing a plausible mechanism for their effect. In addition, the genotype-phenotype relationships are consistent with several previously reported observations of raised plasma levels of soluble *ICAM1* in patients with SLE,¹⁷⁻²¹ suggesting an underlying aetiological role.

Soluble *ICAM1* is a cleaved form consisting of the extracellular domain of the membrane-bound full-length *ICAM1* that is capable of binding to ligands. Thus soluble and membrane-bound *ICAM1* can compete with each other for binding to ligands,¹² but the immunological role of either soluble or membrane-bound *ICAM1* in SLE pathogenesis has not been characterised. The level of soluble *ICAM1* probably correlates with that of membrane-bound *ICAM1*,^{12 19} but it is not known whether the increase of soluble *ICAM1* in SLE accompanies with the concurrent increase of the membrane form due to increased gene expression or with the reciprocal decrease of the membrane form due to increased proteolytic processing, as the membrane-bound *ICAM1* levels have not been measured in patients with SLE. None of the Genevar SNPs in the *ICAM1-ICAM4-ICAM5* locus was associated with *ICAM1* mRNA expression. Instead, the non-synonymous *ICAM1* SNP rs5498 (encoding Lys469Glu) was associated with both increased levels of the soluble form of *ICAM1* protein and increased SLE susceptibility. Of note is the fact that it is located at an Ig superfamily domain, which may affect the proteolytic conversion to the soluble *ICAM1*. However, the level of statistical significance for the SLE association of rs5498 was weaker than that of rs3093030 in all populations, and statistical significance was not reached in the African or Korean ancestry populations. Thus it is possible that rs5498 affects SLE susceptibility in some ancestry populations but not others.

In this study, we also showed the association of *ITGAM* rs1143679 in European, African and Hispanic descendants, which was also found in previous association studies for systemic sclerosis.²² This common variant was monomorphic in Koreans, which is consistent with the literature of other Asian populations,¹¹ perhaps reflecting selective pressures in this population. *ITGAM* (also known as CD11b, Mac-1 and complement receptor type 3) is a well-characterised molecule in the integrin α chain family that is expressed in a number of myeloid cells including macrophages, monocytes and neutrophils.^{23–25} *ITGAM* encodes the α chain of $\alpha_M\beta_2$ integrin, which regulates neutrophil and monocyte adhesion and migration from the bloodstream via interactions with a wide range of structurally unrelated ligands, including but not limited to ICAM1 and ICAM2.^{26 27}

The previously identified SLE-susceptibility *ITGAM* SNP, rs1143679, encodes a change in amino acid from arginine at position 77 to histidine (R77H). This conversion of amino acids induces an alteration in the tertiary and quaternary structures of the ligand-binding $\alpha_M\beta_2$ domain, thereby modifying its overall binding affinity.²⁸ Nath *et al* suggest that the R77H conversion may ultimately influence the conformation of the α 1 domain (specific to the ligand-binding domain for ICAM1), with subsequent consequences on $\alpha_M\beta_2$ ligand binding.²

Indeed, evidence suggests that $\alpha_M\beta_2$ levels are increased on neutrophils in SLE patients with active disease and may contribute to endothelial injury, consistent with the cumulative organ damage associated with SLE.²⁹ $\alpha_M\beta_2$ is also involved in immune complex clearance, a process that is impaired in patients with SLE,³⁰ suggesting that the function of $\alpha_M\beta_2$ may be altered in these people, either directly by structural modification or indirectly by alterations in receptor–ligand binding, possibly resulting from structural modification.²

In addition, this conversion is a target for alloantibodies present in mothers of neonates affected with neonatal autoimmune neutropenia.²⁸ Notably, alloantibodies reactive against the polymorphic $\alpha_M\beta_2$ molecule are capable of blocking the $\alpha_M\beta_2$ -dependent adhesion of neutrophils and monocytic U937 cells to a number of molecules including ICAM1.²⁸

The functional association of *ICAM* and *ITGAM* prompted us to examine the associated SNPs in these genes for epistatic effects. Although a joint role for both *ICAM1* and *ITGAM* in SLE aetiology appears to be plausible, they independently affected SLE susceptibility in this study. This may indicate that the functional role of the *ICAM* variants is involved in a non-integrin-mediated pathway or that some factors are affected by ICAM–ITGAM binding through different pathways of ICAM and ITGAM without a common pathway.

In summary, we identified the association of several SNPs within the *ICAM1–ICAM4–ICAM5* locus with SLE susceptibility. Although the effect size was modest, the association was consistent across all four ancestries. Our findings suggest that common variations in genes involved in ICAM-mediated or ICAM/integrin-mediated adhesion play a causal role in SLE susceptibility.

Contributors KK, EEB, JAK, JOO, SKN, BR, AHS, KLM, PMG, JBH, CK, RPK and S-CB designed the study. EEB, C-BC, MEA-R, H-SL, SAB, LAC, GSA, JCE, AMS, COJ, GSG, DLK, BPT, J-MA, JMG, YMK, SCS, C-HS, S-KL, C-SK, JTM, MP, RRG, LMV, TBN, JM, BAP-E, TJV, BIF, KLM, PMG, JDR, JAJ, RHS, JBH, RPK and S-CB recruited and characterised the SLE cases and controls. KK, AA and KMK performed genotyping. SBG, AW, MC and CDL performed quality control analyses. KK and EEB performed statistical analyses. KK, EEB, C-BC, CK, RPK and S-CB prepared the manuscript. All authors approved the final draft.

Acknowledgements We thank the study participants and physicians who provided samples (Peter K Gregersen, So-Young Bang, BIOLUPUS Network and GENLES Network) and technical support (Taehyeung Kim). The members of the BIOLUPUS Network who provided samples used in this study are Sandra D Alfonso, Raffaella

Scorza, Mauro Galleazzi, Gian Domenico Sebastiani, MD Danieli, and Sergio Migliarese in Italy, Bernard R Lauwerys in Belgium, Eموke Endrefly and L. szl–Kov cs in Hungary, Carlos Vasconcelos and Berta Martins da Silva in Portugal, I–igo R a Figueroa, Carmen Gutierrez, Ana Su rez, Norberto Ortego, Jos Luis Calleja, Enrique de Ram–n Garrido, Julio Sanchez Rom n, and Mario Sabio in Spain, Helle Lastrup and Peter Junker in Denmark, and Marc Bijl and Cees Kallenberg in Holland; Dr Lennart Tuedsson provided controls from Southern Sweden (Lund). The members of the GENLES Network who provided samples used in this study are Hugo R Scherbarth, Pilar C Marino, Estela I Motta, Susana Gamron, Cristina Drenkard, Emilia Menso, Alberto Allievi, Guillermo A Tate, Jose L Presas, Simon A Palatnik, Marcelo Abdala, Mariela Bearzotti, Alejandro Alvarellos, Francisco Caeiro, Ana Bertoli, Sergio Paira, Susana Roverano, Cesar E Graf, Estela Bertero, Cesar Caprarulo, Griselda Buchanan, Carolina Guiller–n, Sebastian Grimaudo, Jorge Manni, Luis J Catoggio, Enrique R Soriano, Carlos D Santos, Cristina Prigione, Fernando A Ramos, Sandra M Navarro, Guillermo A Berbotto, Marisa Jorfen, Elisa J Romero, Mercedes A Garcia, Juan C Marcos, Ana I Marcos, Carlos E Perandones, Alicia Eimon and Cristina G Battagliotti in Argentina; Eduardo Acevedo and Mariano Cucho in Per ; Ignacio Garc’a de la Torre, Mario Cardiel R’os, Jos Francisco Moctezuma and Marco Maradiaga Cece–a in Mexico.

Funding The work was supported by grants from US NIH (AI063622, AI071651, AI082714, AI083194, AI094377, AI24717, AR042460, AR043274, AR043814, AR044804, AR048940, AR052300, AR053483, AR058554, AR060366, AR43727, AR62277, CA141700-01, GM063483, HD07463, K08-AI083790, K24-AR002138, M01-RR00079, N01-AR-6-227, P01-AR49084, P30-DK42086, P30-AR055385, P60-AR049459 P60-AR053308, P60-2-AR30692, P01-AR49084, PR094002, R01-AR33062, R21-AI070304, RR015577, RR020143, UL1RR024999, UL1RR025005, UL1RR025741, UL1RR029882 and 5UL1RR025777), the Lupus Foundation of America, the Alliance for Lupus Research, a Kirkland Scholar Award, the US Department of Veterans Affairs, the Korean Healthcare Technology Research and Development Project (A111218-11-GM01), the Korea Research Program for New Drug Target Discovery (20090083335), the National Research Foundation of Korea (2010-0014162), the Lupus Research Institute Novel Research Grant, the Alliance for Lupus Research Target Identification in Lupus Grant, the Arthritis National Research Foundation Eng Tan Scholar Award, the ESF in the framework of the Research Networking Programme European Science Foundation The Identification of Novel Genes and Biomarkers for Systemic Lupus Erythematosus (BIOLUPUS) 07-RNP-083, the Swedish Research Council, the Swedish Association against Rheumatism, the Swedish International Development Agency, the Gustaf Vth 80th-Jubilee Foundation, the Instituto de Salud Carlos III (PS09/00129) partly financed by the FEDER funds of the European Union, and a grant from the Consejer’a de Salud de la Junta de Andaluc’a.

Competing interests None.

Patient consent Obtained.

Ethics approval The study was approved by the Institutional Review Board of all participating institutions.

Provenance and peer review Not commissioned; externally peer reviewed.

Author affiliations ¹Department of Biological Sciences, Korea Advanced Institute of Science and Technology, Daejeon, Korea

²Departments of Medicine and Epidemiology, Schools of Medicine and Public Health, The University of Alabama at Birmingham, Birmingham, Alabama, USA

³Department of Rheumatology, Hanyang University Hospital for Rheumatic Diseases, Seoul, Korea

⁴Arthritis and Clinical Immunology Program, Oklahoma Medical Research Foundation, Oklahoma City, Oklahoma, USA

⁵Area of Human DNA Variability, Centro de Gen–mica e Investigaci–n Oncol–gica (GENYO), P zer-Universidad de Granada-Junta de Andaluc’a, Granada, Spain

⁶Division of Rheumatology, University of Colorado Denver School of Medicine, Aurora, Colorado, USA

⁷Rosalind Russell Medical Research Center for Arthritis, University of California, San Francisco, California, USA

⁸University of Washington, Seattle Children’s Hospital, Seattle, Washington, USA

⁹The Lupus Genetic Group, University of Southern California, Los Angeles, California, USA

¹⁰Department of Medicine, Medical University of South Carolina, Charleston, South Carolina, USA

¹¹David Geffen School of Medicine, University of California, Los Angeles, California, USA

¹²Center for Autoimmune Diseases Research, Universidad del Rosario, Bogota, Colombia

¹³Division of Rheumatology, University of Michigan and US Department of Veterans Affairs Medical Center, Ann Arbor, Michigan, USA

¹⁴Department of Medicine, University of Oklahoma Health Sciences Center, Oklahoma City, Oklahoma, USA

¹⁵Department of Medicine, US Department of Veterans Affairs Medical Center, Oklahoma City, Oklahoma, USA

¹⁶Department of Internal Medicine (Rheumatology), Kyungpook National University School of Medicine, Daegu, Korea

¹⁷Division of Rheumatology, Department of Medicine, Eulji Medi-Bio Research Institute, Eulji University, Daejeon, Korea

¹⁸Department of Rheumatology, Ajou University School of Medicine, Suwon, Korea

¹⁹Department of Internal Medicine, Yonsei University College of Medicine, Seoul, Korea

²⁰Department of Ophthalmology, Chungnam National University School of Medicine, Daejeon, Korea

²¹Department of Medicine, Johns Hopkins University School of Medicine, Baltimore, Maryland, USA

²²Northwestern University, Feinberg School of Medicine, Chicago, Illinois, USA

²³Department of Medicine University of Puerto Rico Medical Sciences Campus, San Juan, Puerto Rico

²⁴Gwen Knapp Center for Lupus and Immunology Research, University of Chicago, Chicago, Illinois, USA

²⁵Department of Immunology, Instituto de Biomedicina y Parasitología López-Neyra, CSIC, Granada, Spain

²⁶Department of Medicine, Sanatorio Parque, Rosario, Argentina

²⁷King's College London, Divisions of Genetics and Molecular Medicine and Immunology, Infection and Inflammatory Disease, Guy's Hospital, London, UK

²⁸Departments of Biostatistical Sciences and Internal Medicine, Wake Forest School of Medicine, Winston-Salem, North Carolina, USA

²⁹Division of Rheumatology, The University of Texas Health Science Center at Houston, Houston, Texas, USA

³⁰Pediatrics and US Department of Veterans Affairs Medical Center, Cincinnati Children's Hospital Medical Center, Cincinnati, Ohio, USA

REFERENCES

- Harley JB, Alarcón-Riquelme ME, Criswell LA, et al. Genome-wide association scan in women with systemic lupus erythematosus identifies susceptibility variants in ITGAM, PXK, KIAA1542 and other loci. *Nat Genet* 2008;**40**:204–10.
- Nath SK, Han S, Kim-Howard X, et al. A nonsynonymous functional variant in integrin- α (M) (encoded by ITGAM) is associated with systemic lupus erythematosus. *Nat Genet* 2008;**40**:152–4.
- Hochberg MC. Updating the American College of Rheumatology revised criteria for the classification of systemic lupus erythematosus. *Arthritis Rheum* 1997;**40**:1725.
- Adrianto I, Wen F, Templeton A, et al. Association of a functional variant downstream of TNFAIP3 with systemic lupus erythematosus. *Nat Genet* 2011;**43**:253–8.
- Purcell S, Neale B, Todd-Brown K, et al. PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* 2007;**81**:559–75.
- Huedo-Medina TB, Sánchez-Meca J, Marín-Martínez F, et al. Assessing heterogeneity in meta-analysis: Q statistic or I² index? *Psychol Methods* 2006;**11**:193–206.
- Mägi R, Morris AP. GWAMA: software for genome-wide association meta-analysis. *BMC Bioinformatics* 2010;**11**:288.
- Cordell HJ. Detecting gene-gene interactions that underlie human diseases. *Nat Rev Genet* 2009;**10**:392–404.
- Dimas AS, Deutsch S, Stranger BE, et al. Common regulatory variation impacts gene expression in a cell type-dependent manner. *Science* 2009;**325**:1246–50.
- Yang TP, Beazley C, Montgomery SB, et al. Genevar: a database and Java application for the analysis and visualization of SNP-gene associations in eQTL studies. *Bioinformatics* 2010;**26**:2474–6.
- Han S, Kim-Howard X, Deshmukh H, et al. Evaluation of imputation-based association in and around the integrin- α -M (ITGAM) gene and replication of robust association between a non-synonymous functional variant within ITGAM and systemic lupus erythematosus (SLE). *Hum Mol Genet* 2009;**18**:1171–80.
- van de Stolpe A, van der Saag PT. Intercellular adhesion molecule-1. *J Mol Med* 1996;**74**:13–33.
- Gahmberg CG, Tian L, Ning L, et al. ICAM-5: a novel two-faceted adhesion molecule in the mammalian brain. *Immunol Lett* 2008;**117**:131–5.
- Toivanen A, Ihanus E, Mattila M, et al. Importance of molecular studies on major blood groups: intercellular adhesion molecule-4, a blood group antigen involved in multiple cellular interactions. *Biochim Biophys Acta* 2008;**1780**:456–66.
- Paré G, Chasman DI, Kellogg M, et al. Novel association of ABO histo-blood group antigen with soluble ICAM-1: results of a genome-wide association study of 6,578 women. *PLoS Genet* 2008;**4**:e1000118.
- Barbalic M, Dupuis J, Dehghan A, et al. Large-scale genomic studies reveal central role of ABO in sP-selectin and sICAM-1 levels. *Hum Mol Genet* 2010;**19**:1863–72.
- Egerer K, Feist E, Rohr U, et al. Increased serum soluble CD14, ICAM-1 and E-selectin correlate with disease activity and prognosis in systemic lupus erythematosus. *Lupus* 2000;**9**:614–21.
- Sabry A, Sheashaa H, El-Husseini A, et al. Intercellular adhesion molecules in systemic lupus erythematosus patients with lupus nephritis. *Clin Rheumatol* 2007;**26**:1819–23.
- Sfikakis PP, Tsokos GC. Clinical use of the measurement of soluble cell adhesion molecules in patients with autoimmune rheumatic diseases. *Clin Diagn Lab Immunol* 1997;**4**:241–6.
- Tso TK, Huang WN. Elevated soluble intercellular adhesion molecule-1 levels in patients with systemic lupus erythematosus: relation to insulin resistance. *J Rheumatol* 2007;**34**:726–30.
- Tulek N, Aydinoglu O, Ozoran K, et al. Soluble intercellular adhesion molecule-1 (sICAM-1) in patients with systemic lupus erythematosus. *Clin Rheumatol* 1996;**15**:47–50.
- Anaya JM, Kim-Howard X, Prahalad S, et al. Evaluation of genetic association between an ITGAM non-synonymous SNP (rs1143679) and multiple autoimmune diseases. *Autoimmun Rev* 2012;**11**:276–80.
- Abbas AR, Baldwin D, Ma Y, et al. Immune response in silico (IRIS): immune-specific genes identified from a compendium of microarray expression data. *Genes Immun* 2005;**6**:319–31.
- Hynes RO. Integrins: versatility, modulation, and signaling in cell adhesion. *Cell* 1992;**69**:11–25.
- Lu H, Smith CW, Perrard J, et al. LFA-1 is sufficient in mediating neutrophil emigration in Mac-1-deficient mice. *J Clin Invest* 1997;**99**:1340–50.
- Fagerholm SC, Varis M, Stefanidakis M, et al. α -Chain phosphorylation of the human leukocyte CD11b/CD18 (Mac-1) integrin is pivotal for integrin activation to bind ICAMs and leukocyte extravasation. *Blood* 2006;**108**:3379–86.
- Li Y, Zhang L. The fourth blade within the beta-propeller is involved specifically in C3bi recognition by integrin α M β 2. *J Biol Chem* 2003;**278**:34395–402.
- Sachs UJ, Chavakis T, Fung L, et al. Human alloantibody anti-Mart interferes with Mac-1-dependent leukocyte adhesion. *Blood* 2004;**104**:727–34.
- Buyon JP, Shadick N, Berkman R, et al. Surface expression of Gp 165/95, the complement receptor CR3, as a marker of disease activity in systemic lupus erythematosus. *Clin Immunol Immunopathol* 1988;**46**:141–9.
- Hepburn AL, Mason JC, Wang S, et al. Both Fc γ and complement receptors mediate transfer of immune complexes from erythrocytes to human macrophages under physiological flow conditions in vitro. *Clin Exp Immunol* 2006;**146**:133–45.