The distinct pathogenesis of IgG4 Mikulicz's disease and Sjogren's syndrome in Japan; the role of IL-21 and IL-6

Robert Irving Fox

Maehara *et al* present an interesting study on 'Interleukin-21 contributes to germinal centre formation and immunoglobulin G4 production in IgG4-related dacryoadenitis and sialoadenitis, so-called Mikulicz's disease (MD)'.¹ They have studied 12 Japanese patients with MD (IgG4-DS), 15 with Sjogren's syndrome (SS) and 15 healthy controls. The labial salivary biopsies (LSB) of 15 SS patients were chosen from a larger group of SS patients (n=66) based on the presence of ectopic germinal centres (GC) in 15 SS LSBs studied. All the MD patients were chosen on the basis of their elevated IgG4. The SS patients were chosen on the basis of ectopic GC in their biopsies. The mRNA expression of IL-21 and BCL-6 was significantly higher in MD than in SS labial salivary gland biopsies. The IL-21 was detected by immunohistochemistry in and around GC only in the MD patients. These results were consistent with the hypothesis that follicular T-helper cells (Tfh) play a key role in the formation of GC in MD. By contrast, IL-17 was not found in either SS or MD. This article extends a recent study by the same group² in Kyushu that distinguished MD and SS by showing elevation of Th2 and Tregs (ie, IL4, IL10 and FoxP3 expression by immunohistochemistry); and IgG4 plasma cells in the MD labial biopsies. Over the past decade, this group has published a number of excellent papers on IgG4-related diseases, including sclerosing cholangitis, multicentric Castleman's disease and other lymphoproliferative disorders.³⁻⁵

The article is carefully presented by experts in the field, but is a bit confusing to the practicing rheumatologist due to the myriad of different cytokines and

receptors described. Several key issues for the general rheumatologist include:

- What is the relationship between Japanese MD patients (now called IgG4-related dacryoadenitis) and MD patients outside Japan?
- What is the relationship of the IgG4-related MD dacryoadenitis patients and SS patients in Japan?
- What is the significance of finding MD-like lesions in biopsies of SS patients, as these lesions have been reported over the past 50 years;^{6,7} are we dealing with a characteristic IL-21-based process or a different pathogenesis leading to a similar histopathology?

MD was originally considered to be a subtype of SS, based on histopathologic similarities between the two diseases.^{8,9} The term refers to a 'benign' lymphoepithelial lesion of the parotid and/or lacrimal glands. The name was originally derived from Jan Mikulicz-Radecki, the Polish surgeon who described these conditions in 1892.¹⁰ There is a marked lymphoplasmacytic infiltration of the glands. Lymphoid follicles surround solid epithelial nests, giving rise to the 'epimyoepithelial islands' that are mainly composed of degenerating ductal and myoepithelial cells.¹¹ Excess hyaline basement membrane material is deposited between cells, and there is also acinar atrophy and destruction. When the myoepithelial islands were found in a biopsy, it was felt that an inflammatory disease was more likely than a lymphoma.¹²

After Mikulicz's initial report of pathologic specimens, biopsies with similar dense lymphoid infiltrates and myoepithelial islands were reported in association with other diseases, such as tuberculosis, sarcoidosis, lymphoma¹³ and, subsequently, a similar pathology was noted in parotid biopsies from HIV-infected patients.¹⁴ Thus, the term Mikulicz syndrome fell out of favour by the 1930s as lacking specificity.¹⁵ Indeed,

the report by Henrik Sjogren in 1933¹⁵ described a more limited spectrum of patients with lymphoid infiltrates of salivary and lacrimal glands associated with keratoconjunctivitis sicca (KCS) and by excluding the above conditions; his initial reports were largely ignored in the medical literature for the next 20 years. The term SS gained a rebirth after pathologic reports by Morgan in 1953,¹³ and a *New England Journal* clinical pathology conference (CPC case) in 1954;¹⁶ literally, the diagnosis of SS was 'rediscovered' 20 years after the initial reports by Sjogren, since the initial report was not translated from the German literature.

Benign lymphoepithelial lesions labelled as MD (in patients lacking features of SS) were initially reported predominantly in Caucasian males around 50 years of age.¹⁶ In this study of IgG4 MD from Japan, the majority of patients (66%) were female. The lesions were usually unilateral in previous reports of Caucasian patients, and the symmetry of lesions is not specified in the present report from Japan. In the initial MD reports by Mikulicz, some patients with biopsies containing lymphoepithelial lesions had clinical features of KCS.¹⁶ When KCS features were present, the swelling was usually bilateral, and patients were predominantly female with a slightly younger age distribution.¹³ The concept of 'benign' lymphoepithelial lesion was later questioned,¹² since MD patients among Caucasian patients were uniformly found to have clonal immunoglobulin rearrangements and elevated frequency of lymphoma.¹⁷ The monocytic B-cell infiltrates in the SS biopsies with myoepithelial lesions exhibited IgM clonal rearrangements.¹⁸ Thus, the IgG4 MD patients in Japan may represent a distinct subgroup from non-Japanese MD patients previously reported.¹² Only future reports of non-Japanese MD patients will allow us to make comparisons.

Maehara *et al* present an elegant 'bench to bedside' translation of molecular biology to histology.¹ Due to the preferential and characteristic localisation of lymphoid infiltrates in the lacrimal and parotid glands, either vascular adhesion molecules or their ligands must direct the 'homing' of lymphocytes. In the IgG4-related MD reported in this article, the homing receptor seems to involve CXCR5. Their particular histologic organisation into their ectopic GC appears to involve Th2 cells (IL-4 and Tregs),² follicular T-helper cells and IL-21.

By contrast, SS biopsies of salivary glands (in Europe) exhibit CXCL and

Scripps Memorial, La Jolla, California, USA

Correspondence to Dr Robert Irving Fox, Scripps Memorial, Suite #911 9850 Genesee Ave, #910, La Jolla, CA 92037, USA; robertfoxmd@mac.com

type 1 interferon signature.²⁰ In a recent paper by the same group (Tanaka *et al*²), a different pattern of adhesive molecules was shown in IgG4 MD and SS. Similarly, the cytokine expressions (IL-21 and IL-6) also differ between the MD and SS patients in their cohort. Thus, IgG4 MD and SS patients in Japan differ in the factors that initiate and maintain the lymphoid infiltrates as well as the clinical features of each group.

A key issue for western rheumatologists is the association of IgG4 with MD, SS or sclerosing cholangitis in our practice. Little hard data exist on the global incidence and prevalence of IgG4-related disease, including MD or SS.²¹ Virtually all studies, including the present study, pertain to the disease in Japan. Although it has been a decade since the IgG4-related diseases were reported from Japan, there remains a paucity of reports of IgG4-related disease from other parts of the world.²¹ It may be that particular environmental and epigenetic factors in Japan cause immunologically predisposed individuals to develop these particular histologic responses. By analogy, an unusual variant of SS is found in Kyushu (Japan), and was later found to be due to a variant of HTLV-1 viral infection.²²

In summary, Maehara *et al* have extended their previous papers on IgG4-related MD.¹ IgG4 MD in Japan is clinically and pathogenetically distinct from SS in the same geographic region. Although some biopsies from SS patients contain histologic features of myoepithelial islands and ectopic GC, the pathogenetic processes leading to this histopathology are distinct from IgG4 MD. The frequency of IgG4-related MD in regions outside Japan remains unknown, and so, it is difficult to assess the significance of these findings in our MD, SS or sclerosing cholangitis

patients. Nevertheless, it is an elegant study in describing the factors leading to a particular pathologic biopsy in a group of Japanese patients. For the readers, it is time to urge our pathologists to review our MD cases and determine the relationship with the patients in this study.

Competing interests None.

Provenance and peer review Commissioned; externally peer reviewed.

Accepted 8 September 2012

Ann Rheum Dis 2012;**71**:1919–1920.
doi:10.1136/annrheumdis-2012-201866

REFERENCES

1. Maehara T, Moriyama M, Nakashima H, *et al*. Interleukin-21 contributes to germinal centre formation and immunoglobulin G4 production in IgG4-related dacryoadenitis and sialoadenitis, so called Mikulicz's disease. *Ann Rheum Dis* 2012;**71**:2011–19.
2. Tanaka A, Moriyama M, Nakashima H, *et al*. Th2 and regulatory immune reactions contribute to IgG4 production and the initiation of Mikulicz disease. *Arthritis Rheum* 2012;**64**:254–63.
3. Masaki Y, Dong L, Kurose N, *et al*. Proposal for a new clinical entity, IgG4-positive multiorgan lymphoproliferative syndrome: analysis of 64 cases of IgG4-related disorders. *Ann Rheum Dis* 2009;**68**:1310–5.
4. Sato Y, Kojima M, Takata K, *et al*. Systemic IgG4-related lymphadenopathy: a clinical and pathologic comparison to multicentric Castleman's disease. *Mod Pathol* 2009;**22**:589–99.
5. Hamano H, Kawa S, Horiuchi A, *et al*. High serum IgG4 concentrations in patients with sclerosing pancreatitis. *N Engl J Med* 2001;**344**:732–8.
6. Fishleder A, Tubbs R, Hesse B, *et al*. Uniform detection of immunoglobulin-gene rearrangement in benign lymphoepithelial lesions. *N Engl J Med* 1987;**316**:1118–21.
7. Morgan WS. The probable systemic nature of Mikulicz's disease and its relation to Sjögren's syndrome. *N Engl J Med* 1954;**251**:5–10.
8. Yamamoto M, Takahashi H, Sugai S, *et al*. Clinical and pathological characteristics of Mikulicz's disease (IgG4-related plasmacytic exocrinopathy). *Autoimmun Rev* 2005;**4**:195–200.
9. Yamamoto M, Takahashi H, Ohara M, *et al*. A new conceptualization for Mikulicz's disease as an IgG4-related plasmacytic disease. *Mod Rheumatol* 2006;**16**:335–40.
10. Mikulicz JH. Über eine eigenartige symmetrische Erkrankung der Tränen- und Mundspeicheldrüsen. In: Billroth GT, ed. *Beitr Chir Fortschr Stuttgart* 1892;610–30.
11. Dardick I, Rippstein P, Skimming L, *et al*. Immunohistochemistry and ultrastructure of myoepithelium and modified myoepithelium of the ducts of human major salivary glands: histogenetic implications for salivary gland tumors. *Oral Surg Oral Med Oral Pathol* 1987;**64**:703–15.
12. Batsakis JG, Bernacki EG, Rice DH, *et al*. Malignancy and the benign lymphoepithelial lesion. *Laryngoscope* 1975;**85**:389–99.
13. Morgan W, Castleman B. A clinicopathologic study of Mikulicz's disease. *Am J Pathol* 1953;**29**:471–503.
14. Ulirsch RC, Jaffe ES. Sjögren's syndrome-like illness associated with the acquired immunodeficiency syndrome-related complex*. *Hum Pathol* 1987;**18**:1063–8.
15. Sjögren HS. Zur kenntnis der keratoconjunctivitis sicca (Keratitis follicularis bei hypofunktion der tranendrüsen). *Acta Ophthalmol (Copenh)* 1933;**2**:1–151.
16. Morgan WS. The probable systemic nature of Mikulicz's disease and its relation to Sjögren's syndrome. *N Engl J Med* 1954;**251**:5–9.
17. Fishleder A, Tubbs R, Hesse B, *et al*. Immunoglobulin-gene rearrangement in benign lymphoepithelial lesions. *N Engl J Med* 1987;**316**:1118–21.
18. Shin SS, Sheibani K, Fishleder A, *et al*. Monocytoid B-cell lymphoma in patients with Sjögren's syndrome: a clinicopathologic study of 13 patients. *Hum Pathol* 1991;**22**:422–30.
19. Diehl SA, Schmidlin H, Nagasawa M, *et al*. STAT3-mediated up-regulation of BLIMP1 is coordinated with BCL6 down-regulation to control human plasma cell differentiation. *J Immunol* 2008;**180**:4805–15.
20. Hjelmervik TOR, Petersen K, Jonassen I, *et al*. Gene expression profiling of minor salivary glands clearly distinguishes primary Sjögren's syndrome patients from healthy control subjects. *Arthritis Rheum* 2005;**52**:1534–44.
21. Stone JH, Zen Y, Deshpande V. IgG4-Related Disease. *N Engl J Med* 2012;**366**:539–51.
22. Ohya Y, Nakamura S, Hara H, *et al*. Accumulation of human T lymphotropic virus type I, Æinfected T cells in the salivary glands of patients with human T lymphotropic virus type I, Æassociated Sjögren's syndrome. *Arthritis Rheum* 1998;**41**:1972–8.