subsequent studies have shown that citrullination of α -enolase is crucial for its autoantigenicity. α -enolase is an evolutionary conserved protein implicated both in glycolysis pathway and as a plasminogen receptor. Here, the authors have evaluated, in the well-known collagen induced arthritis model, the clinical, immunological and histological effects of both recombinant non-citrullinated α -enolase and immunodominant peptides from human and bacterial species.

Methods Different doses of α -enolase (10 and 100 μ g) or immunodominant enolase peptide 1 from human (hEP1) or *porphyromonas gingivalis* (pEP1) (10 or 100 μ g) were intraperitoneally injected to 6 week-old DBA/1 mice one day prior to collagen II arthritis induction (CIA). Both clinical (weight, arthritis score, tarsal thickness) and biological (anticollagen II and anti- α -enolase antibodies) were assessed during the 90 days follow-up period. Four histological score were also assessed: inflammation, synovial thickening, cartilage resorption and bone resorption.

Results Prophylactic injection of recombinant α -enolase was able to significantly prevent weight loss and to decrease the severity of arthritis evaluated by the arthritis score as well as the tarsal thickness. There was a dose-effect since 100 μ g led to better results. Levels of anticollagen II antibodies were significantly lower whereas titers of anti- α -enolase antibodies were significantly higher in mice treated with 100 μ g of α -enolase compared to control mice. Moreover, histological score were in agreement with clinical score. As regards to hEP1 and pEP1, the authors established a dose-dependant protective effect in CIA which is significant for pEP1. This protective effect is not due to once again a decrease of anti-collagen II antibodies titer.

Conclusion Prophylactic treatment with recombinant α -enolase suggest a protective role of this molecule. The clinical effect is not due to an immunological response mediated by anti- α -enolase antibodies. Prophylactic injection could induce either an immune deviation or an emergence of regulatory lymphocyte population, responsible of a decrease of anti-CII antibodies production. These results suggest α -enolase has an immunomodulatory effect in CIA mice. Those results suggest that non-citrullinated α -enolase could constitute a potential new therapeutic approach in RA.

4 PROPHYLACTIC INJECTION OF NON-CITRULLINATED α -ENOLASE HAS IMMUNOMODULATORY EFFECTS IN COLLAGEN-INDUCED ARTHRITIS MICE

Guillou C,¹ Avenel G,² Derambure C,^{1,2} Le Loarer F,³ Verdet M,² Hiron M,¹ Maho M,^{1,2} Le-Loet X,² Adriouch S,¹ Sabourin J C,³ Boyer O,¹ Lequerre T,^{1,2} Vittecoq O.^{1,2} ¹INSERM U905, Institut for Biomedical Research, University of Rouen, Rouen, France; ²Department of Rheumatology, Rouen University Hospital, Rouen, France; ³Laboratory of Histology, Rouen University Hospital, Rouen, France

10.1136/annrheumdis-2011-201237.4

Background Identification of autoantibodies associated with rheumatoid arthritis (RA) has been of major interest. In this context, the authors have previously identified for the first time α -enolase as a new auto-antigen in early RA. Moreover,