production and increased OC, two arthritis hallmarks. Here, we investigated whether miR-146a was differentially regulated in both Ly6C^{high} and Ly6C^{low} monocyte subsets in healthy and arthritic conditions. Moreover, we developed techniques to deliver siRNA to Ly6C^{high} monocytes in vivo and thus determined the specific impact of miR-146a overexpression in this particular monocyte subset on inflammation and osteoclastogenesis, in the context of arthritis.

Materials and Methods Subset monocytes were isolated from peripheral blood of arthritic and healthy mice by FACS sorting following membrane stainings. Transcriptomic analyses for miRNA expression were performed (n = 6) and differential miRNA expression levels were validated on individual samples using multiplex RT-qPCR (n = 13). Collagen-induced arthritic (CIA) mice were injected intravenously at disease onset with lipoplex containing either control or miR-146a mimic (0.5 mg/kg). Arthritis severity was monitored and OC differentiation was performed by stimulating bone marrow-derived monocytes (BMDM) with M-CSF and RANKL. miR-146a expression levels measured during OC differentiation. TRAP activity, OC numbers and nuclei per OC were quantified.

Results Transcriptomic analyses showed higher expression of miR-146a in Ly6C^{low} monocytes when compared to Ly6C^{high} monocytes, in both healthy and arthritis mice. In arthritis mice, expression of miR-146a in Ly6C^{high} monocytes is down-regulated as compared to healthy controls. During differentiation of BMDM into OC, miR-146a expression levels are down-regulated on day 2. Enforced expression of miR-146a in Ly6C^{high} monocytes upon intravenous injection of miR-146a-containing lipoplex leads to decreased bone erosion in mouse CIA. This effect was associated with a decreased number of mature OC and TRAP activity, as well as a reduced number of nuclei per OC.

Conclusions Overall, our results show that specific over-expression of miR-146a in Ly6C^{high} monocytes alters OC differentiation and decreased bone erosion in mouse CIA. These data also suggest that Ly6C^{high} monocytes might be the monocyte subset precursor of OC, and that targeting this specific monocyte subset might represent a therapeutic option in the context of arthritis to inhibit bone loss.

A9.18 POTENTIAL THERAPEUTIC APPLICATION OF HUMAN UMBILICAL CORD WHARTON JELLY DERIVED MESENCHYMAL STEM CELLS IN PRIMARY SJÖGREN'S SYNDROME

doi:10.1136/annrheumdis-2013-203223.18

¹Alessia Alunno, ²Pia Montanucci, ¹Sara Caterbi, ¹Onelia Bistoni, ²Giuseppe Basta, ¹Elena Bartoloni, ²Teresa Pescara, ¹Ilaria Pennoni, ¹Riccardo Calafiore, ¹Roberto Gerli. ¹Rheumatology Unit, Department of Clinical and Experimental Medicine; ²Section of Internal Medicine, Endocrine and Metabolic Sciences, University of Perugia, Perugia, Italy

Background and Objectives hUCMS are adult stem cells easy to retrieve in bulk, under acceptable ethical conditions. Their immunomodulatory and pro-differentiation properties have been widely demonstrated. Modulation of the immune system is mediated by both cell contact and soluble factors such as interferon (IFN)- γ produced by hUCMS. Very few evidence is currently available with respect to potential therapeutic application of hUCMS in systemic autoimmune disorders. In particular, no data have been published in primary Sjögren's syndrome (pSS) to date. Furthermore, we have recently developed an endotoxin-free alginate matrix which can be used to microencapsulate (CpS) different cell types and transfer them into a non-immunosuppressed host. These microcapsules containing pancreatic islets, upon approval by the Italian Ministry of Health for in vivo use, have been grafted into a cohort of patients with type 1 diabetes mellitus with no adverse effects, while proving to be immunoprotective. On this background, we aimed to assess the in vitro immune system modulation by IFN- γ pretreated CpS-hUCMS on T cells from pSS, with special regard to Th17- and Treg-cell subsets.

Materials and Methods Ten pSS patients and 5 healthy donors (HD) were enrolled. Peripheral blood mononuclear cells (PBMC) were obtained by density gradient from heparinised venous blood. Co-cultures of CpS-hUCMS and PBMCs were arranged at different ratios. Lymphocyte proliferation was assessed by CFSE dilution assay. Phenotypic analysis by flow cytometry for regulatory and effector T cell subpopulations was performed after culture. Real time PCR analysis and the evaluation of culture supernatants for cytokine expression are currently ongoing.

Results CpS-hUCMS were able to inhibit HD and pSS PBMC cell proliferation and this effect was inversely correlated to hUCMS number. Phenotypic analysis revealed a predominant Th17-cell response in SS which was promptly hampered by CpS-hUCMS. Moreover, Th1-cell proliferation was fair and reduced CpS-hUCMS FoxP3 and IL-17 expression among CD4⁺ T cells was also modulated by CpS-hUCMS, thereby suggesting that a Treg/Th17 rebalance.

Conclusions This is the first study evaluating the effects of hUCMS on T cells in pSS. It appeared that CpS-hUCMS rebalance Treg/Th17 ratio and therefore may exert therapeutic effects in such disease. Furthermore, it is the first study that employs a new technology of drug delivery which may be applied in vivo in pSS patients.

10. Others

A10.1 ANALYSIS OF THE MIGRATORY POTENTIAL OF DERMAL FIBROBLASTS OF PATIENTS WITH SYSTEMIC SCLEROSIS

doi:10.1136/annrheumdis-2013-203224.1

¹S Lefèvre, ¹FMP Meier, ²A Günther, ¹U Müller-Ladner, ¹E Neumann. ¹Dept Internal Medicine and Rheumatology, Justus-Liebig-University Gießen, Kerckhoff Clinic, Bad Nauheim, Germany; ²Dept Internal Medicine, Medical Clinic II, Justus-Liebig-University Gießen, Gießen

Background Systemic sclerosis (SSc) is characterised by fibroblast-mediated progressive skin fibrosis. Subsequent involvement of internal organs leads to severe impairments of their function and lung fibrosis represents the most common cause of death in SSc. Molecular mechanisms of the spreading of SSc are largely unknown. Recent results showed the migratory potential of rheumatoid arthritis (RA) synovial fibroblasts (SFs). Based on these results, the aim of this study was the analysis of migratory and adhesive properties of SSc dermal fibroblasts (DFs) and their role in SSc-spreading with focus on organ involvement.

Methods SScDFs (in part GFP-transfected) were injected intracutaneously into SCID mice. After 14–25 days, parts of the skin, internal organs and blood were analysed. Besides fluorescence analysis, immunohisto- and -cytochemistry was performed using species-specific antibodies to detect human cells. To avoid cross-reactivity of the antibodies, additional real time PCR analysis with primers for human β 2-microglobulin was performed after RNA isolation out of the respective organs and subsequent reverse transcription.

To analyse and to compare the adhesive behaviour of SScDFs with fibroblasts from other diseases and origins, multi-well culture plates were coated with Matrigel® (MG), growth factor-reduced (GFR) MG, or remained untreated. Cellular adhesion of SScDFs (n = 5), RASFs (n = 5), RADFs (n = 5), SFs (n = 2) and DFs of healthy individuals (n = 4) was analysed.

Results SScDFs or human cDNA, respectively, were not detected in any internal organ by fluorescence analysis, immunohistochemistry or real time PCR. Human SScDFs and human cDNA were only detectable in the murine skin at the injection site.

Fibroblasts of SSc patients, healthy SFs and DFs showed an increased adhesion to GFR MG compared to MG (SSc: GFR MG: 8.5 fold, MG: 8.2 fold; healthy SFs: GFR MG: 6.6 fold, MG: 4 fold; healthy DFs: GFR MG: 10 fold, MG: 7.6 fold). In contrast a reduction of cellular adhesion of RASFs (GFR MG: 4 fold, MG: 5.4 fold)