Response to: 'Bivalent HPV vaccine safety depending on subtypes of juvenile idiopathic arthritis' by Dr Akioka

We thank Dr Akioka for the comments¹ on our recent study on the bivalent human papillomavirus (HPV) vaccine in patients with juvenile idiopathic arthritis (JIA).² Akioka and coworkers describe a patient with enthesitis-related JIA after receiving the bivalent HPV vaccine. It was unclear whether this patient newly developed JIA or experienced a flare of pre-existing JIA.

With regard to the former, vaccinations are indeed often described as potential environmental triggers inducing autoimmune diseases in genetically susceptible individuals.³⁻⁴ For example, cases of systemic lupus erythematosus have been described after the quadrivalent HPV vaccine.⁵ However, in a large register-based cohort study, including 997 585 girls among whom 296 826 received a total of 696 420 quadrivalent HPV vaccine doses, no association between exposure to the quadrivalent HPV vaccine and autoimmune adverse events was found.⁶ Thus, large controlled studies often do not support a relation between vaccinations and the induction of autoimmune diseases except for the Guillain-Barré syndrome after the swine-flu vaccine and the oral polio vaccine, and idiopathic thrombocytopenia after the measles, mumps, rubella (MMR) vaccine.⁷ Nevertheless, the involvement of vaccinations in the pathogenesis of autoimmune diseases deserves further study, as it can improve our understanding of immunopathogenesis and thereby contribute to the prevention of disease. By no means the (alleged) association between vaccinations and autoimmune diseases should result in refraining from current immunisation practice, because the actual prevalence of autoimmune reactions to vaccinations is remarkably low, their course is often milder than those associated with natural infections and we cannot identify the individuals who are at risk of developing an autoimmune reaction after vaccination. Therefore, the benefits of infection prevention outweigh the small risk of developing an autoimmune disease in genetically susceptible individuals.

With regard to the induction of JIA disease flares after vaccination, case reports exist of JIA flares after rubella or influenza vaccination.⁷ Drawing conclusions from these case reports is impossible, because JIA has a relapsing and remitting disease course. To distinguish between a causal or temporal relation between vaccination and JIA disease flares, a (randomised) controlled study design is needed to correct for this relapsing disease course. We recently showed in a randomised controlled study that the MMR vaccination did not aggravate JIA.⁸ In the current study involving the HPV vaccine, patients served as their own controls in order to detect potential aggravation of JIA disease after vaccination. No detrimental effect of HPV vaccination on JIA disease activity was detected, not even in patients with high baseline disease activity, or patients using methotrexate or anti-tumour necrosis factor- α .

Aikoka points out that patients with JIA reported longer durations of arthralgia, fatigue and headache after vaccination compared with healthy controls and fear that JIA patients are more at risk for (severe) adverse events after HPV vaccination. However, we also show that the frequency of these complaints was similar in patients and healthy controls and was transient within 2 weeks in

all cases. We therefore conclude that patients with JIA and healthy controls tolerated the HPV vaccine equally well.

Aikoka suggests that polyenthesitis is an important factor in developing pain and adverse events caused by HPV vaccination, based on the fact that three patients developed inflammatory bowel disease (IBD), one uveitis and one lower back pain (of a non-inflammatory cause). We included only one patient with enthesitis-related JIA in this study. This patient did not develop severe adverse events. The four HLA-B27-positive patients also did not develop severe adverse events. We cannot draw conclusions on the risk of adverse events in enthesitis-related JIA because of these small patient numbers. Most importantly, we do not believe that these serious adverse events were related to the HPV vaccine, because the patients diagnosed with IBD already had abdominal pain, diarrhoea and intermittent rectal bleeding prior to the first HPV vaccine dose. Similarly, the treatment for uveitis in one oligoarticular JIA patient had been planned prior to HPV vaccination.

Although the serious adverse events in our study were in our opinion unrelated to HPV vaccination, we agree with Aikoka that it is important to remain vigilant for (severe) adverse events and to detect patients who are prone to develop (severe) adverse events after (HPV) vaccination. More studies are needed to discover means to identify patients who are individually predisposed to disease flares after vaccination. This might enable more personalised vaccination strategies in future.

M W Heijstek, N M Wulffraat

Department of Pediatric Immunology, University Medical Centre Utrecht, Utrecht, The Netherlands

Correspondence to Dr M W Heijstek, Department of Pediatric Immunology, University Medical Centre Utrecht, Room KC 03.063.00, PO BOX 85090, Utrecht 3508 AB, The Netherlands; m.w.heijstek@umcutrecht.nl

Contributors MWH and NMW wrote this letter and had final responsibility for the decision to submit for publication.

Competing interests None.

Patient consent Obtained.

Ethics approval CCMO The Hague, The Netherlands.

Provenance and peer review Commissioned; internally peer reviewed.

CrossMark

To cite Heijstek MW, Wulffraat NM. *Ann Rheum Dis* 2014;**73**:e76.

Received 19 August 2014

Accepted 12 September 2014

Published Online First 29 September 2014

► <http://dx.doi.org/10.1136/annrheumdis-2014-206426>

Ann Rheum Dis 2014;**73**:e76. doi:10.1136/annrheumdis-2014-206446

REFERENCES

- 1 Akioka S. Bivalent HPV vaccine safety depending on subtypes of juvenile idiopathic arthritis. *Ann Rheum Dis* 2014;**73**:e75.

- 2 Heijstek MW, Scherpenisse M, Groot N, *et al.* Immunogenicity and safety of the bivalent HPV vaccine in female patients with juvenile idiopathic arthritis: a prospective controlled observational cohort study. *Ann Rheum Dis* 2014;73:1500–7.
- 3 Wahren-Herlenius M, Dorner T. Immunopathogenic mechanisms of systemic autoimmune disease. *Lancet* 2013;382:819–31.
- 4 Wraith DC, Goldman M, Lambert PH. Vaccination and autoimmune disease: what is the evidence? *Lancet* 2003;362:1659–66.
- 5 Gatto M, Agmon-Levin N, Soriano A, *et al.* Human papillomavirus vaccine and systemic lupus erythematosus. *Clin Rheumatol* 2013;32:1301–7.
- 6 Arnheim-Dahlstrom L, Pasternak B, Svanstrom H, *et al.* Autoimmune, neurological, and venous thromboembolic adverse events after immunisation of adolescent girls with quadrivalent human papillomavirus vaccine in Denmark and Sweden: cohort study. *BMJ* 2013;347:f5906.
- 7 Heijstek MW, Ott de Bruin LM, Bijl M, *et al.* EULAR recommendations for vaccination in paediatric patients with rheumatic diseases. *Ann Rheum Dis* 2011;70:1704–12.
- 8 Heijstek MW, Kamphuis S, Armbrust W, *et al.* Effects of the live attenuated measles-mumps-rubella booster vaccination on disease activity in patients with juvenile idiopathic arthritis: a randomized trial. *JAMA* 2013;309:2449–56.