480 rabbits have found that by injecting young animals intra-peritoneally with the blood of patients with acute rheumatism there follows after an incubation of six weeks a thrombo-endocarditis that is transmissible in series from rabbit to rabbit up to five passages. They have never isolated any germ from the cardiac lesions or the blood, and conclude that the endocarditis is transmissible by contagion from one rabbit to another and that it is due to a filterable virus. This affection has never been seen in the normal control rabbits, but it has been found in rabbits injected with blood from normal subjects and those suffering from diseases other than rheumatism. Further work is in progress.

M. H. G.

CORRESPONDENCE

The Editor of THE RHEUMATIC DISEASES.

DEAR SIR,

I am very grateful for the review of my book, "The Synovial Membrane and the Synovial Fluid," which appeared in your journal.

Deeply convinced as I am of the shortcomings of human endeavour, I sincerely appreciate criticism of my own work. Unfortunately, Dr. Bywaters made errors which cast undeserved doubt on the work of qualified investigators quoted in my book. On p. 64, line 5 from the bottom, Dr. Bywaters writes: "Miller is quoted as saying that normal human joints are under a negative pressure of 6 to 12 mm. of water. This would need a magician to prove; and unfortunately Miller is not listed in the bibliography." In my book, p. 92, line 7 from the bottom, the paragraph referred to reads as follows: "Miller has found that normal joints of living humans and dogs are under a negative pressure which amounts to about 6 to 12 centimetres of water." Dr. Bywaters mistook centimetres for millimetres. The method employed by the author quoted, far from being magical, is based on simple and generally accepted physical determination of differences in pressure. In my text, the printer's proof-reader substituted the English spelling of the name of the author as Miller. In the bibliography the name is spelled correctly: "Mueller, Walter: *Biologie der Gelenke*. Leipzig, Verlag von Johann Ambrosius Barth, Leipzig, 1929."

I would like to point out one other statement concerning the work of two authors besides my own which is misleading. On p. 63, line 12 from the bottom, we read, "He cites Curtis and Brunschwig that cases of osteo-arthritis show a high rate of absorption (borne out by his own case in Table III., which gave the highest phenolsulphonephthalein clearance). This is taken to mean an increased local blood supply." The authors quoted (p. 53) are not Curtis and Brunschwig, but Curtis and Brandenburg. Neither they nor myself interpreted the rate of absorption of phenolsulphonephthalein from the joints as an indication of increased blood supply. It is only pointed out that adhesions which destroy the synovial membrane decrease the rate of excretion.

You will understand that in requesting a correction of these mis-statements I only act out of a feeling of obligation towards the authors whom I have quoted.

Thanking you for any courtesy,

I am, Very truly yours,

DAVID H. KLING, M.D.