DIFFERENTIAL AGGLUTINATION TITRE (D.A.T.) IN JUVENILE RHEUMATOID ARTHRITIS

BY

E. G. L. BYWATERS, MARY E. CARTER, AND F. E. T. SCOTT

*From the Rheumatism Research Unit, Medical Research Council,
Canadian Red Cross Memorial Hospital, Taplow*

This study concerns the titre of differential sheep cell agglutination (Rose-Waaler) in rheumatoid arthritis affecting children, its variability and its correlation with other clinical features, during a period of at least 4 years' observation.

Method

The technique employed from 1949-1956, which has been described by one of us (Scott, 1952), was a slight modification of the method originally proposed by Rose, Ragan, Pearce, and Lipman (1948). Since 1956, a further modification has been employed, as suggested by Greenbury (1957), using Perspex trays (M.R.C. pattern) in place of tubes and racks. This method will be described in detail elsewhere (Bywaters and Scott, unpublished). The results by the two methods are closely comparable, the newer method giving more consistent results.

The results are expressed as a differential titre (ratio) in both methods, and the lowest ratio accepted as positive in a series of doubling dilutions is 1 : 16. Slight changes in the sensitivity of the method have been found to occur because of variations in agglutinability of red cell suspensions from different sheep, amounting usually to not more than a one-tube difference. A single positive result occurring in the midst of a series of negative tests in one individual is therefore suspect and may well be due to technical factors, although in an attempt to control this, sera of known titre were included as a safeguard in each series of titrations.

Material

This series consists of 142 patients (85 females and 57 males) admitted to this Unit from its inception in 1947 to the end of 1953, and followed to December 31, 1957, giving, for the purpose of this study, a maximum follow-

up period of 10 years and a minimum of 4 years; since the D.A.T. test was not started here until January, 1949, the period of D.A.T. observation ranges from 4 to 8 years in all cases, except in two patients who died and were followed for 3 years only. The 142 cases were all under 16 years of age at the onset of their illness.

75 patients were first seen more than one year after the onset of disease (referred to as "late" cases) and 67 within one year of onset ("early" cases). Patients were seen after discharge as a rule at least yearly when blood was taken. The mean number of D.A.T.s done per patient in this study was 9.5 (ranging from 2 to 24).

The diagnostic criteria for admission to the series were as follows:

- (i) Polyarthritis affecting four or more joints, past or present, other diagnoses having been excluded.
- (ii) Less than four joints involved, but diagnosis confirmed by synovial biopsy. Four patients had three joints involved, six had two joints involved, and three had one joint involved. Twelve of these thirteen patients had biopsy confirmation; the thirteenth was a child with monarthritis who also had iritis and had been followed for 8 years.

These criteria included five patients with psoriasis, (four with negative D.A.T.s and one with one D.A.T. at 1 : 32), one patient with ulcerative colitis, and one with sacro-iliac involvement.

Disease activity was considered to be present if there was active joint involvement with two out of the four following:

- (1) Joint pain.
- (2) Joint warmth.
- (3) Joint limitation of movement.
- (4) Joint swelling.

Results

Table I shows the results taking all D.A.T.s done (Groups I-III) and Table II the results of only the first D.A.T. done (Groups IV and V). The distribution of sex within these groups and the mean duration of follow-up are also shown. The difference between the mean duration of follow-up of Group I "all negative" (5.2 yrs) and of Group III "two or more positive" (6.1 yrs) is statistically significant ($p = <0.01$), but this does not affect the conclusions drawn from subsequent Tables. The only possible inference is that a slightly greater number of patients might become sero-positive if followed for longer periods.

The intermediate Group II of 36 patients with only one positive D.A.T. (Table I) we have found to resemble more closely the negative group as regards age and sex incidence (see Tables III and IV), whether the lone positive result was 1 : 16 (21 cases)

or 1 : 32 or more (15 cases); 32 out of the 36 had between six and 21 estimations made, the remaining four having two, three, four, and four respectively. Altogether, in these 36 patients, the single positive result represented in most cases under 20 per cent. of all D.A.T.s done, whereas in the 28 cases with more than one positive result, the positive results represent 60 per cent. of all D.A.T.s done. We feel that some of these single positive results in the midst of a negative series might possibly be due to technical error.

Age Incidence.—Table III (opposite) shows that the proportion of cases in which the first D.A.T. was positive increased markedly with age at onset. A similar relationship to age at onset is seen in the positive cases (Group III), but not in the negative cases (Group I) or in the doubtful positive cases (Group II), although the difference between the positive and

TABLE I
RESULTS OF ALL TESTS DONE IN 142 CASES, BY SEX, MEAN DURATION OF FOLLOW-UP,
AND MEAN DURATION FROM ONSET TO TIME FIRST EXAMINED

Group	D.A.T. Results	Cases		Sex		First Examined		Mean Duration of Follow-up (yrs)	Mean Duration of Disease before First Examination (yrs)
		No.	Per cent.	Male	Female	"Late" > 1 yr from Onset	"Early" < 1 yr from Onset		
I	All Negative	78	54.9	28	50	38	40	5.2 { 5.3 (M) 5.1 (F)	2.4
II	One Positive Only	36	25.4	12	24	18	18	5.9 { 5.8 (M) 5.9 (F)	2.3
III	Two or More Positive	28	19.7	17	11	19	9	6.1 { 6.3 (M) 5.7 (F)	2.3
Total		142	—	57	85	75	67	5.7	2.3

* Difference between these two means = 0.9
Standard error of this difference = 0.27
 $\therefore t = 0.9/0.27 = 3.33$ ($p = <0.01$)

TABLE II
RESULTS OF FIRST D.A.T. IN 142 CASES, BY SEX, MEAN DURATION OF FOLLOW-UP,
AND MEAN DURATION FROM ONSET TO TIME FIRST EXAMINED

Group	First D.A.T. Results	Cases		Sex		First Examined		Mean Duration of Follow-up (yrs)	Mean Duration of Disease before First Examination (yrs)
		No.	Per cent.	Male	Female	"Late" > 1 yr from Onset	"Early" < 1 yr from Onset		
IV	Positive	19	13.4	11	8	11	8	5.9 { 5.7 (M) 6.1 (F)	2.2
V	Negative	123	86.6	46	77	64	59	5.5 { 5.6 (M) 5.4 (F)	2.4
Total		142	—	57	85	75	67	5.7	2.3

TABLE III
AGE INCIDENCE

Group		IV	I	II	III
Age at Onset (yrs)	Total No. of Cases	First D.A.T. Positive	D.A.T.s All Negative	Only One Positive D.A.T.	Two or More Positive D.A.T.s
1-4	43	1 (2%)	29 (67.4%)	10 (23.3%)	4 (9.3%)
5-9	43	6 (14%)	20 (46.4%)	14 (32.6%)	9 (21%)
10-16	56	12 (21.5%)	29 (51.6%)	12 (21.4%)	15 (27%)
	142	19 (13.4%)	78 (54.9%)	36 (25.4%)	28 (19.7%)
Mean Duration of Follow-up (yrs)		5.9	5.2	5.9	6.1

negative groups is just not significant ($0.1 > p > 0.05$) at the 5 per cent. level. Table III also shows the mean duration of follow-up for each of the groups. The positive cases (Group III) were followed slightly longer than the all-negative cases (Group I), which may have exposed them to a slightly greater risk of giving a positive result, but this would not have influenced those whose first D.A.T. was positive (Group IV), who were followed as long as the more highly-positive group and who also showed a tendency for more positive results in the older age groups.

Table IIIA shows the mean duration of follow-up

TABLE IIIA
AGE INCIDENCE AND MEAN DURATION OF DISEASE AND LENGTH OF FOLLOW-UP IN THREE GROUPS

Age at Onset (yrs)	Total No. of Cases	Mean Duration of Follow-up (yrs)	Mean Duration of Disease from Onset to End of Follow-up (yrs)
1-4	43	5.5	9.2
5-9	43	5.8	7.7
10-16	56	5.3	6.7

for each of the three age groups, 1-4, 5-9, and 10-16 yrs, and the mean duration of disease from onset. The oldest group, which was most frequently positive, was followed for a slightly shorter period than the younger groups, so that the significance of the trend for positives to occur in the older group is not altered by length of follow-up, nor by the duration of disease.

Sex Incidence.—In Table IV the observed numbers of positive and negative cases in each of the various groups have been compared with the expected numbers on the nul hypothesis, and the χ^2 values calculated.

The proportion of males to females is the same in Groups I and II, but in Groups IV and III males are more frequently D.A.T.-positive than females, although this only approaches statistical significance in Group III (χ^2 , Yates' modification = 3.36; $0.1 > p > 0.05$). Although the males were followed for 5.8 years (mean) and the females for 5.5 years (mean), this does not alter the conclusions stated above, as in Group III both sexes fell into the longest follow-up period (mean 6.1 yrs).

TABLE IV
SEX INCIDENCE

Group		IV	I	II	III
Sex	Total No. of Cases	First D.A.T. Positive	D.A.T.s All Negative	Only One D.A.T. Positive	Two or More D.A.T.s Positive
Male	57	11	28	13	16
Female	85	8	50	23	12
Total	142	19	78	36	28
		NS $\chi^2 = 2.08$	NS $\chi^2 = 0.93$	NS $\chi^2 = 0.14$	NS $\chi^2 = 3.36$

Variation of D.A.T. during Course of Disease.— This has been examined for each 2-year period of observation, in respect both of tests (Table V) and of persons (Table VI), in the group of 64 children with at least one positive D.A.T. during observation. Tables VA and VIA show the total series of 142 cases.

These four Tables show a tendency for the D.A.T. to become negative with duration of disease. This tendency was slightly more marked in the early cases. In both early and late cases some initially positive changed to negative and *vice versa*. Only nine of the 64 patients showed at least one positive result each year.

TABLE V
PROPORTION OF POSITIVE D.A.T.s TO TOTAL NUMBER DONE IN 2-YEAR PERIODS ON 64 PATIENTS WITH AT LEAST ONE POSITIVE D.A.T. DURING OBSERVATION

First Examined	No. of Cases	Years of Observation								Total D.A.T.s Done on 64 Patients
		1st and 2nd		3rd and 4th		5th and 6th		7th and 8th		
		No.	Per cent.	No.	Per cent.	No.	Per cent.	No.	Per cent.	
Early ..	27	38/175	21·7	13/78	16·7	5/44	11·4	1/21	4·8	318
Late ..	37	60/185	32·4	32/105	30·5	22/86	25·6	6/40	15	416
Total ..	64	98/360	27·2	45/183	24·6	27/130	20·8	7/61	11·5	734

TABLE VA
PROPORTION OF POSITIVE D.A.T.s TO TOTAL NUMBER DONE IN 2-YEAR PERIODS IN THE TOTAL SERIES OF 142 PATIENTS

First Examined	No. of Cases	Years of Observation								Total D.A.T.s Done on 142 Patients
		1st and 2nd		3rd and 4th		5th and 6th		7th and 8th		
		No.	Per cent.	No.	Per cent.	No.	Per cent.	No.	Per cent.	
Early ..	67	38/325	11·7	13/157	8·3	5/103	4·8	1/39	2·6	638
Late ..	75	60/336	17·9	32/207	15·5	22/129	17·0	6/52	11·5	711
Total ..	142	98/611	16·0	45/364	12·4	27/232	11·6	7/92	7·6	1,349

TABLE VI
PROPORTION OF PATIENTS WITH A POSITIVE D.A.T. IN EACH 2-YEAR PERIOD TO NUMBER OF PATIENTS TESTED AT LEAST ONCE IN EACH 2-YEAR PERIOD

First Examined	No. of Cases	Years of Observation							
		1st and 2nd		3rd and 4th		5th and 6th		7th and 8th	
		No.	Per cent.	No.	Per cent.	No.	Per cent.	No.	Per cent.
Early ..	27	20/27	74·1	7/25	28	5/21	23·8	1/9	11·1
Late ..	37	19/37	51·2	20/34	58·8	15/29	51·7	5/19	26·3
Total ..	64	39/64	61	27/59	45	20/50	40	6/28	21·5

TABLE VIA
PROPORTION OF PATIENTS WITH A POSITIVE D.A.T. IN EACH 2-YEAR PERIOD TO TOTAL NUMBER OF PATIENTS TESTED IN SERIES (142 CASES)

First Examined	No. of Cases	Years of Observation							
		1st and 2nd		3rd and 4th		5th and 6th		7th and 8th	
		No.	Per cent.	No.	Per cent.	No.	Per cent.	No.	Per cent.
Early ..	67	20/66	32	7/56	12·5	5/48	10·4	1/17	5·8
Late ..	75	19/75	25·2	20/64	31·2	15/50	30	5/26	19·3
Total ..	142	39/141	27·6	27/120	22·3	20/98	22	6/43	14

Correlation of D.A.T. with Disease Activity.—Table VII compares disease activity and changes in D.A.T. during the observation period in three groups: Group I “negative”, Group II “doubtful”, and Group III “more positive”. Each group is subdivided into early and late cases, but this subdivision shows little relevance.

The same proportion of sero-positive and sero-negative patients show activity which subsides, but a higher proportion of sero-positive patients remain active throughout their period of observation. None of the patients with a positive D.A.T. at some time were inactive throughout the observation period. Since those remaining active were followed for a slightly shorter period (mean 5·3 yrs) than those who became inactive (mean 5·7 yrs), it is reasonable to expect that the former group would eventually become inactive given more time.

Of the 64 patients who had a positive D.A.T. at some time, 35 had negative results in the first year and became positive later; 29 were positive at some time in the first year of observation and became negative later. Of the 35 cases in which the D.A.T.s were at first negative and later became positive, eleven were seen within the first year of their illness

(“early”) and information on the duration of “negativity” is therefore available. This averaged nearly 4 years (Table VIII) with a range from 2 to 7 years, nearly equal numbers changing over in each of the three age groups (0-9, 10-16, and 17-20). Ten out of eleven patients had active disease when the first D.A.T. (negative) was done and five out of eleven patients had active disease when the D.A.T. became positive.

The D.A.T. became negative between 0 and 8 years of age in three cases, between 9 and 16 years in eight cases, and between 17 and 20 years in three cases. All fourteen patients had active disease when the first D.A.T. (positive) was done and five of them still had active disease when the D.A.T. became negative. The mean duration of “positivity” in these fourteen cases was 2·3 years.

Table VIII and Table IX (overleaf) show that as many patients became negative as became positive in the older groups. These Tables also show that there is no strict correlation between D.A.T. “positivity” and disease activity. When the D.A.T. is negative, this is not unexpected, since a large number of patients had negative D.A.T.s throughout, despite disease activity at some time in 69 out of 78 cases.

TABLE VII
CORRELATION OF D.A.T. WITH DISEASE ACTIVITY

Group	No. of Cases	Results of D.A.T. during Observation	First Examined	Disease Activity							
				Inactive Becoming Active		Inactive Throughout		Active Becoming Inactive		Active Throughout	
				No.	Per cent.	No.	Per cent.	No.	Per cent.	No.	Per cent.
I	78	Negative	Early .. Late ..	3} 3}	6 7·7	6} 3}	9 11·5	29} 19}	48 61·5	1} 14}	15 19·2
II	36	Only One Positive	Early .. Late ..	1} 1}	2 6	0		14} 13}	27 75	3} 4}	7 19
III	28	Two or More Positive	Early .. Late ..	0} 1}	1 3·6	0		7} 8}	15 53·6	2} 10}	12 42·8
Total ..	142	—	—	9	—	9	—	90	—	34	—

TABLE VIII
DURATION OF NEGATIVE RESPONSE IN ELEVEN “EARLY” CASES WHO CONVERTED TO POSITIVE LATER

Age at Onset when D.A.T. was Negative (yrs)	Disease Activity when First Examined	Age when D.A.T. First Positive (yrs)	Disease Activity when D.A.T. First Positive	Time Interval before D.A.T. Positive for First Time (yrs)
1	—	3	+	2
5	+	7	—	2
5	+	9	+	4
5	+	7	—	2
5	+	12	+	7
6	+	8	—	2
8	+	11	—	3
9	+	12	—	3
12	+	18	—	6
14	+	20	+	6
15	+	19	+	4
11	—	—	—	Mean 3·7

TABLE IX
DURATION OF POSITIVE RESPONSE IN FOURTEEN "EARLY" CASES WHO CONVERTED TO NEGATIVE LATER

Age at Onset when D.A.T. was Positive (yrs)	Disease Activity when First Examined	Age when D.A.T. was First Negative (yrs)	Disease Activity when D.A.T. was First Negative	Time Interval before D.A.T. Negative for First Time (yrs)
1	+	3	-	2
2	+	5	-	3
6	+	8	-	2
7	+	9	-	2
9	+	11	+	2
10	+	12	-	2
11	+	15	-	4
12	+	15	+	3
12	+	14	+	2
13	+	15	+	2
14	+	16	+	2
15	+	17	-	2
15	+	18	-	3
15	+	17	-	2
14				Mean 2·3

Note: The remaining two "Early" cases out of the total 27 "Early" cases who had a positive D.A.T. at some time did not convert from positive to negative during their period of observation.

Figure.—Variation of titre in relation to disease activity in sixteen patients with Still's disease.

Ann Rheum Dis: first published as 10.1136/ard.18.3.225 on 1 September 1959. Downloaded from <http://ard.bmj.com/> on April 26, 2023 by guest. Protected by copyright.

The Figure shows the pattern of D.A.T. response in the sixteen patients who had two or more positive D.A.T.s at 1 : 32 or more. These patterns show that there is no particular trend in the behaviour of D.A.T. titres for individuals, and no two cases showed any similarity of pattern; this supports the statement made above that there is no strict correlation between D.A.T. positivity and disease activity. (The white dots in the Figure indicate the sequence of tests done, but are not related to the date when they were performed in each year.)

Although, as shown in Table III, serological positivity is more frequent in children whose disease starts in their 'teens, there seems to be little tendency for the group of children who are younger at the time of onset to convert to sero-positivity when they reach their 'teens. However, our period of observation is short and our numbers few. This is an important point which will be decided by further observation.

We have also tried to correlate the D.A.T. in these children with other clinical features, such as rash, nodules, erosions, lymphadenopathy, and splenomegaly (Table X). Although splenomegaly was

found more commonly in the positive groups to a significant degree ($0.05 > p > 0.02$), many comparisons are being made using the same data and a higher level of significance is needed.

If, however, we select the 28 highly positive cases (Group III: two or more positive D.A.T.s during observation) for comparison with the rest (Table XI), a greater probability emerges that nodules are related to D.A.T. positivity ($p = > 0.001$), as is known to be the case in adult rheumatoid arthritis (Ball, 1952).

Summary

142 patients suffering from Still's disease were followed for from 4 to 8 years; 13.4 per cent. had a positive D.A.T. (1 : 16 or more) at the first test, and 19.7 per cent. had two or more positive D.A.T.s. The mean number of D.A.T.s performed on each patient was 9.5.

Sero-positivity was more frequent in those patients who were older at the time of onset, but there was little tendency for those who were younger at onset to convert to sero-positivity when they reached their 'teens. Males tended to be more frequently sero-positive than females.

TABLE X
CORRELATION OF D.A.T. WITH OTHER CLINICAL FEATURES

Group	D.A.T. Results	Total No. of Cases	Clinical Features									
			Rash		Nodules		Splenomegaly		Lymphadenopathy		Erosions	
			No.	Percent.	No.	Percent.	No.	Percent.	No.	Percent.	No.	Percent.
III	Two or More Positive	28	6	22	7	25	8	29	18	64	14	50
II	One Only Positive	36	10	28	0	0	13	36	14	39	12	33
I	All Negative	78	21	27	4	5	13	17	35	45	22	56
Total		142	37	—	11	—	34	—	77	—	48	—

TABLE XI
CORRELATION OF D.A.T. WITH OTHER CLINICAL FEATURES

Group	D.A.T. Results	Total No. of Cases	Clinical Features									
			Rash		Nodules		Splenomegaly		Lymphadenopathy		Erosions	
			No.	Percent.	No.	Percent.	No.	Percent.	No.	Percent.	No.	Percent.
III	Two or More Positive	28	6	21.8	7	25	8	28.6	18	64.2	14	50
I and II	All Negative or Only One Positive	114	31	27.2	4	3.5	26	22.8	49	42.9	34	29.8
Total		142	37	—	11	—	34	—	77	—	48	—
			$\chi^2 = 0.15$		$\chi^2 = 11.6$		$\chi^2 = 0.1$		$\chi^2 = 3.2$		$\chi^2 = 3.2$	
			NS		$P = < 0.001$		NS		$P = 0.1-0.05$		$P = 0.1-0.05$	

There was no typical pattern of D.A.T. response throughout the period of follow-up; there was no correlation with duration of disease and little correlation with disease activity.

An attempt was made to correlate the D.A.T. with other clinical features, such as rash, nodules, erosions, lymphadenopathy, and splenomegaly. When the highly-positive group (two or more positive D.A.T.s) was compared with the remainder, nodules appeared to be related to sero-positivity, as in adult patients.

REFERENCES

- Ball, J. (1952). *Ann. rheum. Dis.*, 11, 97.
 Bywaters, E. G. L., and Scott, F. E. T. In "Recent Advances in Clinical Pathology," ed. S. C. Dyke. Unpublished.
 Greenbury, C. L. (1957). *Ass. clin. Path.*, Broadsheet No. 18.
 Rose, H. M., Ragan, C., Pearce, E., and Lipman, M. O. (1948). *Proc. Soc. exp. Biol. (N.Y.)*, 68, 1.
 Scott, F. E. T. (1952). *Lancet* 1, 392.

Titre d'agglutination différentielle (D.A.T.) dans l'arthrite rhumatismale juvénile

RÉSUMÉ

On observa 142 cas de maladie de Still pendant 4 à 8 ans; 13,4% de ces cas accusèrent un D.A.T. positif (1 : 16 ou plus) à la première détermination et 19,7% eurent deux ou plus titres positifs. En moyenne, le D.A.T. fut effectué 9,5 fois chez chaque malade.

La séro-positivité fut plus fréquente chez des enfants plus âgés au moment du début de la maladie, mais il n'y eut qu'une tendance faible chez des enfants à début précoce à devenir séro-positifs après avoir dépassé l'âge de 10 ans. Les garçons tendaient à être séro-positifs plus souvent que les filles.

Pendant toute la période d'observation on ne vit aucun tableau typique de D.T.A.; il n'y eut pas de corrélation avec la durée de la maladie et peu de corrélation avec son évolution.

On essaya de mettre en corrélation le D.A.T. et les autres caractères cliniques, tels que nodules, exanthème, érosions, lymphadenopathie et splénomégalie. En comparant le groupe fortement positif (deux ou plus D.A.T. positifs) avec le reste, on trouva, comme chez des adultes, une plus grande fréquence des nodules dans ce groupe.

Cifra de aglutinación diferencial (D.A.T.) en la artritis reumatoide juvenil

SUMARIO

Se observaron 142 casos de enfermedad de Still durante 4 a 8 años; en un 13.4% ellos la reacción de aglutinación diferencial fué positiva (1 : 16 o más) en la primera determinación y en un 19.7% se obtuvo dos o más reacciones positivas. En promedio se procedió a 9,5 determinaciones de la D.A.T. en cada enfermo.

La sero-positividad fué más frecuente en niños de mayor edad al tiempo del comienzo de la enfermedad, pero niños enfermos desde la edad más tierna tuvieron poca tendencia a convertirse en sero-positivos después de rebasar la edad de 10 años. Los chicos tendían a presentar reacciones sero-positivas más frecuentemente que las chicas.

Durante el entero período de observación no se dibujó ningún cuadro típico de la D.A.T.; tampoco se observó correlación alguna con la duración de la enfermedad, ni mucha con su evolución.

Se tentó correlacionar la D.A.T. con otros rasgos clínicos, como exantema, nódulos, erosiones, linfadenopatía y esplenomegalia. Al comparar el grupo fuertemente positivo (dos o más D.A.T. positivas) con el resto, se vió, como en enfermos adultos, una relación entre nódulos y sero-positividad.