Hepatic effects of fenclozic acid

F. DUDLEY HART, LOGIE S. BAIN, E. C. HUSKISSON, T. R. LITTLER, AND R. T. TAYLOR

From the Westminster Hospital, London, S.W.1, the City Hospital, Aberdeen, and the General Hospital, Birkenhead

Fenclozic acid (I.C.I. 54,450: Myalex) has recently been withdrawn from trial because of hepatotoxicity. In a published trial (Chalmers, Kellgren, and Platt, 1969), which suggests that it has some effect as an anti-inflammatory agent in rheumatoid arthritis, up to 400 mg. fenclozic acid were given daily for 10 days to 40 patients and compared with 3.6 g. aspirin daily. Two of these patients showed abnormal serum glutamic-oxaloacetic transaminase (SGOT) values after treatment with fenclozic acid. As more patients had raised SGOT values after treatment with aspirin it was concluded that further evaluation of fenclozic acid was justified.

Present investigation

We have recently completed trials of fenclozic acid in rheumatoid arthritis. Eight patients developed jaundice or abnormalities of hepatic function, as

detailed in the Table. As these investigations were performed in different laboratories, the normal values for each laboratory are also given. The four patients who became jaundiced had received fenclozic acid for 17 to 35 days.

Comment

As there was an unacceptable incidence of jaundice and other abnormalities of liver function in patients who had had fenclozic acid for more than 14 days, the drug was withdrawn. In short trials of drugs of this nature serum enzyme levels appear to be unsatisfactory for predicting or excluding longer-term liver toxicity.

Summary

The effects of fenclozic acid on liver function are described. These have lead to withdrawal of the drug

Table *Liver function tests*

<i>Case no.</i>	<i>Bilirubin (mg./100 ml.)</i>	<i>Alkaline Phosphatase (KA units)</i>	<i>SGPT (units)</i>	<i>SGOT (units)</i>
Normal (Aberdeen)	< 1.0	< 13	< 40	< 35
1*	2.0	50	160	81
2*	4.0	39	170	64
3	1.2	26	53	70
4	0.2	10	100	187
5	0.2	16	63	70
6	0.2	13	83	70
Normal (Birkenhead)	< 1.0	< 10	< 19	< 19
7*	2.9	40	17	46
Normal (Westminster)	< 1.0	< 17	< 12	< 17
8*	8.5	30	35	40

*Jaundiced

Reference

CHALMERS, T. M., KELLGREN, J. H., AND PLATT, D. S. (1969) *Ann. rheum. Dis.*, **28**, 595 (Evaluation in man of fenclozic acid (I.C.I. 54,450: Myalex), a new anti-inflammatory agent.II. Clinical trial in patients with rheumatoid arthritis).