The dosage of radioactivity received by the technician in the course of preparation of the isotope, handling syringes, and initial nursing would be approximately the same.

Now the maximum permissible dosage for the hands and forearms is 15 rem. in 3 months, or about 1,200 m.rem in a week. The maximum permissible irradiation for a worker in this field is 6,200 m.rem. These data show that the amount of radioactivity absorbed by the medical staff is well within the safety margin, and will help to deter-

mine how many patients can safely be treated each week.

DR. ANSELL I think we have had an extremely interesting day. We have learned that there is still an enormous field for further work in the use of these isotopes—to find the best type of isotope for each joint and to carry out controlled studies of the use of the many techniques that are now available to us. I am sure that further excellent work will be done and that we shall have to re-convene in a year or two's time to see how we are all getting on.

Afterthoughts

J. M. GUMPEL

It is clear from the discussion that there is general acceptance of the efficacy of radiocolloids in the treatment of chronic synovitis, and general agreement about the indications for treatment and the selection of patients. Although one suitable controlled study has confirmed the beneficial use of ^{90}Y in the knee, there is still a need for further controlled studies of the treatment of other joints, and a need to clarify the effect of certain other factors such as the previous removal of fibrin from the joint.

Interest is at present focused on two main points: the possibility of long-term hazards of treatment, and the leakage of radio-pharmaceuticals from the joint. These two points are linked, as the long-term hazards seem to be related more to whole body irradiation than to irradiation of the synovium. As a

result, the measurement of the retention of radiocolloids in the joint becomes a valuable research procedure rather than an academic exercise, and should certainly replace the measurement of extra-articular spread with its potential 20 per cent. "dead space". The adoption of any measure that will reduce irradiation of other structures is important, and amongst these are the avoidance of radiocolloids with significant γ emission, the evaluation of the available radiocolloids to compare their relative retention and common patterns of spread, and probably most important of all the investigation of the variable permeability of the individual synovial membrane, and the application of methods of reducing transport across the membrane, as for example by rest and by premedication.