Correspondence

Sir,

The now common use of penicillamine in the treatment of rheumatoid arthritis makes it important to minimize the risk of serious adverse reaction to the drug. A few severe and/or fatal blood dyscrasias have already occurred.

I am collecting information on patients who have developed aplastic anaemia, leucopenia, granulocytopenia, or thrombocytopenia attributable to penicillamine. I would be most grateful to your readers for details of any such patients under their care.

APRIL KAY

Medical Officer, ARC Epidemiology Research Unit, New Cross Hospital, Avonley Road, London SE14 5ER.

Notes

Workshop on Immune Complexes

Kennedy Institute of Rheumatology, London, January 9, 1976

The workshop was sponsored by the Arthritis and Rheumatism Council. Two sessions were held, on the detection and measurement of circulating soluble immune complexes, and on the detection and measurement of anti-DNA antibodies. The proceedings will be published, edited by Dr. R. N. Maini and Dr. E. J. Holborow, as a supplement to the *Annals*.

Symposium on the Fibrotic Process

Royal Postgraduate Medical School, London, April 23, 1976

The symposium was organized by Professor E. G. L. Bywaters and Dr. M. I. V. Jayson for the Arthritis and Rheumatism Council. Four sessions were held, on the development of fibrosis, fibrosis in disease, chemical fibrosis and immunological reaction, and treatment of fibrosis. The proceedings will be published, edited by Dr. M. I. V. Jayson, as a supplement to the *Annals*.