III. Chemical fibrosis and immunological reaction

CHAIRMAN: L. E. GLYNN

Practolol peritonitis

A. E. READ

From the Department of Medicine, University of Bristol

The inflammatory process in the peritoneum related to the administration of practolol is characterized by dense fibrosis. First described by Brown *et al.* (1975), the disease will disappear with the introduction of restrictions on the prescription of the drug. However, the involvement of the peritoneal membrane in drug reactions is unusual and several clinical and pathological features of the disorder are worthy of review.

The disease is one of the middle-aged and elderly—that is, those who have cardiac disorders, usually myocardial ischaemia, that have been treated with the drug. It is found equally in the two sexes, and in the 16 patients that we have treated or reviewed the symptoms have usually been of small intestinal obstruction. Thus vomiting and weight loss, sometimes considerable, have been seen in 15 of them, while one has presented with malabsorption associated with small intestinal bacterial contamination.

In all except four patients these symptoms were accompanied by the presence of one or more abdominal masses. These have been in various sites in the abdomen and have mimicked a variety of neoplastic and other disorders. For example, the women with lower abdominal masses had been invariably referred for gynaecological operations (Trudinger and Fitchett, 1976). Many investigations were performed on these patients, particularly before the nature of the disorder was recognized, but the most helpful information was obtained by barium examination of the small bowel. Here dilatation of the small bowel starting in the duodenum, constricting and fixation of small bowel loops, delay in small bowel transit, and sacculization of some of

the small bowel, which appeared almost like normal barium-filled colon, were all seen. Sometimes the separation of loops of barium-filled small bowel indicated undue peritoneal thickening.

All but one of the patients has been subject to one or more laparotomies. The findings have usually been identical. The visceral peritoneum is grossly thickened and loops of gut are densely adherent, and in some cases the whole of the small bowel is cocooned in a fibrous bag which encircles the whole of the small bowel from duodenum to ileum, sparing the colon. There is no involvement of the retroperitoneal tissues, thus distinguishing this disorder from retroperitoneal fibrosis. Surgical removal of the thickened peritoneum allows the constricted small bowel to assume its normal length.

Histological examination of resected tissue confirms the dense fibrosis (Fig) and shows a mild underlying inflammatory process. Examination of the tissue biochemically has confirmed the presence of increased cross-links of a stable type.

Patients treated surgically have all eventually obtained relief from chronic intestinal obstruction though two patients died after laparotomy, one after the first operation, the second after the third. Postoperative complications such as further myocardial infarction, pulmonary embolus, and wound dehiscence were also seen, while other problems of practolol toxicity such as ocular involvement made general progress slow.

Skin (9 cases) and ocular (11 cases) involvement were common in this group of 16 patients, while pleural involvement, sometimes with effusion, was

FIGURE Superficial peritoneal fibrosis with vascular adipose tissue and low-grade inflammation. Haematoxylin and eosin. $\times 22$

seen in three cases. Parenchymatous pulmonary fibrosis was also seen in two other patients.

Practolol peritonitis was seen between one and four years after the start of practolol therapy in the

usual therapeutic dosage. At first it certainly proved to be a difficult disease to diagnose. It is an unpleasant and dangerous condition and one whose precise cause is speculative.