Encore

Interleukin 1 and cartilage metabolism

Interleukin 1 in bovine cartilage culture leads to prostaglandin E₂ release and stimulation of proteoglycan breakdown. This effect is much more potent in inhibiting synthesis of proteoglycan than in breaking it down however, and the effect on proteoglycan occurred earlier than that on prostaglandin release. Non-steroidal anti-inflammatory drugs seemed to have little effect on proteoglycan metabolism, but other drugs such as chloroquine did appear to inhibit proteoglycan synthesis. The effects of interleukin 1 on proteoglycan breakdown, its synthesis, and prostaglandin release seem to be independently regulated. *Arthritis Rheum* 1989; 32: 288-97.

Raynaud phenomenon improved by the cold

A letter in the *Lancet* reports a smoker who used vibratory machinery whose Raynaud phenomenon improved with the cold. Cooling his hand increased blood flow by 30%. The authors are intrigued but baffled. Has anyone else seen this, or can throw light on it? *Lancet* 1989; i: 905.

Pregnancy and systemic lupus erythematosus (SLE)

Young women with SLE want to know whether it is safe to become pregnant and whether harm will result to their babies. A study on 80 pregnant women with this condition suggests that worsening of the disease is uncommon and that prophylactic prednisolone treatment is unnecessary. *Arthritis Rheum* 1989; 32: 665-70.

Cranial subluxation of the odontoid peg in rheumatoid arthritis

When cranial subluxation of the odontoid peg occurs it intrudes progressively into the foramen magnum and may even be fractured. Compression of the medulla oblongata and a subsequent progressive myelopathy is likely to occur. Bilateral collapse of the lateral facet joints as well as ligamentous weakening seems to be the cause. A study of 18 affected patients shows that surgical stabilisation arrests progress of the lesion and in about three quarters of the patients there is functional improvement. This is not the case with conservative management.

J Bone Joint Surg [Am] 1989; 71: 189-95.

Osteoporosis and pulsed electromagnetic fields

The use of electromagnetic fields to regulate bone mass and induce healing of bone fractures has been considered even since 1954 and with mixed enthusiasm. A study on turkey ulnae, however, showed a demonstrable osteogenic response, though the frequency and power of these pulses seemed to be critical. Shall we be seeing a clinical application of this observation in osteoporosis in future years? Time will tell. This form of treatment has had a chequered career so far.

J Bone Joint Surg [Am] 1989; 71: 411-7.

Cyclosporin and gout

A study in renal transplant recipients treated with cyclosporin showed a significant increase in hyperuricaemia and clinical gout compared with those treated with azathioprine. The concomitant use of diuretics exaggerated this effect. It seemed to be due to a reduction in renal urate clearance. Gouty arthritis seems to be a real problem in patients with a renal transplant receiving this drug.

N Engl J Med 1989; 321: 287-92

Cartilage collagens

There is a useful review of collagen types in articular disease in a recent issue of *Arthritis and Rheumatism*. Thirteen different collagens have now been identified, but a structural function for some of them is not obvious as yet. The article is a helpful source of recent references on the subject and attempts, briefly, to introduce some daylight into an ever more complex situation.

Arthritis Rheum 1989; 32: 241-6.

Total hip replacement, warfarin, and pulmonary embolism

The risk of a postoperative pulmonary embolus in a patient undergoing total hip replacement is an ever present worry. A study of 3000 patients undergoing this operation in Los Angeles, who had warfarin prophylaxis from the night of operation onwards, showed only a 0.5% rate with no fatalities. Bleeding occurred in 1.5%. The study was uncontrolled, but these results are enviably good when compared with rates as high as 5.2% in other series where anticoagulants were not used.

J Bone Joint Surg [Am] 1989; 71: 321-6.

Negative studies

A leader in the *New England Journal of Medicine* considers the problem. Negative studies it is believed are less likely to be published than positive ones, thereby distorting scientific reports. The failure to publish these negative results may be due more to recognition of basic failures in the work by the authors than a bias against publishing negative reports. Another reason for not publishing negative results is that too often the hypothesis tested is not very important. Good negative studies of important issues should be published, nevertheless. What do our readers think?

N Engl J Med 1989; 321: 464-6.

Stabilisation of the wrist in rheumatoid disease

The unstable subluxed wrist is a major handicap to patients with rheumatoid arthritis, though appropriate splinting is often a practical answer. Arthrodesis may be a lengthy procedure and tiresome to the patient. This paper describes a method of internal fixation with an intramedullary Rush nail. The procedure requires no external immobilisation apparently and is simple, long lasting, and looks to be a useful operation where destructive changes are advanced.

J Hand Surg [Br] 1989; 14: 288-90.

GEMAENNES