Whereas IL-1ra can completely prevent the latter process,⁹ blocking activation of the mature osteoclast requires simultaneous inhibition of TNF and IL-1.¹⁰ It would therefore be logical to contemplate clinical trials in RA, inhibiting both of these cytokines and observing the effect on RA related osteoporosis as an end point.

GILES V CAMPION
Synergen BV
PO Box 85515
2508 CE Den Haag
The Netherlands

- 1 Gowen M, Wood D D, Ihrie E J, McGuire M K B, Russell R G G. An interleukin-1-like factor stimulates bone resorption in vitro. *Nature* 1983; **306**: 378–80.
- 2 Boyce B F, Aufdemorte T B, Garrett I R, Yates A J P, Mundy B R. Effects of interleukin-1 on bone turnover in normal mice. *Endocrinology* 1989; **125**: 1142–50.
- 3 Pfeilschifter J, Chenn C, Bird A, Mundy G R, Roodman G D. Interleukin-1 and tumor necrosis factor stimulate the formation of human osteoclast-like cells in vitro. *J Bone Miner Res* 1989; **4**: 113–8.
- 4 Pacifici R, Rifas R, McCracken I, et al. Ovarian steroid treatment blocks a postmenopausal increase in blood monocyte interleukin-1 release. *Proc Natl Acad Sci USA* 1989; **86**: 2398–402.
- 5 Pacifici R, Rifas L, Teitelbaum S, et al. Spontaneous release of interleukin 1 from human blood monocytes reflects bone formation in idiopathic osteoporosis. *Proc Natl Acad Sci USA* 1987; **84**: 4616–20.
- 6 Pacifici R, Vannice J L, Rifas L, Kimble R B. Monocytic secretion of interleukin-1 receptor antagonist in normal and osteoporotic women: effects of menopause and estrogen/progesterone therapy. *J Clin Endocrinol Metab* 1993; **77**: 1135–41.
- 7 Firestein G S, Boyle D L, Yu C, et al. Synovial interleukin 1 receptor antagonist and interleukin 1 balance in rheumatoid arthritis. *Arthritis Rheum* 1994; **37**: 644–52.
- 8 Kitazawa R, Kimble R B, Vannice J L, Kung V T, Pacifici R. Interleukin-1 receptor antagonist and tumor necrosis factor binding protein decrease osteoclast formation and bone resorption in ovariectomized mice. *J Clin Invest* 1994. In press.
- 9 Kimble R B, Vannice J L, Bloedow D C, Thompson R C, Pacifici R. Interleukin-1 receptor antagonist decreases bone loss and bone resorption in ovariectomized rats. *J Clin Invest* 1994; **93**: 1959–67.
- 10 Kimble R B, Matoyoshi A B, Vannice J L, Kung V, Pacifici R. Simultaneous inhibition of TNF and IL-1 is required to block bone loss in the early postovariectomy period. *J Bone Miner Res* 1994; **9** (suppl): Abstract 153.

Antibodies to nuclear antigens in three patients with scleroderma and asthma

We read with interest the report of Martin *et al*¹ regarding autoantibodies to centromere and histone in patients with scleroderma and severe pulmonary and vascular disease. During the past two years we have been following up three female patients with scleroderma (two with localised scleroderma and one with diffuse scleroderma) and asthma, aged 40, 48, and 77, respectively; the onset of scleroderma had been preceded by asthma in each of them. The first two patients were atopic. Accompanying diseases were acne rosacea (first patient), thyroid adenoma (second), and diabetes mellitus type II (third patient). The diabetes in the last patient had begun 30 years ago after stress. Symptoms typical of asthma and radiographic evidence of pulmonary emphysema were found in all patients; none had symptoms suggestive of Sjögren's syndrome. The serum IgE values

were increased in the first two patients. All patients had positive antinuclear antibodies (ANA) detected by indirect immunofluorescence on HEp-2 cells in titre 1:80 with a speckled pattern of immunofluorescence. The second patient had positive DNA antibodies (enzyme linked immunosorbent assay (ELISA)), and the second and third patients positive antibodies to histone 2A (ELISA). None of the three had Ro(SS-A) or La(SS-B) antibodies. The son of the second patient also has asthma.

These cases cannot be included in the group of any overlap syndromes. The question is whether asthma is a manifestation of fibrosis, or the two diseases are independent entities. The scleroderma involves only the skin in two of the patients. Spirometry showed an obstructive defect of ventilation in all three patients. The increase in IgE in the sera of two patients suggests involvement of reaginic type reactions. The ANA positivity suggests type II and III immune reactions. We could find no published reports of a combination of scleroderma, asthma, and positive ANA. The presence of two diseases—scleroderma and asthma—raises the following issues:

- (1) The role of scleroderma as a cause for the development of asthma in these patients. The pulmonary involvement in scleroderma is very common. It is of a restrictive type, while in asthma, obstruction of the airways is a characteristic feature. Our data support the notion of an independent development of asthma.
- (2) The role of the endocrine system in the development of scleroderma. It is well known that rheumatic diseases are more frequent in females. Females also suffer from rosacea more often and the appearance of the disease may be linked to the climacteric. It is notable that two of our patients had accompanying endocrine diseases, thyroid adenoma and diabetes. The combination of asthma with scleroderma and endocrine disorders in our three patients warrants further investigation of this association.

K NIKOLOV

Department of Dermatology, Faculty of Medicine
1 G Sofiiski str Sofia 1431 PO Box 45
Bulgaria

M BALEVA

T VULCHEVA

Centre of Allergy

P JAKOVLEV

Department of Internal Medicine, St Zagora

- 1 Martin L, Pauls J D, Ryan J P, Fritzler M J. Identification of a subset of patients with scleroderma with severe pulmonary and vascular disease by the presence of autoantibodies to centromere and histone. *Ann Rheum Dis* 1993; **52**: 780–4.

AUTHOR'S REPLY: Dr Nikolov and his colleagues raise interesting questions. The three patients they describe had asthma diagnosed on clinical grounds and confirmed by spirometry. We also are not aware of any reports of asthma co-existing with scleroderma, but as asthma occurs in 4–6% of the population,¹ it is not surprising that the two diseases coexist. However, as Nikolov *et al* imply, a key question is whether scleroderma and asthma may have a common pathogenic basis.

In addition to the observations offered in their letter, it is interesting that scleroderma and asthma do share certain pathogenic features; for example increased dermal mast cells¹ and increased ability of mast cells to

release histamine² have been reported in scleroderma. Both these findings suggest that the pathogenesis of scleroderma may include a hypersensitive reaction similar to that described in asthma.¹ It is not clear if a similar process involving mast cells occurs in the lung in scleroderma, although circumstantial evidence suggests that it does.

Our report⁴ suggested that scleroderma patients with centromere and histone antibodies had more severe disease characterised by pulmonary and vascular involvement. It is not clear if the patients referred to by Nikolov *et al* had centromere antibodies, but two did have antibodies to histone H2a. Although review of our charts has not identified asthma as a clinical feature in our patients, based on the observations of Nikolov *et al*, in future studies it may be important to determine if there are clinical correlations between the two diseases.

M J FRITZLER

Faculty of Medicine

The University of Calgary

3330 Hospital Dr NW

Calgary, AB Canada T2N 4N1

- 1 Hawkins R A, Claman H N, Clark R A, Steigerwald J C. Increased dermal mast cell populations in progressive systemic sclerosis: a link to chronic fibrosis? *Ann Intern Med* 1985; **102**: 182–6.
- 2 Pearson M E, Huff J C, Giorno R C, Panicheewa S, Claman H N, Steigerwald J C. Immunologic dysfunction in scleroderma: evidence for increased mast cell releasability and HLA-DR positivity in the dermis. *Arthritis Rheum* 1988; **31**: 672–7.
- 3 Bigby T D, Wasserman S I. Asthma. In: Stein J H, ed. *Internal medicine*, 4th edn, ch. 300. Chicago: Mosby, 1994: 2387–98.
- 4 Martin L, Pauls J D, Ryan J P, Fritzler M J. Identification of a subset of patients with scleroderma and severe pulmonary and vascular disease by the presence of antibodies to centromere and histone. *Ann Rheum Dis* 1993; **52**: 780–4.

LETTERS TO THE EDITOR

Ehlers Danlos syndrome and osteoporosis

Ehlers Danlos syndrome (EDS) is a group of inherited connective tissue disorders with extreme genetic and clinical variability.¹ The clinical manifestations of EDS are a result of abnormalities in collagen types 1 and 3, the major proteins of skin, ligaments, tendons, blood vessels, and internal viscera. Type 1 collagen is also the main protein constituent of the bone matrix and its abnormalities form the molecular basis of osteogenesis imperfecta (OI).² EDS is therefore quite closely linked to OI and the two conditions are known to coexist.^{3 4} In spite of this there are no detailed studies available on bone mineral content in patients with EDS.

In the past three years we have seen seven patients with EDS and assessed bone densities in the lumbar spine and hip using dual energy x ray absorptiometry.

A 65 year old lady was referred from the orthopaedic department for investigations of a wedge fracture of the L2 vertebra. She did not have any obvious precipitating causes for osteoporosis in her past medical, menstrual,