Hydrostatic pressure induces expression of interleukin 6 and tumour necrosis factor α mRNAs in a chondrocyte-like cell line

Kenji Takahashi, Toshikazu Kubo, Yuji Arai, Isao Kitajima, Masaharu Takigawa, Jiro Imanishi, Yasusuke Hirasawa

Abstract

Objective—To clarify the effect of pressure on the expressions of proteoglycan core protein and metabolism related cytokines in a chondrocyte-like cell line, HCS-2/8.

Methods—HCS-2/8 cells were exposed to 1, 5, 10, or 50 MPa of hydrostatic pressure (HP) for two hours, and mRNA expressions of interleukin 6 (IL6) and tumour necrosis factor α (TNF α) were examined by using reverse transcription-polymerase chain reaction (RT-PCR) method with specific primer sets; and mRNA of proteoglycan core protein, stromelysin, and tissue inhibitor of metalloproteinase 1 (TIMP1) were measured with northern blotting.

Results—HP exposure caused temporal morphological changes of the cells, but did not affect cellular viability. IL6 and TNF α mRNA expressions were not observed in the control cells under the atmospheric pressure, whereas in the cells treated with HP, pressure dependent enhancement of IL6 mRNA expression was observed between 30 minutes and four hours after the HP release. TNF α mRNA expression also increased 30 minutes after the exposure to 50 MPa of HP and disappeared four hours later. Proteoglycan core protein mRNA levels increased between 30 minutes and four hours after the exposure to 1 or 5 MPa of HP, whereas the levels decreased after 10 or 50 MPa of HP. Stromelysin and TIMP1 mRNA signals did not respond to HP.

Conclusions—HP at excessively high levels induced IL6 and TNF α expression and reduced the expression of proteoglycan core protein, while physiological levels of HP increased the expression of proteoglycan core protein. These findings are important when considering the pathology of osteoarthritis.

(Ann Rheum Dis 1998;57:231-236)

could occur. This process of disease progression is confirmed by the fact that (a) clinical deformation in OA starts from the weight bearing area,¹ and (b) many OA related animal models are prepared by making mechanical changes on the joints.²⁻³ Chondrocytes play an important part in the metabolism and maintenance of cartilage matrix, therefore, it is quite important to elucidate how chondrocytes respond to pressure.

Recently, as a cause of onset and progression of OA, locally produced humoral factors, for example, cytokines and proteases, have been considered important, in addition to mechanical stresses. Among them, tumour necrosis factor α (TNF α) and interleukin 6 (IL6) were reported to be upregulated in cartilage of an experimental OA model or OA patients.⁴⁻⁵ TNF α promote connective tissue degradation through the inhibition of extracellular matrix synthesis, and induces expression of collagenase and stromelysin in chondrocytes, as well as interleukin 1 β .⁶ IL6 plays a prominent part in the coordinated systemic host defence response to injury. IL6 production in chondrocytes markedly increases when IL1 β , TNF α , or transforming growth factor β 1 (TGF β 1) are present⁷; and IL6 induces tissue inhibitor of metalloproteinase1 (TIMP1).⁸

However, to our knowledge, the pathology of OA has not been investigated in terms of the relation between pressure and cytokine expression. This prompted us to examine whether mechanical stress changed non-matrix chondrocyte expression, such as expression of OA related cytokines.

This study investigated the effect of pressure in mRNA expression of proteoglycan core protein, cytokines such as TNF α and IL6, which are implicated in OA,⁹ stromelysin, and TIMP1, by using a chondrocyte-like cell line, HCS-2/8,¹⁰⁻¹¹ which can be subcultured and possesses characteristic features of normal chondrocytes.

Department of
Orthopaedic Surgery,
Kyoto Prefectural
University of
Medicine,
Kawaramachi-Hirokoji,
Kamigyo-ku, Kyoto
602-0841, Japan
K Takahashi
T Kubo
Y Arai
Y Hirasawa

Department of
Laboratory and
Molecular Medicine,
Kagoshima University
School of Medicine,
Kagoshima Japan
I Kitajima

Department of
Biochemistry and
Molecular Dentistry,
Okayama University
Dental School,
Okayama, Japan
M Takigawa

Department of
Microbiology, Kyoto
Prefectural University
of Medicine, Kyoto
Japan
J Imanishi

Correspondence to:
Dr Kubo.

Accepted for publication
27 February 1998

Articular cartilage is an important component of the joint, and it is always exposed to pressure produced by weight bearing and muscle contraction. Cartilage matrix mainly consists of proteoglycan and collagen, and this realises a cushioning effect of the matrix against mechanical stresses. However, if the mechanical stresses are increased to a level excessively higher than the physiological levels, cartilage matrix will be impaired and osteoarthritis (OA)

Table 1 Cytokine specific oligonucleotide primers for PCR

Cytokine	Primer	Position	cDNA	Reference
IL6	S	242-261	819	15
	AS	4535-4554		
TNF α	S	1521-1540	468	16
	AS	2270-2289		
β actin	S	103-122	540	17
	AS	619-642		

S: sense. AS: antisense.

Figure 1 Morphological changes of HCS-2/8 after exposure to HP. Phase contrast photomicrograms of cells cultured in DMEM containing 10% FBS after HP exposure. (A) Control cells under atmospheric pressure. (B) Thirty minutes after applying 50 MPa of HP. (C) Twenty-four hours after applying 50 MPa of HP. (D) Thirty minutes after applying 5 MPa of HP. Bar = 100 μ m.

Figure 2 Pressure dependent enhancement of IL6 mRNA expression. (A) IL6 RT-PCR products from the cells at 30 minutes (a) and (b) and four hours (c) and (d) after HP exposure. Two point five microgram (a) and (c) and 1 μ g (b) and (d) of total RNA were reverse transcribed in 10 μ l reaction mixture. (B) β actin RT-PCR products from the cells at 30 minutes (a) and two hours (b), which confirm cDNA isolation. One microgram of total RNA was reverse transcribed in 10 μ l reaction mixture. C: control cells treated under the atmospheric pressure.

Methods

CELL CULTURE

HCS-2/8 cells were seeded in 60 mm plastic Petri dishes (Falcon Plastic Corp, Oxnard, CA), at a density of 2.0×10^6 cells/3 ml of Dulbecco's modified Eagle's medium (DMEM: Nissui Pharmaceutical Co, Tokyo, Japan) containing 10% fetal bovine serum (FBS: GIBCO BRL Co, Gaithersburg, MD), 60 μ g/ml of kanamycin, and 0.292 mg/ml of L-glutamine (Wako Pure Chemical Industries Ltd, Osaka, Japan). The cells were maintained at 37°C in a humidified atmosphere of 5% CO₂. The medium was changed twice a week.

PRESSURE APPLICATION

The cells reached confluence after approximately one week of culture, and then they were exposed to hydrostatic pressure (HP) ranging from 1 MPa to 50 MPa as previously described.¹² The Petri dishes were placed in a deformable Teflon pouch, which was filled with serum free DMEM containing the same concentrations of kanamycin and L-glutamine as mentioned above, and packed completely after removing air bubbles. The pouch then was placed in a stainless steel pressurisation vessel, exposed to 1, 5, 10, or 50 MPa of HP for two hours. After depressurisation, 3 ml of fresh serum free DMEM was given to each Petri dish, and the dish was maintained at atmospheric pressure in the air containing 5% CO₂ until analysis. Cells seeded in Petri dishes and placed in the same apparatus under the same conditions but not treated with HP exposure were used as non-pressurised controls.

ANALYSIS OF CYTOKINES AND PROTEOGLYCAN CORE PROTEIN MRNA IN THE CONTROLS AND HP EXPOSED CELLS

Total RNA was extracted from the controls and HP exposed cells between 30 minutes and eight hours after HP exposure by using the acid guanidinium thiocyanate-phenol chloroform method.¹³ The RNA preparations were analysed with a reverse transcription-polymerase chain reaction (RT-PCR) method for mRNA expressions of IL6 and TNF α ; and with northern blotting for mRNA expressions of proteoglycan core protein, stromelysin and TIMP1. β actin mRNA expression was examined by both RT-PCR and northern blotting, to ensure the quality of the RNA preparation and to prove the reliability of RT-PCR and northern blot protocols used in this study. RT-PCR assay was performed as previously described.¹⁴ Specific primer sets for IL6, TNF α , and β actin were designed based on the published sequence data.¹⁵⁻¹⁷ Table 1 summarises their primer positions. Each of the four doses of total RNA (0.2, 1, 2.5, and 5 μ g) was reverse transcribed with oligo (dT)16 primer by using MoMuLV RNase H-reverse transcriptase (GIBCO BRL Co, Gaithersburg, MD) in 10 μ l reaction mixture under the conditions recommended by the manufacturer. Specific sequences of cytokine mRNAs were amplified by using specific primers for them and Taq DNA polymerase (Wako Pure Chemical Industries), as previously described.¹⁴ RT-PCR products

Figure 3 Induction of TNF α mRNA expression after HP exposure. The cells at 30 minutes after the exposure to 50 MPa of HP expressed TNF α mRNA. Two point five microgram (a) and 1 μ g (b) of total RNA were reverse transcribed in 10 μ l reaction mixture. No TNF α mRNA expression was detected in the cells between four and eight hours after HP exposure. C: control cells treated under the atmospheric pressure.

were analysed in ethidium bromide containing agarose gels and evaluated for the expression of each cytokine's mRNA.

Northern blotting was performed as described elsewhere.¹⁸ The total RNA (7 μ g) was fractionated in a formaldehyde containing agarose gel, transferred to a nylon membrane, and prehybridised. The prehybridisation mixture was replaced with fresh solution containing 200 ng/ml of probes. Hybridisation was performed overnight at 65°C. Chemiluminescent detection by using DIG luminescent detection kits (Boehringer-Mannheim Biochemica, Mannheim, Germany) and Renaissance nucleic acid chemiluminescence reagent kit (DuPont NEN) was performed according to the manufacturer's specifications. Kodak

Figure 4 Effect of HP exposure on proteoglycan core protein mRNA expression. Cells were exposed to 1, 5, 10 or 50 MPa of HP for two hours, and total RNA was extracted at 30 minutes (A) and four hours (B) after the release of HP. Equal amounts of total RNA were fractionated by gel electrophoresis, transferred to a nylon membrane, and hybridised to RNA probes. For quantification, the signals were measured by using a densitometer. A photograph of gel staining (right side) shows equal amounts of RNA were applied. β actin mRNA expression is also shown. C: control cells treated under the atmospheric pressure.

XAR films were exposed to the nylon membranes for 5 to 30 minutes, and the films were analysed by using a densitometer, comparing the relative mRNA level of several samples on a single blot. We confirmed equal volumes of RNA were loaded by checking the intensity of ethidium bromide stained 18S and 28S rRNA bands. RNA loading was also examined by probing for β actin mRNA. The RNA probes for proteoglycan core protein, stromelysin, TIMP1, and β actin mRNA were prepared as described in a previous study.¹⁹

Results

In all duplicated or triplicated experiments for each test item, almost identical results were obtained. Figures are those of the representative case of the repeated experiments.

MORPHOLOGY OF HCS-2/8 CELLS AFTER HP EXPOSURE

After HP exposure, the cells were observed under a phase contrast light microscope. The HCS-2/8 cells have a polygonal shape under atmospheric pressure (fig 1A). This shape changed to almost spherical after the exposure to 50 MPa of HP for two hours (fig 1B). After the pressure was released, the cells regained their original polygonal shape within 24 hours (fig 1C). The cells that were exposed to lower magnitudes of HP (5 to about 10 MPa), demonstrated milder morphological changes (fig 1D), and returned to the polygonal shape soon after the pressure release. Exposure to 1 MPa of HP resulted in less morphological changes on the cells. Examination by using trypan blue dye exclusion assay showed that cellular viability was not affected by these HP levels.

EFFECTS OF HP ON CYTOKINE mRNA EXPRESSIONS IN HCS-2/8 CELLS

All PCR products having an expected size are evaluable in the agarose gel. mRNAs for β actin were clearly detected at a constant level in all samples. After the exposure to 1, 5, 10, or 50 MPa of HP for two hours, total RNAs were isolated from the cells and analysed identically by using PCR. IL6 mRNA and TNF α mRNA expressions were not observed in the control cells treated under the atmospheric pressure. In the PCR assays using various RNA concentrations, pressure dependent enhancement of IL6 mRNA expression on the cells was observed between 30 minutes and four hours after HP exposure (fig 2). No IL6 mRNA expression was observed eight hours after HP. TNF α mRNA expression was also increased at 30 minutes after the exposure to 50 MPa of HP and disappeared four hours after the release of HP (fig 3).

Northern blotting was performed to analyse the expressions of proteoglycan core protein, stromelysin, and TIMP1 in the cells exposed to 1, 5, 10, or 50 MPa of HP. Expression of β actin mRNA was found at an almost constant level in all samples. Proteoglycan core protein was detected in the cells grown under the atmospheric pressure, and its mRNA levels were increased 30 minutes and four hours after the exposure to 1 MPa or 5 MPa of HP. In particu-

Figure 5 Stromelysin and TIMP1 mRNA signals in HCS-2/8 cells after HP exposure. Representative signals at four hours after HP exposure are shown. C: control cells treated under the atmospheric pressure.

lar, the signal level for this mRNA in the cells exposed to 1 MPa was increased sixfold four hours after the release of HP. On the other hand, in the cells exposed to 10 MPa or 50 MPa of HP, proteoglycan core protein mRNA levels decreased 30 minutes and four hours after the release of HP (fig 4). At eight hours, the mRNA levels returned to almost the same as those under atmospheric pressure. Stromelysin and TIMP1 mRNA signals did not show major response to HP during eight hours after the release of HP (fig 5).

Discussion

The effect of some types of pressure, for example, tension,²⁰ shearing force,⁴ and HP,²¹⁻²³ has been studied using tissue sections²¹⁻²² and cultured cells.²³ HP is considered to have the most influential effect on the metabolism of cartilage.²⁴ A recent study²⁵ demonstrated that chondrocytes and extracellular matrix in different zones of a tissue react differently to mechanical loading. Although tissue culture better represents *in vivo* conditions, direct compressive force given to cartilage tissue will result in such phenomena as deformation of tissues and cells,²⁶ effusion of tissue fluid,²⁷ and increase of proteoglycan levels followed by the increase of cation.²⁴ These factors will affect experimental results, and make the analysis of pressure effect quite difficult. Therefore, we used cultured cells, instead of tissue sections, in the investigations of the effect of HP on the expression of metabolism related cytokines.

HCS-2/8 is a chondrosarcoma cell line, established in 1989 by Takigawa *et al.*¹⁰ HCS-2/8 produces Type II, IX, and XI collagens, as well as cartilage type proteoglycan (aggrecan), as seen in normal chondrocytes. In addition, the reaction of HCS-2/8 to various vitamins and growth factors is similar to those of normal chondrocytes.¹¹ This cell line can maintain these characteristics of normal chondrocytes for a long culture period, and can be a useful model of human chondrocytes. Our study, for the first time, demonstrated that HP can induce expression of IL6 and TNF α mRNAs in the cultured cells that possess characteristic features of normal chondrocytes. In the measurements using RT-PCR, IL 6 and TNF α mRNA expressions in the control cells without giving exposure to HP were all negative, while the expressions became positive after exposure to HP regardless of the RNA concentrations and this change was reproducible. Therefore, our RT-PCR findings were evaluated to be valid.

During walking, pressure on a human hip joint ranges between 3 MPa and 10 MPa.²⁸ It

sometimes reaches approximately 20 MPa²⁹ and the level varies continuously. In a pathological condition, for example, OA, much higher HP could occur. We previously reported¹² that constant 50 MPa of HP induced heat shock protein in HCS-2/8, the expression of which is increased in OA cartilage,^{30, 31} and this magnitude of HP adversely affects the metabolism of these cells. Therefore, constant 50 MPa was considered to represent an abnormally high and pathological level of stimulus. IL6 and TNF α have been considered as cytokines that are induced by inflammation or immune reactions. In this study, these cytokines were also induced after exposure to a unphysiologically high pressure level, 50 MPa. This indicates that IL6 and TNF α , which are increased in OA chondrocytes, can be induced by unphysiologically high, excessive pressure. This is important when considering the mechanism of OA development because of an extremely high pressure level.

In this study, cell morphology changed to spherical after giving 50 MPa, while this change was mild under low pressure levels. On the other hand, IL6 mRNA expression was most noticeable at 50 MPa. Our findings on morphological changes after applying HP agree with those of previous studies, and show cytoskeletons, such as actin filament, changed under the HP.^{32, 33} Mohtai *et al* reported that 1.6 Pa of fluid-induced shear induced the expression of IL6 mRNA and IL6 protein,⁴ and chondrocytes exposed to fluid induced shear for 48 hours appeared to be elongated and elliptical and their major axes aligned primarily circumferentially.³⁴ On the other hand, cytoskeletons, for example, actin filament, are believed to be involved in the mRNA expression and play quite an important part in the local expression and function of mRNA.³⁵⁻³⁷ These findings suggest that changes in actin filament could be involved closely with IL6 induction by mechanical stress. In the same way, the changes in actin filament could also be involved in TNF α induction under 50 MPa of HP. Mechanotransduction of cells has been reported by many researchers, and it is thought that ion channels take an important part in the initial response to mechanical stress, and changes in intracellular calcium ion concentrations and cAMP production are largely involved in the regulation of gene expression caused by mechanical stress.³⁸ It has been hypothesised that mechanical stress is transferred from extracellular matrix to nuclei via cytoskeleton, which is connected to integrins on cellular surface.³⁹ In OA cartilage tissues, the number of chondrocytes that are in the dynamic process of polymerisation and depolymerisation of cytoskeleton elements, is reported to be increased.⁴⁰ Buschmann *et al*⁴¹ demonstrated the importance of cell matrix interaction in the cellular reaction against pressure by using the primary chondrocyte/agarose culture. HCS-2/8 in plate culture and at a confluent level is reported to express integrin and to generate large volume of chondrocyte-like matrix, and this matrix is accumulated around the cell.^{10, 42} In this study, we loaded HP to

HCS-2/8 at a confluent level, but this cell at a sparse or subconfluent level reacted less apparently to HP (unpublished data). Therefore, we consider that integrin and extracellular matrix may also take an important place in the present findings.

Cytoskeleton may also be involved in the regulation of proteoglycan synthesis in chondrocytes.⁴³ We previously reported that 1 or 5 MPa of HP induced an increase of ³⁵S sulphate uptake into chondrocyte-like cells, and 10 or 50 MPa of HP induced a decrease.¹² In this study, expression of proteoglycan core protein mRNA increased after giving 1 or 5 MPa of HP, and it decreased after giving 10 or 50 MPa. This change in the expression volume caused by HP was larger than the changes in ³⁵S sulphate uptake. On this point, CS-Szabo *et al* reported that changes in the levels of proteoglycan core protein mRNA were severalfold greater than expected.⁴⁴ Therefore, our findings on proteoglycan core protein mRNA are thought to be consistent to the previous findings, and as shown in another study,⁴⁵ this shows that HP affects extracellular matrix production on transcriptional level. TNF α is reported to induce IL6,⁶ but in the present study, expression of TNF α mRNA occurred transiently immediately after the exposure to HP. Therefore, we cannot conclude whether TNF α directly induced IL6. Our chronological findings also suggest that these cytokines would not affect proteoglycan expression.

In this study, HP did not affect the expression of stromelysin and TIMP1 mRNAs, and this was in contrast with our expectation. In OA, imbalance between proteases and inhibitors is thought to be implicated,⁴⁶ but constant HP for two hours did not influence the imbalance at transcriptional level. We also reported¹² that HP increased the expression of TGF β 1, which can induce TIMP1,⁴⁷ and shear stress to chondrocytes is reported to increase TIMP1 mRNA expression.³⁴ TNF α increases stromelysin expression,⁶ and IL6 increases TIMP1 expression.⁸ The difference between our findings and expectation could be attributable to the constant HP used in this study, and to the monitoring of mRNAs after the exposure to HP. We applied constant pressure instead of cyclic pressure, because our purpose was to isolate the specific regulatory signals after giving abnormal pressure. Because Smith *et al*⁴⁵ reported that cells react to cyclic pressure more noticeably than to constant pressure, future studies using cyclic HP and that monitor gene and protein expressions would be of interest. In addition, HCS-2/8 is not only a chondrocyte-like cell line but also a transformed cell line, and transformed cell lines are known to be resistant to HP in comparison to primary cultures.⁴⁸ To better understand our findings, mRNA studies should be repeated in primary chondrocyte cultures. Monitoring the IL6 and TNF protein productions in addition to mRNA, and comparison between IL6 induction by HP and that by such an inflammatory cytokine as IL1, merit future studies.

This study received a grant in aid for scientific research from the Ministry of Education, Science, and Culture of Japan.

- 1 Dippe P, Kirwan J. The localization of osteoarthritis. *Br J Rheumatol* 1994;33:201-4.
- 2 Muir H, Carney SL. Pathological and biochemical changes in cartilage and other tissues of canine knee resulting from induced joint instability. In: Helminen HJ, Kirivanta I, Tammi M, Saamanen A-M, Paukkonen K, Urvelin J, eds. *Joint loading: biology and health of articular structures*. Bristol: John Wright, 1987; 47-63.
- 3 Hoch DH, Grodzinsky AJ, Koob TJ, Albert ML, Eyre DR. Early changes in material properties of rabbit articular cartilage after meniscectomy. *J Orthop Res* 1983;1:4-12.
- 4 Mohtai M, Gupta MK, Donlon B, Ellison B, Cooke J, Gibbons G, et al. Expression of interleukin-6 in osteoarthritic chondrocytes and effects of fluid-induced shear on this expression in normal human chondrocytes in vitro. *J Orthop Res* 1996;14:67-73.
- 5 Kammermann JR, Kincaid SA, Rumph PF, Baird DK, Visco DM. Tumor necrosis factor- α (TNF- α) in canine osteoarthritis: Immunolocalization of TNF- α , stromelysin and TNF receptors in canine osteoarthritic cartilage. *Osteoarthritis Cartilage* 1996;4:23-34.
- 6 Tyler JA, Bolis S, Dingle JT. Mediators of matrix catabolism. In: Kuettner KE, Schleyerbach R, Peyron JG, Hascall VC, eds. *Articular cartilage and osteoarthritis*. New York: Raven Press, 1992: 251-64.
- 7 Guerne PA, Carson DA, Lotz M. IL-6 production by human articular chondrocytes: modulation of its synthesis by cytokines, growth factors, and hormones in vitro. *J Immunol* 1990;144:499-505.
- 8 Lotz M, Guerne PA. Interleukin-6 induces the synthesis of tissue inhibitor of metalloproteinases-1/erythroid potentiating activity (TIMP-1/EPA). *J Biol Chem* 1991;266: 2017-20.
- 9 Venn G, Nietfeld JJ, Duits AJ, Brennan FM, Arner E, Covington M, et al. Elevated synovial fluid levels of interleukin-6 and tumor necrosis factor associated with early experimental canine osteoarthritis. *Arthritis Rheum* 1993;36:819-26.
- 10 Takigawa M, Tajima K, Pan H-O, Emoto M, Kinoshita A, Suzuki F, et al. Establishment of a clonal human chondrosarcoma cell line with cartilage phenotypes. *Cancer Res* 1989;49:3996-4002.
- 11 Enomoto MI, Takigawa M. Regulation of tumor-derived and immortalized chondrocytes. In: Adolphe M, ed. *Biological regulation of the chondrocytes*. Boca Raton: CRC Press, 1992:321-38.
- 12 Takahashi K, Kubo T, Kobayashi K, Imanishi J, Takagawa M, Arai Y, et al. Hydrostatic pressure influences mRNA expression of transforming growth factor- β 1 and heat shock protein 70 in chondrocyte-like cell line. *J Orthop Res* 1997;15:150-8.
- 13 Chomczynski P, Sacchi N. Single-step method of RNA isolation by acid guanidinium thiocyanate-phenol-chloroform extraction. *Anal Biochem* 1987;162:156-9.
- 14 Sotozono C, Kinoshita S, Kita M, Imanishi J. Paracrine role of keratinocyte growth factor in rabbit corneal epithelial cell growth. *Exp Eye Res* 1994;59:385-92.
- 15 Yasukawa K, Hirano T, Watanabe Y, Muratani K, Matsuda T, Nakai S, et al. Structure and expression of human B cell stimulatory factor-2 (BSF-2/IL-6) gene. *EMBO J* 1987;6: 2939-45.
- 16 Shirai T, Yamaguchi H, Ito H, Todd CW, Wallace RB. Cloning and expression in *Escherichia coli* of the gene for human tumor necrosis factor. *Nature* 1985;313:803-6.
- 17 Albino AP, Davis BM, Narus DM. Induction of growth factor RNA expression in human malignant melanoma: markers of transformation. *Cancer Res* 1991;51:4815-20.
- 18 Kobayashi K, Ohgitani E, Tanaka Y, Kita M, Imanishi J. Herpes Simple virus-induced expression of 70 kD heat shock protein (HSP70) requires early protein synthesis but not viral DNA replication. *Microbiol Immunol* 1994;38: 321-5.
- 19 Arai Y, Kubo T, Kobayashi K, Takeshita K, Takahashi K, Ikeda T, et al. Adenovirus vector-mediated gene transduction to chondrocytes: In vitro evaluation of therapeutic efficacy of transforming growth factor- β 1 and heat shock protein 70 gene transduction. *J Rheumatol* 1997;24:1787-95.
- 20 Dewitt MT, Handley CJ, Oakes BW, Lowther DA. In vitro response of chondrocytes to mechanical loading. The effect of short term mechanical tension. *Connect Tissue Res* 1984;12:97-110.
- 21 Hall AC, Urban JPG, Gehl KA. The effects of hydrostatic pressure on matrix synthesis in articular cartilage. *J Orthop Res* 1991;9:1-10.
- 22 Lipiello L, Kaye C, Neumata T, Mankin HJ. In vitro metabolic response of articular cartilage segments to low levels of hydrostatic pressure. *Connect Tissue Res* 1985;13:99-107.
- 23 Van Kampen GPJ, Veldhuizen JP, Kuijter R, van der Stadt RJ, Schipper CA. Cartilage response to mechanical forces studied in high density cultures. *Arthritis Rheum* 1985;28: 419-24.
- 24 Urban JPG. The chondrocyte: a cell under pressure. *Br J Rheumatol* 1994;33:901-8.
- 25 Wong M, Wuethrich P, Buschmann MD, Eggli P, Hunziker E. Chondrocyte biosynthesis correlates with local tissue strain in statically compressed adult articular cartilage. *J Orthop Res* 1997;15:189-96.
- 26 Broom ND, Myers DB. A study of the structural response of wet hyaline cartilage to various loading stresses. *Connect Tissue Res* 1980;7:227-37.
- 27 Mow VC, Holmes MH, Lai WM. Fluid transport and mechanical properties of articular cartilage. *J Biomech* 1984;17:377-94.
- 28 Afohe NYP, Byers PD, Hutton WC. Contact pressures in the human hip. *J Bone Joint Surg Am* 1987;69B:536-41.
- 29 Hodge WA, Fijan RS, Carlson KL, Burgess RG, Harris WH, Mann RW. Contact pressures in the human hip joint measured in vivo. *Proc Natl Acad Sci USA* 1986;83:2879-83.
- 30 Takahashi K, Kubo T, Arai Y, Imanishi J, Kawata M, Hirasawa Y. Localization of heat shock protein in osteoarthritic cartilage. *Scand J Rheumatol* 1997;26:368-75.
- 31 Takahashi K, Kubo T, Goomer RS, Amiel D, Kobayashi K, Imanishi J, et al. Analysis of heat shock proteins and cytokines expressed during early stages of osteoarthritis in a mouse model. *Osteoarthritis Cartilage* 1997;5:321-9.
- 32 Parkkinen JJ, Lammi MJ, Inkinen R, Jortikka M, Tammi M, Virtanen I, et al. Influence of short-term hydrostatic pressure on organization of stress fibers in cultured chondrocytes. *J Orthop Res* 1995;13:495-502.
- 33 Haskin C, Cameron I. Physiological levels of hydrostatic pressure alter morphology and organization of cytoskeletal and adhesion proteins in MG-63 osteosarcoma cells. *Biochem Cell Biol* 1993;71:27-35.
- 34 Smith LR, Donlon BS, Gupta MK, Mohtai M, Das P, Carter DR, et al. Effects of fluid-induced shear on articular chondrocyte morphology and metabolism in vitro. *J Orthop Res* 1995;13:824-31.
- 35 Fey EG, Ornelles DA, Penman S. Association of RNA with the cytoskeleton and the nuclear matrix. *J Cell Sci Suppl* 1986;5:99-119.
- 36 Bagchi T, Lawson DE, Sells BH. Cytoskeletal association of muscle-specific mRNAs in differentiating L6 rat myoblast. *Exp Cell Res* 1987;168:160-72.
- 37 Lawrence JB, Singer BH. Intracellular localization of messenger RNAs for cytoskeletal proteins. *Cell* 1986;45: 407-15.
- 38 Wright MO, Stockwell RA, Nuki G. Response of plasma membrane to applied hydrostatic pressure in chondrocytes and fibroblasts. *Connect Tissue Res* 1992;28:49-70.
- 39 Wang N, Butler JP, Ingber DE. Mechanotransduction across the cell surface and through the cytoskeleton. *Science* 1993;260:1124-7.
- 40 Kouri JB, Jimenez SA, Quintero M, Chico A. Ultrastructural study of chondrocytes from fibrillated and non-fibrillated human osteoarthritic cartilage. *Osteoarthritis Cartilage* 1996;4:111-25.
- 41 Buschmann MD, Gluzband YA, Grodzinsky AJ, Hunziker EB. Mechanical compression modulates matrix biosynthesis in chondrocyte/agarose culture. *J Cell Sci* 1995;108: 1497-508.
- 42 Tuckwell DS, Ayad S, Grant ME, Takigawa M, Humphries MJ. Conformation dependence of integrin-typeII collagen binding. Inability of collagen peptides to support $\alpha_2\beta_1$ binding, and mediation of adhesion to denatured collagen by a novel $\alpha_2\beta_1$ -fibronectin bridge. *J Cell Sci* 1994;107: 993-1005.
- 43 Newman P, Watt FM. Influence of cytochalasin D-induced changes in cell shape on proteoglycan synthesis by cultured articular chondrocytes. *Exp Cell Res* 1988;178:199-210.
- 44 CS-Szabo G, Melching LI, Roughley PJ, Glant TT. Changes in messenger RNA and protein levels of proteoglycans and link protein in human osteoarthritic cartilage samples. *Arthritis Rheum* 1997;40:1037-45.
- 45 Smith LR, Rusk SF, Ellison BE, Wessells P, Tsuchiya K, Carter DR, et al. In vitro stimulation of articular chondrocyte mRNA and extracellular matrix synthesis by hydrostatic pressure. *J Orthop Res* 1996;14:53-60.
- 46 Dean DD, Martel-Pelletier J, Pelletier J-P, Howell DS, Woessner JFJ. Evidence for metalloproteinase inhibitor imbalance in human osteoarthritic cartilage. *J Clin Invest* 1989;84:678-85.
- 47 Gunter M, Haubeck H-D, van de Leur E, Blaser J, Bender S, Gutgenann I, et al. Transforming growth factor β 1 regulates tissue inhibitor of metalloproteinases-1 expression in differentiated human articular chondrocytes. *Arthritis Rheum* 1994;37:395-405.
- 48 Dibb W, Morild E, Laerum OD. Effects of high pressure on normal and neoplastic rat cells in culture. *Virchows Arch [Cell Pathol]* 1981;38:169-76.