SAT0002 COMPARISON OF TWO EXERCISE REGIMENS IN PATIENTS WITH ANKYLOSING SPONDYLITIS

KK Trnavsky, Y Sulcova, J Knebertova. *Rheumatology, Postgraduate Medical School, Arthocentrum, Praha, Czech Republic*

10.1136/annrheumdis-2001.354

Background Exercise alone can produce adequate relief of symptoms in ankylosing spondylitis (AS) and manipulative therapy has also been shown to improve range of movement in short ? time.

Objectives To compare the effect of a dynamic individual rehabilitation regimen with a simple conservative exercise regimen (exercies in goupes).

Methods Forty patients with established AS, mean age 51 years and mean disease duration 10 years were randomly assigned to group of 20 patients each. The treatment programme consisted of two thre ? months periods.: A. Intensive individual programme (assisted by an experienced physiotherapist) included thermotherapy. Soft techniques, relaxation of shortened muscles, breathing exercises. B. Current exercise programme in groups. Metrologic assessment and questionnaires were used to evaluate the activity of the process (BASDAI) and the functional handicap of the patient (BASFI). In the first group of patients both the programme A and B were applied. The second group underwent only programme B. After 3 months the therapeutic programmes in the two groups were exchanged.

Results The combined exercise regimen involving individual therapeutic care resulted in improvement of parameters assessing spinal column mobility ? Schober ??s distance, tips to the floor distance, inclination and retro ? and anteflexion of cervical column. The indicators of function and disease activity evaluated according to BASDAI and BASFI questionnaires showed also improvement but the differences did no reach statistical significance over the three-months period of follow-up.

Conclusion Intensive individual exercise programme including soft techniques and relaxation of shortened muscles is more effective in AS than a conservative programme based on exercise in groups.

SAT0003 INTESTINAL COLONISATION WITH SULPHATE-REDUCING BACTERIA: A CO-FACTOR IN THE AETIOPATHOGENESIS OF ANKYLOSING SPONDYLITIS?

¹SM Stebbings, ¹J Highton, ²G Tannock. ¹*Rheumatology*; ²*Microbiology, University of Otago School of Medicine, Dunedin, New Zealand*

10.1136/annrheumdis-2001.355

Background Sulphate-reducing bacteria (SRB) have only recently been identified as a constituent of the normal colonic microflora humans. Such colonisation however is far from universal. Small studies in patients with ulcerative colitis (UC) have found SRB to be present in up to 92% of patients and a pathogenic role has been speculated. Approximately 65% of patients with ankylosing spondylitis (AS) have colonic inflammation, and this is now recognised as an independent risk factor for the disease. Furthermore, studies in germfree HLA-B27 transgenic rats have shown that the normal intestinal microflora is essential to the development of spondyloarthritis in these animals.

Objectives We aimed to discover whether intestinal colonisation with SRB was more common in AS patients than controls, and thus whether SRB may play a role in the aetiology of AS.

Methods Fifteen patients fulfilling the Modified New York criteria for AS and 15 age and gender matched controls were recruited. Each subject provided 3 faecal specimens at monthly intervals. Bacterial DNA extracts were obtained from 3 faecal samples from 14 patients and 14 controls, and one each from the remaining patient and control. PCR was performed using primers targeting a conserved region of subunit A of the adenosine-5?-phosphosulphate (APS) reductase gene a gene conserved across all SRB species.

Results (See Table 1)

Chi-square with Yates? correction demonstrated a highly significant association between AS and colonisation with SRB (p = 0.003).

Abstract SAT0003 Table 1

	PCR positive for SRB	PCR negative for SRB	Total
Ankylosing spondylitis	24 (56%)	19 (44%)	43
Controls	7 (16%)	36 (84%)	43
Total number of samples			86

Identification of sulphate-reducing bacteria in faecal samples using PCR.

Conclusion We have discovered a much higher prevalence of faecal carriage of SRB in patients with AS compared with healthy controls. This raises the possibility that these organisms may play a role in the aetiopathogenesis of AS. An alternative possibility is that SRB reflect the presence of colonic inflammation in these individuals.

REFERENCES

- 1 Gibson GR, Cummings JH, Macfarlane GT. Growth and activities of sulphate-reducing bacteria in gut contents of healthy subjects and patients with ulcerative colitis. *FEMS Microbiol Ecol.* 1991;**86**:103-12
- 2 Veys EM, Mielants H, De Vos M, Cuvelier C. Spondylarthropathies: from gut to target organs. *Baillieres Clin Rheumatol.* 1996;**10**(1):123-46
- 3 Taurog JD, Richardson JA, Croft JT, et al. The germfree state prevents development of gut and joint inflammatory disease in HLA-B27 transgenic rats. *J Exp Med.* 1994;**180**(6):2359-64

SAT0004 IS THE ASSOCIATION SJÖGREN'SYNDROME-SPONDYLARTHROPATHY FORTUITOUS?

C Scotto di Fazano, E Quennesson, S Quennesson, RM Grilo, D Couaral, P Vergne, C Bonnet, P Bertin, R Treves. *Rheumatology, University Hospital Dupuytren, Limoges, France*

10.1136/annrheumdis-2001.356

Background Secondary Sjögren's syndrome is known to occur with rheumatoid arthritis, systemic lupus erythematosus, but apart from a few cases, has not been reported with spondylarthropathy (SpA).

Objectives To determine the prevalence of Sjögren's syndrome in spondylarthropathy patients.

Methods Patients with SpA (n = 62) with inflammatory back pain and/or peripheral arthritis were diagnosed as Ankylosing Spondylitis (AS), undifferentiated SpA (uSpA), psoriatic Arthritis (PA), and arthritis associated with inflammatory bowel diseases (IBD). The investigation of sicca symptoms in the group SPD and healthy controls (n = 102) was based on a valited questionnaire. Patients with SpA with sicca symptomes and/or positive antinuclear antibodies (ANA) were investigated for Sjögren's syndrome by minor salivary gland biopsy.

In the spondylarthropathy group, the laboratory variables were: HLA B27, HLA DR, DQ, anti ENA, serology CMV, EBV, HIV, hepatitis B, C.

Results In 17/62 SpA patients (11 uSpA, 5 AS, 1 PA, 17 females, 1 male) Sjögren's syndrome diagnosis by the European criteria indicated a frequency of 27.4%. Of 62 SpA, 12 were ANA positive (19.3%), of whom 5 had Sjögren's syndrome. ANA were detected in 8/17 (47%) patients with Sjögren's syndrome. Of 62 SpA, 20 had sicca symptoms (32.2%), compared to 3 of 102 (2.9%) controls. HLA B27 is positive in 12/17 patients (70.5%). The results of HLA DR, DQ seem to show an high frequency of HLA DR4, and DQ3 in the association group but the results are not significant.

Conclusion The frequency of Sjögren's syndrome is 27.4% in the spondylarthropathy group, thus more than in controls. Thus, the association seems to be not fortuitous. And it's interesting to see more females in a group with more undifferentiated spondylarthropathy: 16/17 are females.

SAT0005 FACTORS INFLUENCING FATIGUE AND IMPACT OF FATIGUE ON QUALITY OF LIFE IN ANKYLOSING SPONDYLITIS (AS) PATIENTS

¹A Van Tubergen, ¹J Coenen, ¹R Landewé, ¹A Spoorenberg, ²A Chorus, ¹A Boonen, ¹S Van der Linden, ¹D Van der Heijde. ¹Rheumatology, University Hospital Maastricht, Maastricht; ²Division Public Health, TNO Prevention and Health, Leiden, The Netherlands

10.1136/annrheumdis-2001.357

Background

Objectives 1. To identify factors that influence fatigue in AS. 2. To assess whether fatigue is associated with quality of life in AS.

Methods A total number of 812 AS patients were included. Fatigue was measured with a single-item question elicited from the Bath AS Disease Activity Index (BASDAI) and with the Multidimensional Fatigue Inventory (MFI). The patients also completed questionnaires on functional ability (BASFI), global well-being (BAS-G), and quality of life (SF-36). In addition, patients were asked to identify factors that influence fatigue.

Patients were dichotomously divided into a F+ group if the BASDAI fatigue score was >5.0, labelled as experiencing fatigue as a major symptom, and a F- group in case the fatigue score was <5.0 (fatigue a minor symptom). Logistic regression analysis (dependent variable F+/F-) was used to determine which factors were associated with fatigue. Multiple regression analysis was used to investigate whether fatigue contributes in explaining quality of life (dependent variable dimensions from the SF-36; independent variable BASDAI fatigue question). In both analyses adjustments were made for age, sex, disease duration, and the presence of co-morbidity.

Results 53% of the patients were categorised to the F+ group. They scored significantly worse compared to the F- group with respect to each dimension of the MFI and to all other questionnaires studied (all $p < 0.001$). Fatigue was, in the opinion of the patients, highly influenced by pain (70% of the patients) and stiffness (54%). The hierarchical order of factors perceived to be associated with fatigue was grossly similar in the F+ and F- groups, but pain, poor sleep and work were significantly more often considered important by patients from the F+ group. Logistic regression analysis showed that scores on BASDAI (without fatigue question), BASFI and BAS-G were independently associated with fatigue (R^2 : 0.52). Multiple regression analysis showed that fatigue was significantly associated with all separate dimensions of the SF-36 studied ($p < 0.001$).

Conclusion Fatigue is a major symptom in AS and is associated with disease activity, functional ability, and global well-being. Fatigue independently contributes negatively to different dimensions of quality of life.

SAT0006 ASSOCIATION OF THORACIC KYPHOSIS WITH FUNCTIONAL ABILITY, GLOBAL WELL-BEING, AND QUALITY OF LIFE IN ANKYLOSING SPONDYLITIS

¹A Van Tubergen, ¹S Van der Linden, ¹R Landewé, ¹D Van der Heijde, ¹D Vosse, ²P Geusens. ¹Rheumatology, University Hospital Maastricht, Maastricht, The Netherlands; ²Rheumatology, Limburg University Center, Diepenbeek, Belgium

10.1136/annrheumdis-2001.358

Background

Objectives To determine the association of increased thoracic kyphosis with functional ability, global well-being, and quality of life in ankylosing spondylitis (AS) patients.

Methods Baseline data from 120 Dutch AS outpatients participating in a trial on the efficacy of spa therapy were used. Patients were selected on having an occiput-wall distance of 0.0 cm (group A) or an occiput-wall distance > 10.0 cm (group B), defining no and severe kyphosis respectively.

The Bath Ankylosing Spondylitis Functional Index (BASFI) (scale 0–10), Dougados Functional Index (DFI) (scale 0–40) and the Health Assessment Questionnaire for Spondylarthropathies (HAQ-S) (scale 0–3) were used to assess functional ability. Global well-being was assessed with a 10 cm visual analogue scale (scale 0–10), and quality of life with the AS Quality of Life questionnaire (ASQoL) (scale 0–18). For each instrument counts that 0 is the best score.

Results Group A consisted of 21 patients (9 men, 12 women, mean age 44 (8) years), group B of 26 patients (23 men, 3 women, mean age 51 (10) years). After adjustment for age and gender with multiple regression analysis similar results were found.

Abstract SAT0006 Table 1

	Group A	Group B	Difference (95% CI); p-value
BASFI	4.3 (1.8)	6.0 (1.5)	1.7 (0.7 to 2.7); $p = 0.001$
DFI	9.9 (4.1)	15.3 (4.2)	5.4 (3.0 to 7.8); $p < 0.001$
HAQ-S	0.8 (0.4)	1.2 (0.4)	0.4 (0.2 to 0.6); $p = 0.001$
Global well-being	5.3 (2.6)	5.7 (1.7)	0.4 (-1.0 to 1.7); $p = 0.596$
ASQoL	8.7 (4.5)	9.2 (4.4)	0.5 (-2.1 to 3.1); $p = 0.710$

Conclusion An increased occiput-wall distance is associated with deteriorating functional ability. Global well-being and perceived quality of life seem to be generally affected in AS patients, but an increased occiput-wall distance is not associated with further deterioration. The latter merits further investigation.

SAT0007 ANKYLOSING SPONDYLITIS INITIATED AS A REACTIVE ARTHRITIS

ZA Curcic. Institute of Rheumatology, School of Medicine, Novi Sad, Yugoslavia

10.1136/annrheumdis-2001.359

Background