Conclusion Taken together, these data show that alleles at class I loci are associated with the disease, although it is difficult to assess their specificity to PsA genetic susceptibility. No particular role of the HLA class II region could be demonstrated.

SAT0024 ENTHESITIS AND DACTYLITIS AS THE BEGINNING STAGE OF PSORIATIC ARTHRITIS

C Montilla, CM González, FJ López-Longo, M Moreno-Zazo, R Castillo, N Caro, L Cebrián, C González-Montagut, S Gómez, I Monteagudo, L Carreño. *Rheumatology, Hospital General Universitario Gregorio Marañón, Madrid, Spain*

10.1136/annrheumdis-2001.376

Background Enthesitis and dactylitis are characteristic features of psoriatic arthritis and the others spondyloarthropathies. Both may appear from the onset of the disease.

Objectives To analyze the clinical characteristic of a population of patients with psoriatic arthritis who presented enthesitis and dactylitis as their only clinical symptoms at the onset of the disease.

Methods We have studied a group of 204 patients diagnosed with psoriatic arthritis following the Moll and Wright criteria, selecting only those who expressed either Enthesitis or dactylitis as initial symptoms of the disease.

Results Twenty-eight patients (14%) proved to have Enthesitis or dactylitis 21 enthesitis, 7 dactylitis and 1 Enthesitis and dactylitis as disease onset, and seventeen were men (60.7%). The mean age of psoriatic arthritis onset was 38 years (range: 17–58). The mean time from Enthesitis and dactylitis onset and the Moll and Wright criteria was 6 months (range: 1–180). Enthesitis was most frequent in achilles tendon (81%) and the plantar fascia (72.7%). Twenty patients (71.4%) presented arthritis involving peripheral joints, 4 (14.3%) presented and axial skeleton distribution, and the remaining (14.3%) showed a mixed peripheral and axial distribution.

Conclusion Enthesitis and dactylitis are initial clinical features of psoriatic arthritis, even though the majority of the patients with these symptoms at the beginning of the disease evolve towards a peripheral arthritis pattern.

SAT0025 EVOLUTION OF PSORIATIC ARTHRITIS WITH DISTAL INTERPHALANGEAL JOINTS INVOLVEMENT AT ONSET

CA Montilla, CM González, FJ López-Longo, R Castillo, N Caro, C González-Montagut, L Cebrián, S Gómez, E Carmona, I Monteagudo, L Carreño. *Rheumatology, Hospital General Universitario Gregorio Marañón, Madrid, Spain*

10.1136/annrheumdis-2001.377

Background Distal interphalangeal joints (DIP) arthritis described by Moll and Wright as a psoriatic arthritis (Psa) pattern is now considered an evolutionary step towards other classification patterns.

Objectives To study the clinical and radiological evolution of a group of patients with arthritis limited to the DIP as onset of Psa.

Methods We have studied the clinical and radiological evolution of 204 patients diagnosed with Psa (Moll and Wright criteria) during the first year of the disease selecting those patients whose initial clinical features were exclusively arthritis of the DIP. The radiological findings after one-year follow-up were classified into stages according to Steinbrocker's criteria.

Results Arthritis of DIP as onset of the disease was found in 19/204 (9.3%) of the patients. The mean age at onset was 49 years (range: 34–64). Psoriasis before arthritis onset was present in 8/19 (42.1%) and in 9/19 (47.3%) the onset of arthritis and psoriasis was simultaneous. Onychodystrophy (pitting) was present in 7/19 (36.8%). After one-year 9/19 (47.3%) patients continued with arthritis as the only manifestations of Psa, 8/19 (42.1%) patients evolved toward other arthritis patterns (5 asymmetrical oligoarthritis and 3 symmetrical polyarthritis). The joints more commonly involved along with the DIP were metacarpophalangeal (75%) and proximal interphalangeal (62.5%). 2/19 patients (10.4%) did not show DIP arthritis after one year. At the end of the one-year study 9/19 (47.3%) of the patients were classified into III stage radiological criteria, 2/19 (10.4%) were at the stage I and in the remaining patients radiological was considered as normal.

Conclusion Almost half of the patients with DIP arthritis as Psa onset evolve within one year towards others arthritis patterns. Half of the patients revealed an aggressive radiological progression during the first year of the disease.

SAT0026 IS RADIOGRAPHIC EXAMINATION SUFFICIENT FOR DETECTION OF NON TRAUMATIC SPINAL FRACTURES IN ANKYLOSING SPONDYLITIS: COMPARISON OF RADIOGRAPHIC AND MRI FINDINGS

¹H Przepiera-Bedzak, ²L Cyrylowski, ¹M Brzozko. ¹Rheumatology; ²Radiology, Pomeranian Academy of Medicine, Szczecin, Poland

10.1136/annrheumdis-2001.378

Background

Objectives To compare plain radiographic and MRI findings in evaluation of non traumatic spinal fractures in patients with ankylosing spondylitis (AS), and to find correlation between spinal fractures and disease activity, age of patients, and disease duration.

Methods 21 males and 2 females with AS, mean aged 49.8, were studied. Lateral radiographs and MRI examinations of lumbar, thoracic and cervical spine were performed in all patients. To detect vertebral body fractures with radiographs, the anterior, posterior and central heights of each vertebral body were measured. The ratio of each height to that of the two adjacent vertebrae, the ratio of the anterior height to the posterior height and of the central height to the posterior height were calculated. The subjects were considered to have fracture if any of these ratios was less than 0.85 for lumbar spine and less than 0.8 for thoracic and cervical spine. We assessed also the following indexes of disease activity: BASFI, BASDAI, BASG-t, BASG-6, BSMI.

Results Total 25 fractures were found with the radiographic method: 7 fractures in lumbar spine, 15 in thoracic and 3 in cervical. The MRI examinations revealed total 51 fractures: 24 fractures in lumbar spine, 18 in thoracic and 9 in cervical. Out of 25 fractures detected by X-ray examination, MRI confirmed the site of fracture only in 10 (40%). Age of patients and disease duration correlated positively with total number of fractures, number of fractures in lumbar spine and number of fractures in thoracic detected by MRI. There was no such correlation for fractures detected by X-ray. Assessing indexes of disease activity we found positive correlation between BASFI, BASDAI, BASMI and total number of fractures and number of fractures in lumbar spine detected by MRI.

Conclusion The significant difference between radiographic and MRI findings in detecting of vertebral fractures was found. The results suggest that plain X-rays cannot be considered as a sufficient method for detection vertebral fractures in patients with AS. MRI should be considered as the method of choice in revealing spinal fractures in AS patients. Disease activity may contribute in developing of non – traumatic spinal fractures in this group of patients.

SAT0027 HIP JOINT INVOLVEMENT IN PSORIATIC ARTHRITIS (PSAR)

CJ Michet. *Department of Rheumatology, Mayo Clinic, Rochester, USA*

10.1136/annrheumdis-2001.379

Background Long term studies of the spondyloarthropathies have identified early hip joint involvement as a marker for severe disease and disability. Psoriatic spondylitis (PsSp) results in hip arthroplasty more frequently than primary ankylosing spondylitis. It is not clear whether the risk for hip involvement exists only for PsSp or other variants of PsAr.

Objectives To evaluate the prevalence of hip joint involvement cohort of PsAr patients. To determine whether gender, type of psoriasis, PsAr peripheral joint pattern, age of onset of disease, presence of spondylitis are associated with the risk of hip joint involvement. To determine the frequency of hip arthroplasty in these patients.

Methods This is a retrospective cohort study of 417 patients with PsAr treated at a single tertiary academic centre over a 15 year period. Patient characteristics and outcome were abstracted by chart review. Radiographs of patients with hip joint disease were reviewed.

Results 27 PsAr patients (6.5%) were diagnosed with hip joint involvement. 60% had bilateral involvement. Gender, type of psoriasis, nail involvement, pattern of peripheral psoriatic arthritis, DIP joint involvement, or clinical enthesitis were not associated with hip joint disease. Sacroiliitis or spondylitis were strongly associated with hip disease ($p < 0.01$), as was early age of onset of PsAr ($p < 0.001$). 27% of the spondylitis patients developed hip disease. 37% of the patients with hip disease required total hip arthroplasty (THA).

Conclusion Hip joint involvement in psoriatic arthritis is strongly associated with early onset psoriatic arthritis and the presence of psoriatic spondylitis. A high risk of premature THA exists. Better strategies for early diagnosis and medical intervention are needed.

SAT0028 LATE ONSET ANKYLOSING SPONDYLITIS-CLINICAL AND RADIOLOGICAL MANIFESTATIONS

DK Kerimovic-Morina, D Jablanovic. *Clinical V, Institute of Rheumatology-Belgrade, Belgrade, Yugoslavia*

10.1136/annrheumdis-2001.380

Background

Objectives To compare clinical and radiological manifestations of late onset ankylosing spondylitis (LOAS: >51 yrs) with classical onset ankylosing spondylitis (COAS:21–30 yrs).

Methods We identified twenty-nine (2,0%) pts with LOAS out of 1434 AS pts and they were matched for sex and disease duration with COAS pts. All pts fulfilled the diagnostic criteria of AS

of the European Spondyloarthropathy study group (ESSG). Evaluation included clinical examination: inflammatory pain, spinal mobility (Schober test, chest expansion), peripheral arthritis and enthesitis, acute uveitis, family history; laboratory (ESR, HLA-B27); radiographs of the sacroiliac joints (SIJ), anteroposterior and lateral spine and affected peripheral joints). Statistical analysis was performed using Spearman rank correlation test and X-test.

Results Analysis revealed no significant differences ($p > 0.005$) in clinical manifestations between twenty-nine (27 m,2 f) pts with LOAS(mean age and disease duration 56,0 yrs (range 53–62) and 8, o yrs (range 8–12) respectively) and twenty-nine pts with COAS(mean age at onset 24,2 yrs and mean disease duration 7,8 yrs (range 4–16) respectively), except for spinal mobility ($p < 0,005$). No significant differences ($p > 0,005$) were found between groups of the laboratory findings. All pts had bilateral sacroiliitis. The results are shown in the Table 1.

The radiographic changes in SIJ (grade III and IV) were more frequent in LOAS pts than in COAS ($p < 0,005$), as well as on the spine ($p < 0,005$). The radiographic progression in the SIJ and the spine were more rapid in LOAS pts than in COAS pts, the difference was statistically significant ($p < 0,005$).

Abstract SAT0028 Table 1

	LOAS	COAS	p
Sacroiliitis grade I+II	10 (34.3%)	20 (69%)	<0.005
Sacroiliitis grade III+IV	19 (65.5%)	9 (27.5%)	<0.005
Spondylitis L + Th + C	15 (51.7%)	3 (10/3%)	<0.005

Conclusion According to our experience the onset of ankylosing spondylitis is uncommon after 51-st yr; LOAS and COAS have similar clinical manifestations; LOAS had clinically more progressive course and rapid radiographic progression leading to severe changes in the SIJ and spine ($p < 0,005$). the possible explanation for the difference observed is a clinically asymptomatic or ?silent? ankylosing spondylitis in patients with late onset of disease.

REFERENCE

- Caplanne D, Tubach F, Marie Le Parc J. Late onset spondyloarthropathy: Clinical and biological comparison with early onset patients. *Ann Rheum Dis.* 1997;**56**:176–9

SAT0029 THE ROLE OF MICROFLORA IN ANKENT (ANKYLOSING ENTHESOPATHY) IN MICE

¹Z Rehakova, ¹J Sinkora, ¹T Hrnčíř, ¹R Stepankova, ²J Capkova, ³P Ivanyi. ¹Immunology and Gnotobiology, Institute of Microbiology, Academy of Sciences of the Czech Republic, Nový Hradec; ²Institute of Molecular Genetics, Academy of Science; ³Immunogenetics, Institute of Clinical and Experimental Medicine, Prague, Czech Republic

10.1136/annrheumdis-2001.381

Background Murine ankylosing enthesopathy (ANKENT), an experimental model for human ankylosing spondylitis, is strongly dependent on the contact of ANKENT-prone mice (B10 or B10. BR genetic background) with microbes as no disease is observed under germfree (GF) conditions while about 20% of conventional (CV) males are afflicted.

The disease is almost exclusively restricted to male mice, It affects only ankle and tarsal joints. Next risk factor for ANKENT is haplotype.