Objectives To establish the incidence and nature of osteoarthritic disease in patients receiving long term treatment with haemodialysis we surveyed all of our patients who had received dialysis.

Methods The subjects were 123 patients who were dialysed for one year to 22 years. We also analysed clinical factors such as age, haemodialysis duration, underlying disorder of chronic renal insufficiency, haematologic parameters. All patients were seen at an extraclinic attendance when they were asked to evaluate pain and stiffness in each joint. A diagnosis of dialysis associated arthropathy was made in patients by three investigators, using as criteria single or combined presence of carpal tunnel syndrome, erosions and bone cysts of the joints and destructive spondylarthropathy. Plasma urea, creatinin, haemoglobin, ferritin, calcium, phosphorous, urate, parathyroid hormone concentrations, alkaline phosphatase, magnesium, aluminium, B2 microglobulin, osteocalcin, vit D, hepatitis markers were measured before dialysis in patients. Radiographs of the hands were obtained in all patients, in some patients painful joints were also examined by radiography. We used dual-energy X-Ray absorptiometry to detect changes in bone mineral density.

Results We found carpal tunnel syndrome in 29 (23.8%) patients, spondylarthropathy in 14 (11.5%) patients, skeletal deformities in 29 (23.8%) patients, tenosynovitis in 6 (4.8%) patients, peri-arthritis in 10 (8.2%) patients, peri-arthritis in 10 (8.2%) patients and bone cysts in 26 (20.7%) patients. When we compared groups according to duration of hemodialysis treatment; carpal tunnel syndrome, bone cysts and peri-arthritis were found more frequently in patients receiving hemodialysis treatment for longer time. Bone mineral density was reduced in dialysis patients of both sexes in comparison with the healthy subjects. There was significant correlation between bone mineral density and the mean serum B2 microglobulin, magnesium, creatinin and ferritin, alkaline phosphatase levels. Strong relationship was found out among hemodialysis duration and osteoporosis. Depression was observed in 66.3% patients.

Conclusion Changes of the locomotor apparatus in prolonged hemodialysis treatment determine the quality of life with all its consequences for the patient.

REFERENCES

- 1 Levenson JL, Glocheski S. Psychological factors affecting end-stage renal disease. *Psychosomatics* 1993;**32**:382-8
- 2 Ferrari AJL, Rothfuss S, Schumacher HR. Dialysis arthropathy; identification and evaluation of a subset of patients with unexplained inflammatory effusions. *J Rheumatol*. 1997;**24**(9):1780-6
- 3 Loevner LA, Adler RS, Martel W. Dialysis related arthropathy in patients on long term hemodialysis: radiographic features. *J Clin Rheumatol*. 1995;**1**:81-9

SAT0103 ACROMEGALIC ARTHROPATHY: AN ITALIAN STUDY IN 20 PATIENTS

¹F Schiavon, ¹H Marotta, ²P Maffei, ²C Martini, ²C Menegazzo, ²N Siculo, ³R Ragazzi, ¹S Todesco. ¹Division of Rheumatology, Department of Medical and Surgical Sciences, University of Padova; ²Third Medical Clinic, Department of Medical and Surgical Sciences, University of Padova, Padova, Italy; ³First Radiology Unit

10.1136/annrheumdis-2001.486

Background Acromegaly is frequently complicated by axial and peripheral joint abnormalities. Although usually mild, severe handicap is not infrequent. In addition soft tissue, cartilage and bony proliferation make acromegaly a peculiar model of non-inflammatory joint disease, which is reflected by clinical and radiological findings.

Objectives The aim of this study was to compare the clinical and radiographic features of Italian acromegalic patients with the well-known literature studies.

Methods We studied 20 patients (10 men and 10 women) with acromegaly, admitted to our Department. The diagnosis of acromegaly and the status of the disease were based on clinical, hormonal and radiological features. The duration of disease was assumed as the interval between the clinical onset and the time of the diagnosis and was 5.43 ± 2.76 year for men and 8.50 ± 5.66 year for women. Physical examination of peripheral and axial joints and radiological assessment of hands, wrists, shoulders, spine, pelvis hips and knee was performed. Moreover a ultrasonographic study of the shoulder in 8 patients was carried out. All the patients were submitted to a global self assessment of disease status or activity with WOMAC.

Results We found 70% of acromegalic gave a history of peripheral joint pain, often mild and intermittent and 65% described spine involvement. At the examination 12 patients described spine pain, mostly mild and poorly localised, 9 shoulder pain with severely disabling symptoms, 4 mild localised knee joint pain, 2 hip pain. Only 2 patients reported hand pain. Over 50% of patients presented joint hypermobility. We are able to find a correlation between the severity of peripheral joint or spinal abnormalities, WOMAC physical function and stiffness scale score and activity of acromegaly (GH and IGF-1 blood levels) ($p < 0.001$). In our patients the shoulder represents one of the most painful and disabling joint. Radiographic abnormalities included narrowing of glenohumeral joint, beak-like osteophytes and, in 4 patients, a destructive arthropathy with oval deformation of humeral head. Ultrasonographic studies were carried out in 8 patients and revealed tendons involvement in all the patients, even without radiological abnormalities. We found supraspinatus tendon tear in 3 patients, rotator cuff and bicipital tendinitis in 7 patients and subacromial bursitis in 2 patients.

Conclusion

SAT0104 ARE COAGULOPATHIES INVOLVED IN NONTRAUMATIC AVASCULAR NECROSIS?

¹N Mehse, ²JP Vernhes, ³V Guérin, ¹P Germain, ¹L Malterre, ¹J Dehais, ³F Parrot, ¹T Schaevebeke. ¹Rheumatology, ²Rheumatology, CH Libourne, Libourne, France; ³Hematology, CHU Bordeaux, Bordeaux

10.1136/annrheumdis-2001.487

Background Various conditions are considered as classical etiologies of nontraumatic avascular necrosis (AVN), such as endogenous or exogenous hypercorticism, alcoholism, barotraumas, systemic lupus, diabetes mellitus and hyperlipidemia. However, in many cases, none of these conditions are identified and AVN is considered to be idiopathic.

Objectives To seek thrombophilic disorders in a cohort of patients with AVN.

Methods All patients with AVN hospitalised in our departments during a couple of years were enrolled in this prospective study. Diagnosis of AVN was performed on X-rays and/or MRI when necessary and possible. Biological investigations included: glycaemia, cortisolemia, ferritinemia, cholesterolemia, lipoproteinemia, haemoglobin electrophoresis, ANA, antiphospholipid antibodies, anti-B2GP1 antibodies, lupus anticoagulant, antiprothrombin antibodies, antithrombin, protein C, protein S, factor VIII, IX, XI, activated protein C resistance, prothrombin gene mutation, homocysteinemia and methionin load test.