Objectives To establish the incidence and nature of osteoarthritic disease in patients receiving long term treatment with haemodialysis we surveyed all of our patients who had received dialysis.

Methods The subjects were 123 patients who were dialysed for one year to 22 years. We also analysed clinical factors such as age, haemodialysis duration, underlying disorder of chronic renal insufficiency, haematologic parameters. All patients were seen at an extraclinic attendance when they were asked to evaluate pain and stiffness in each joint. A diagnosis of dialysis associated arthropathy was made in patients by three investigators, using as criteria single or combined presence of carpal tunnel syndrome, erosions and bone cysts of the joints and destructive spondylarthropathy. Plasma urea, creatinin, haemoglobin, ferritin, calcium, phosphorous, urate, parathyroid hormone concentrations, alkaline phosphatase, magnesium, aluminium, B2 microglobulin, osteocalcin, vit D, hepatitis markers were measured before dialysis in patients. Radiographs of the hands were obtained in all patients, in some patients painful joints were also examined by radiography. We used dual-energy X-Ray absorptiometry to detect changes in bone mineral density.

Results We found carpal tunnel syndrome in 29 (23.8%) patients, spondylarthropathy in 14 (11.5%) patients, skeletal deformities in 29 (23.8%) patients, tenosynovitis in 6 (4.8%) patients, peri-arthritis in 10 (8.2%) patients, peri-arthritis in 10 (8.2%) patients and bone cysts in 26 (20.7%) patients. When we compared groups according to duration of hemodialysis treatment; carpal tunnel syndrome, bone cysts and peri-arthritis were found more frequently in patients receiving hemodialysis treatment for longer time. Bone mineral density was reduced in dialysis patients of both sexes in comparison with the healthy subjects. There was significant correlation between bone mineral density and the mean serum B2 microglobulin, magnesium, creatinin and ferritin, alkaline phosphatase levels. Strong relationship was found out among hemodialysis duration and osteoporosis. Depression was observed in 66.3% patients.

Conclusion Changes of the locomotor apparatus in prolonged hemodialysis treatment determine the quality of life with all its consequences for the patient.

REFERENCES

- 1 Levenson JL, Glocheski S. Psychological factors affecting end-stage renal disease. *Psychosomatics* 1993;**32**:382-8
- 2 Ferrari AJL, Rothfuss S, Schumacher HR. Dialysis arthropathy; identification and evaluation of a subset of patients with unexplained inflammatory effusions. *J Rheumatol*. 1997;**24**(9):1780-6
- 3 Loevner LA, Adler RS, Martel W. Dialysis related arthropathy in patients on long term hemodialysis: radiographic features. *J Clin Rheumatol*. 1995;**1**:81-9

SAT0103 ACROMEGALIC ARTHROPATHY: AN ITALIAN STUDY IN 20 PATIENTS

¹F Schiavon, ¹H Marotta, ²P Maffei, ²C Martini, ²C Menegazzo, ²N Siculo, ³R Ragazzi, ¹S Todesco. ¹Division of Rheumatology, Department of Medical and Surgical Sciences, University of Padova; ²Third Medical Clinic, Department of Medical and Surgical Sciences, University of Padova, Padova, Italy; ³First Radiology Unit

10.1136/annrheumdis-2001.486

Background Acromegaly is frequently complicated by axial and peripheral joint abnormalities. Although usually mild, severe handicap is not infrequent. In addition soft tissue, cartilage and bony proliferation make acromegaly a peculiar model of non-inflammatory joint disease, which is reflected by clinical and radiological findings.

Objectives The aim of this study was to compare the clinical and radiographic features of Italian acromegalic patients with the well-known literature studies.

Methods We studied 20 patients (10 men and 10 women) with acromegaly, admitted to our Department. The diagnosis of acromegaly and the status of the disease were based on clinical, hormonal and radiological features. The duration of disease was assumed as the interval between the clinical onset and the time of the diagnosis and was 5.43 ± 2.76 year for men and 8.50 ± 5.66 year for women. Physical examination of peripheral and axial joints and radiological assessment of hands, wrists, shoulders, spine, pelvis hips and knee was performed. Moreover a ultrasonographic study of the shoulder in 8 patients was carried out. All the patients were submitted to a global self assessment of disease status or activity with WOMAC.

Results We found 70% of acromegalic gave a history of peripheral joint pain, often mild and intermittent and 65% described spine involvement. At the examination 12 patients described spine pain, mostly mild and poorly localised, 9 shoulder pain with severely disabling symptoms, 4 mild localised knee joint pain, 2 hip pain. Only 2 patients reported hand pain. Over 50% of patients presented joint hypermobility. We are able to find a correlation between the severity of peripheral joint or spinal abnormalities, WOMAC physical function and stiffness scale score and activity of acromegaly (GH and IGF-1 blood levels) ($p < 0.001$). In our patients the shoulder represents one of the most painful and disabling joint. Radiographic abnormalities included narrowing of glenohumeral joint, beak-like osteophytes and, in 4 patients, a destructive arthropathy with oval deformation of humeral head. Ultrasonographic studies were carried out in 8 patients and revealed tendons involvement in all the patients, even without radiological abnormalities. We found supraspinatus tendon tear in 3 patients, rotator cuff and bicipital tendinitis in 7 patients and subacromial bursitis in 2 patients.

Conclusion

SAT0104 ARE COAGULOPATHIES INVOLVED IN NONTRAUMATIC AVASCULAR NECROSIS?

¹N Mehse, ²JP Vernhes, ³V Guérin, ¹P Germain, ¹L Malterre, ¹J Dehais, ³F Parrot, ¹T Schaevebeke. ¹Rheumatology, ²Rheumatology, CH Libourne, Libourne, France; ³Hematology, CHU Bordeaux, Bordeaux

10.1136/annrheumdis-2001.487

Background Various conditions are considered as classical etiologies of nontraumatic avascular necrosis (AVN), such as endogenous or exogenous hypercorticism, alcoholism, barotraumas, systemic lupus, diabetes mellitus and hyperlipidemia. However, in many cases, none of these conditions are identified and AVN is considered to be idiopathic.

Objectives To seek thrombophilic disorders in a cohort of patients with AVN.

Methods All patients with AVN hospitalised in our departments during a couple of years were enrolled in this prospective study. Diagnosis of AVN was performed on X-rays and/or MRI when necessary and possible. Biological investigations included: glycaemia, cortisolemia, ferritinemia, cholesterolemia, lipoproteinemia, haemoglobin electrophoresis, ANA, antiphospholipid antibodies, anti-B2GPI antibodies, lupus anticoagulant, antiprothrombin antibodies, antithrombin, protein C, protein S, factor VIII, IX, XI, activated protein C resistance, prothrombin gene mutation, homocysteinemia and methionin load test.

Results 17 patients were enrolled, 9 men and 8 women, aged 34 to 76, with various stages of AVN (from stage 1 to 4). Antithrombin III deficiency was diagnosed in a woman. Resistance to activated protein C was found in another woman. Hyperhomocysteinemia was identified in 6 patients: two women (associated with antiphospholipid antibodies in one case), and 4 men. A lupus anticoagulant was found in one man and one woman. Two other patients were found to be homozygous to methyl-tetrahydrofolate reductase (MTHFR) mutation, condition which leads to a susceptibility to hyperhomocysteinemia.

Conclusion 59% of our patients with AVN were found to have a thrombophilic disorder. We suggest that such a disorder may precipitate thrombosis when added to other risk factors. Among these thrombophilic disorders, hyperhomocysteinemia seemed to be the most frequent factor associated with AVN. If hyperhomocysteinemia is confirmed to be involved in AVN, these results should be of special interest in prevention of recurrences, as hyperhomocysteinemia can be corrected by folic acid, B6 and B12 supplementation.

SAT0105 LOSARTAN BUT NOT IRBESARTAN REDUCES SERUM URIC ACID IN HYPERTENSIVE PATIENTS WITH GOUT

G Wuerzner, A Chioléro, M Maillard, CL Stubi-Fallab, M Burnier, CA Gerster. *Hypertension, Lausanne University Hospital, Lausanne, Switzerland*

10.1136/annrheumdis-2001.488

Background Losartan is the only angiotensin II receptor antagonist with uricosuric properties and has been shown to decrease serum uric acid (SUA) levels in normal subjects as well as in hypertensive patients.

Objectives To assess and to compare the effects of losartan and irbesartan on serum uric acid in hypertensive patients with gout.

Methods Prospective, randomised, double-blind, cross-over study preceded by a three weeks run-in period with enalapril 20 mg od. During the first four weeks, patients received a morning dose of either losartan 50 mg or irbesartan 150 mg and a evening dose of placebo. The dose was then increased to losartan 50 mg bid or irbesartan bid. The losartan and the irbesartan phases were separated by three weeks of treatment with enalapril 20 mg od.

All drugs were provided in an electronic pill container allowing to monitor compliance (MEMS system).

Results Thirteen male hypertensive (office blood pressure > 140/90 mmHg or under antihypertensive therapy) and hyperuricemic (SUA over 420 µmol/L or over 7.1 g/dL, without any hypouricemic agents) patients completed this study.

The changes in blood pressure induced by losartan and irbesartan were comparable. In contrast to irbesartan which had no effect on SUA, losartan (50 mg od) decreased SUA significantly from 538 ± 26 µmol/L to 491 ± 20 µmol/L ($p < 0.01$) and to 484 ± 25 µmol/L at 50 mg bid. During the first 4 h after drug intake, the first dose of losartan increased the urinary uric acid/creatinine ratio when compared with irbesartan (0.525 ± 0.071 vs 0.358 ± 0.045 , $p = 0.019$). This increase was still present at week 4 (0.485 ± 0.085 vs 0.331 ± 0.057 , $p < 0.01$) but was not found at week 8 when patients received a bid regimen (0.316 ± 0.043 vs 0.331 ± 0.029 , $p = \text{ns}$). The frequency of acute gout episodes did not differ between treatments. The monitoring of compliance showed that the adherence to the morning dose was good ($89.2 \pm 2.8\%$) and significantly greater than that to the evening dose ($78.9 \pm 3.1\%$, $p < 0.001$).

Conclusion These results demonstrate the potential benefits of using losartan in hypertensive patients with hyperuricemia and gout. They also suggest that losartan-treated patients reach a new uric acid steady-state during a sustained administration (>1 month). Hence, the uricosuric effect tends to decrease with time as SUA is reduced. Increasing the dose of losartan to 50 mg bid does not appear to induce a further decrease in serum uric acid.

Pediatric rheumatology

AB0158 MUSCULOSKELETAL PAIN IN CHILDREN AND ADOLESCENTS

MM Costa, P Nero, E Branco, J Branco. *Rheumatology, Egas Moniz, Lisbon, Portugal*

10.1136/annrheumdis-2001.489

Background

Objectives The purpose of this study was the determination of the frequency and consequence of musculoskeletal pain in urban children and adolescents.

Methods This study was performed in unselected population of two urban schools. Two Rheumatologists visited 479 children (216 boys and 263 girls) in their school. Mean age was 11 ± 2.4 years, range 6–17 yr. The survey included a detailed interview on the basis of a standardised questionnaire and physical examination. The severity of their pain was assessed by visual analogue scale and a subjective disability index.

Results About 30.3 percent (145/479) of cases had musculoskeletal pain. The pain was located at lower extremity in 74% of children. Thirty-one case (21.4%) had spinal pain and 19 (13.1%) had upper extremity pain. The severity of their pain, by visual analogue scale, had a mean 3.2 ± 1.8 (range 0.1–8) and $\frac{3}{4}$ of cases had disability. Pain relapses weekly in 38% and persists part of a day in 74%. Twenty eight percent (40/145) consulted their physician and 31 children consumed drugs. The physical activity and schoolbag carriage was associated with pain in 61% (89/145) of cases.

There were no statistically significant demographic differences among cases with and without musculoskeletal pain, except for sex, as is shown in the Table 1:

Abstract AB0158 Table 1

	Age	Sex*	Sports	Weight	Height
Pain	10 ± 2.2	F = 93; M = 52	43	43.3 ± 14.4	144.6 ± 13.3
No pain	11 ± 2.5	F = 170; M = 164	107	42.9 ± 14.5	145.2 ± 14.6

* $p < 0.05$.

Conclusion In our population lower extremity pain was more prevalent. It had a relapsing character and determines disability and consumption of medical services. The relationship with physical activity (practice of sports) was referred by children and was evaluated in other study.

AB0159 JOINT HYPERMOBILITY AND MUSCULOSKELETAL PAIN

MM Costa, P Nero, E Branco, J Branco. *Rheumatology, Egas Moniz, Lisbon, Portugal*

10.1136/annrheumdis-2001.490