THU0019 INTRACELLULAR EXPRESSION OF CXCR3 ON RHEUMATOID ARTHRITIS SYNOVIAL TISSUE CELLS

¹O Krystuffkova, ¹J Vencovsky, ¹S Ruzickova, ²J Sinkora, ²Z Rehakova, ¹J Niederlova, ³CA Power, ³C Plater-Zyberk. ¹Clinical Department, Institute Rheumatology, Praha 2; ²Department Gnotobiology, Institute Microbiology AV CR, Novy Hradek, Czech Republic; ³Immunology Department, SPRI, Geneva, Switzerland

10.1136/annrheumdis-2001.528

Background Inflammatory cell infiltration and synovial activation are important processes in rheumatoid arthritis. Chemotactic gradients of various chemokines are responsible for cell attraction and possibly for their activation. We have previously detected strong expression of chemokine receptor CXCR3 in the rheumatoid joint by immunostaining.

Objectives Characterisation of the cells expressing CXCR3 in RA synovial membrane.

Methods Synovial tissue samples were obtained from RA patients undergoing either synovectomy or a total joint replacement. Cells were released by digestion with collagenase, DNase and briefly with hyaluronidase. A three colour fluorescence analysis was performed with FITC conjugated anti-CXCR3 mouse MAb (R&D) and with a panel of phycoerythrin (PE) conjugated MAb (anti-CD3, CD4, CD8, CD19, CD55, CD31, CD68, CD14 and CD45). Live cells were identified by propidium iodide. PBCs were stained using the same protocol.

Results As expected, a proportion of CD3⁺ and CD4⁺ blood and synovial cells was CXCR3 positive. In addition, CXCR3 was also seen in synovial cells positive for CD55, CD14, CD8 and to a less extent CD31. However, in contrast to the surface staining of cells from peripheral blood, CXCR3 staining was identified only intracellularly in synovial cells. No CXCR3 staining could be detected on the surface of any type of viable synovial cell, including CD3 positive lymphocytes.

Conclusion Flow cytometry identifies synovial cells that display intracellular CXCR3 staining. These cells are comprised of T lymphocytes, macrophages, possibly synovial fibroblasts and endothelial cell populations. The intracellular presence of CXCR3 suggests a possible internalisation of this molecule, which may be a consequence of ligand binding. The significance of this phenomenon and of CXCR3 expression in cell types other than leukocytes remains to be determined.

This work was supported by a grant NI/6459 from IGA MZ CR.

THU0020 ROLE OF FCGAMMARII AND FCGAMMARIII IN INFLAMMATION AND CARTILAGE DAMAGE DURING MURINE EXPERIMENTAL ANTIGEN-INDUCED ARTHRITIS

¹PL Van Lent, ¹KC Nabbe, ¹AB Blom, ¹AE Holthuisen, ¹LB Van de Putte, ²S Verbeek, ¹WB Van den Berg. ¹Rheumatology, University Medical Centre Nijmegen, Nijmegen, The Netherlands; ²Human and Clinical Genetics, University Medical Centre Leiden, Leiden, The Netherlands

10.1136/annrheumdis-2001.529

Background IgG-containing immune complexes, which are found in most RA joints, communicate with haematopoietic cells using three classes of Fc receptors (FcγRI, II, III). In a previous study we found that if an experimental arthritis was elicited in knee joints of FcR γ-chain deficient mice which lack functional FcγRI and III, joint inflammation was lower and severe cartilage destruction was absent.

Objectives To examine the role of FcγRII and III in inflammation and functional cartilage damage in knee joints with chronic experimental antigen-induced arthritis (AIA) using FcγRIII and FcγRII deficient mice.

Methods FcγRII and FcγRIII ^{-/-} mice and their controls were immunised with methylated BSA in complete adjuvant followed by induction of arthritis by local injection of mBSA into the right knee joint. The course of inflammation was studied by ^{99m}Tc uptake. Chronic inflammation and cartilage damage (depletion of proteoglycans, chondrocyte death and matrix erosion) was studied histologically in total knee joints sections stained with haematoxylin or safranin-O. Aggrecan breakdown in cartilage caused by metalloproteinases (MMP) was studied by immunolocalisation (VDIPEN neopeptides) and image analysis.

Results Three weeks after immunisation both humoral (total IgG, IgG1, IgG2a and IgG2b levels) and cellular (T cell responses as measured by lymphocyte proliferation) immunity raised against mBSA was comparable in all groups examined. IgG2a levels were comparable in FcγRIII ^{-/-} but higher in FcγRII ^{-/-} if compared to controls. Joint swelling at day 1, 3 and 7 was similar in FcγRIII ^{-/-} mice and significantly (103%) higher in FcγRII ^{-/-} on day 1 but not on later time-points. Histologically, at day 7 after AIA induction, exudate and infiltrate in the knee joint was similar in FcγRIII ^{-/-} and significantly higher (100% and 388%) in FcγRII ^{-/-} mice if compared to controls. MMP-induced neopeptides determined in various cartilage layers (tibia and femur) were comparable in FcγRIII ^{-/-} and higher (respectively 155–360% and 67–75%) in FcγRII ^{-/-}. Initial depletion of proteoglycans was similar (40–70%) in all groups. In the chronic phase, chondrocyte and cartilage matrix erosion in tibia and femur stage was significantly elevated in FcγRII ^{-/-} but not in FcγRIII ^{-/-} mice.

Conclusion FcγRIII is redundant or not involved in the presence of FcγRI whereas FcγRII is a crucial inhibiting factor in acute and chronic inflammation and cartilage erosion during antigen-induced arthritis.

REFERENCE

- 1 Van Lent, et al. *Arthritis Rheum.* 2000;**43**(4):740–52

THU0021 DEFECTIVE RAP1 ACTIVATION IN SYNOVIAL FLUID T LYMPHOCYTES FROM PATIENTS WITH RHEUMATOID ARTHRITIS

¹PH Remans, ²KA Reedquist, ¹SI Gringhuis, ¹FC Breedveld, ¹JM Van laar, ¹CL Verweij. ¹Rheumatology, Leiden University Medical Center, Leiden, The Netherlands; ²Laboratory for Physical Chemistry, Center for Biomedical Genetics, University Medical Center, Utrecht, The Netherlands

10.1136/annrheumdis-2001.530

Background Rap1 is a small G-protein, member of the Ras GTPase family. Rap1 has been linked to T cell anergy and it is suggested that Rap1 has the potential to inhibit Ras-mediated oncogenic or growth promoting activity. Synovial fluid (SF) T cells from patients with rheumatoid arthritis (RA) display a defective phosphorylation pattern of several pivotal signalling proteins and severe hyporesponsiveness upon antigenic stimulation. Our previous data showed that the hyporesponsive state of the SF T cells in RA correlates with markers of oxidative stress and replenishment of the intracellular level of the anti-oxidant (GSH) by treatment of cells with N-acetyl-cysteine (NAC) seems to partly restore the observed signalling defects.