patients. Vascularity was markedly reduced in the affected areas with relatively increased blood flow in the peripherally-spared areas. We referred to this as a “fried-egg” or “bull’s eye” appearance.

Conclusion Our study demonstrates the use of both grey scale and power Doppler sonography as a helpful non-invasive procedure in the evaluation of HIBM. Their ability to define a unique centrifugal, “fried-egg” myopathic abnormality may be diagnostic for this disorder.

OP0131 CHLOROQUINE MYOPATHY: PREVALENCE AND CLINICAL FEATURES (PRELIMINARY DATA)

¹E Casado, ¹M Larrosa, ¹J Gratacos, ²J Font, ³JM Martinez, ²C Tolosa. ¹Rheumatology; ²Medicine; ³Neurology, Corporacio Sanitaria Parc Tauli. University Hospital of Sabadell, Sabadell, Spain

10.1136/annrheumdis-2001.581

Background

Objectives To analyse the prevalence and clinical features of chloroquine myopathy in a cohort of out-patients in a Rheumatology Unit.

Methods Prospective study of 119 patients (84 female and 35 male), aged 57.8 ± 13.7 , taking antimalarials (112 chloroquine, 7 hydroxychloroquine) for more than 6 months. They were evaluated during 2 years. The underlying disease was RA in 69, connective tissue disease in 22, palindromic rheumatism in 14, spondyloarthropathy in 7 and other rheumatic disease in 7. Muscle enzymes were determined in all patients and an electromyography (EMG) was practised in patients with repeated increased determinations. Patients with altered EMG were biopsied.

Results *Laboratory*: 18 patients (15%) had a repeated increase of muscle enzymes: LDH in 17 patients, CPK in 5 and aldolase in 1 patient. EMG: practised in 13 patients, showed a muscular involvement in 6. Two patients are pending and 3 were dead before the exam. *Muscle biopsy*: 6/7 patients had a chloroquine myopathy (optic and electronic microscopy); 1/7 patients had a steroid myopathy. Two patients (RA) with chloroquine myopathy had an associated inflammatory myositis. *Clinical features*: 6 patients with chloroquine myopathy (5 RA and 1 spondyloarthropathy) represents an accumulated prevalence (2 years) of 5%. The duration of treatment was 36 ± 21 months (rest of the group 40 ± 31), with cumulative doses of 203 ± 25 g (rest of the group 305 ± 243 g). All patients presented weakness and had other simultaneous muscle involvement factors (1 cervical myelopathy, 2 hypothyroidism, 3 hypovitaminosis D and 1 prolonged bedrest) at clinical diagnosis.

Conclusion Toxic chloroquine myopathy is not an exceptional event (accumulated prevalence 5%). When an increase of LDH and/or CPK is present we should rule out this complication. The presence of other factor with muscle involvement probably makes worse the weakness and gets better its detection.

OP0132 MYOSITIS AND PREGNANCY

CA Silva, DA Isenberg. Centre for Rheumatology, University College of London, London, UK

10.1136/annrheumdis-2001.582

Background Polymyositis and dermatomyositis are the most common form of idiopathic inflammatory myopathy, overall the

female to male ratio is 2–3: 1. The age at presentation is most often between 40 and 60 years. Pregnancies in these patients are uncommon, and the overall maternal and foetal risks are not well defined in patients with myositis.

Objectives We assessed the prevalence and outcome of pregnancy in 37 patients with dermatomyositis and polymyositis.

Methods 54 patients with myositis have been treated between 1976 and 2000 at the Bloomsbury Rheumatology Unit, University College/Middlesex Hospital. A prospective record of all such patients had been kept by one of us (DAI). All patients meet the Bohan and Peter Criteria for DM and PM. The presence of another autoimmune disease was noted as appropriate. Patients were reviewed on a regular basis in a specialist outpatient clinic, and for all patients basic demographic data, age at onset, treatment, number of pregnancies, activity of the disease during pregnancy were noted. For the purposes of this study all of the available 37 patients were examined by one of us (CS): 28 females and 9 males.

Results The female patients medium age was 44 years, with age at onset ranging from 16 to 64. 16 have Dermatomyositis/Poly-myositis (7- PM/9- DM) and 12 have myositis as part of an overlap syndrome (SLE (7 patients), 2 patients with RA, 2 with SS, and 2 with Scl). Only 3 of our patients have been pregnant after the onset of the disease. One, with a poor control of her disease, had an abortion during the first trimester, and was also on treatment with MTX. The other two had the disease under control and 2 normal pregnancies with good foetal outcome was achieved. Both suspended immunosuppressants during pregnancy and were able to decrease steroid dosage, without increasing disease activity.

Conclusion Pregnancy in myositis patients is a rare event. Pregnancies occurring in myositis patients with the disease under clinical remission can result in good foetal outcome. The activity of the disease remains stable. These results are consistent with the literature from recent years, and are probable due to better control of disease activity.

Fibromyalgia

AB0166 THE REAL VALUE OF REHABILITATION IN THE MUSCULAR SYNDROMES OF THE CERVICAL SPINE

EL Sidenco. Rehabilitation, University Hospital Bucharest, Bucharest, Romania

10.1136/annrheumdis-2001.583

Background

Objectives We studied a group of 34 patients with algo-functional muscular-ligamentar cervical syndromes based on complex static disorders, diagnostically excluding cervical discal hernia. We propose a clinical -functional score for an unitary evaluation of these cases.

Methods We applied an initial kinetic programme for two weeks; we propose a complex of stretching-strengthening-coordination exercises for the cervical muscular contraction, supported by thermo-electro-massotherapy.

Results Efficiency of the initial kinetic programme was good, the following clinical parameters increased over 63,7%.

The second stage of rehabilitation was applied a month later; we used a kinetic programme analogous to the first stage, for two weeks again, insisting on the strengthening-coordination exercises for the cervical muscles. The kinetic programme of the

second stage of rehabilitation was supported by thermo-massotherapy. The results were superior to the first stage, the improvement of the clinical parameters was over 83,4%.

The third therapeutic stage was applied 3–6 months later, after a new consultation. We used the same kinetic programme of strengthening- co-ordination exercises daily, for another two weeks, thermo-massotherapy supported.

The improvement after the third stage of the kinetic programme was superior, clinical parameters increased over 90% confronted with initial values.

Conclusion We prove the necessity of an unitary score of valuation and the efficiency of a staggered kinetic programme in the complex rehabilitation of the algo-functional muscular-ligamentar cervical syndromes, as the only efficient therapeutic alternative in the functional deficiencies? compensation and for prevention of complications.

AB0167 EFFICIENCY OF REHABILITATION IN THE OLD PATIENTS PERIARTHRITIS

EL Sidenco. *Rehabilitation, University Hospital Bucharest, Bucharest, Romania*

10.1136/annrheumdis-2001.584

Background

Objectives We studied a group of 40 old patients with soft tissues sufferings (periartthritis) at the shoulder or hip. Patients accused pain and limited mobility, and at the clinical examination we found muscular hypotrophy and limitation of function in the shoulder or hip.

Methods We applied an associated kinetic programme of stretching-strengthening- control and co-ordination exercises in two different periods of 15 days, supported by electro-thermo-massotherapy. The valuations were made at the beginning and the end of the programme, on a clinical score of valuation obtained by the clinical parameters; the results were statistically processing.

Results We find from the beginning a large incidence of the periartthritis at the old patients (25,37%) and the preference for the women, larger in comparison with other ages (27 cases from the group of 40).

Application of the kinetic programme, physically supported, significantly improved the pain (completely for the spontaneous and nocturnal pain, and over 70% for the pain at the motion). Limitation of mobility increased over 63,6%, even the functional remaining is limited at the old patient and the real possibilities of the kinetic programme display can be limited by the associated pathologies. Muscular hypotrophy had a medium response (improvement under 33,7%), because of the limited biological capital and also the associated pathologies which imposed the adaptation of the kinetic programme to the real effort's possibilities. Global function of the upper limb increased over 77,5%.

Conclusion We consider the soft tissues sufferings of the old patient indicated for the kinetic programme, adequately supported by physical proceedings; the associated cardiovascular diseases do not forbid the rehabilitation, kinetic programme specially, if the patients are well medically treated and the kinetic programme is well adapted at the real effort's possibilities of the patients.

AB0168 PSYCHOPATHOLOGIC STUDY IN PATIENTS WITH FIBROMYALGIA

¹AR De la Serna, ²M Serra, ²J Wulf. ¹Rheumatology (1)/Psychosomatic (2); ²Psychosomatic, Sant Pau, Barcelona, Spain

10.1136/annrheumdis-2001.585

Background Fibromyalgia is considered a psychosomatic disease.

Objectives This study was conducted to assess psychopathology changes in patients with fibromyalgia.

Methods We evaluated sequential outpatients with ACR criteria of fibromyalgia, through interviews by clinical psychologist, a psychiatrist and test for general psychopathology (Golberg), depression (Beck) and anxiety (Stai).

Results Demographic data: Twenty-nine patients were evaluated (4 men). Twenty-seven were married. Eighteen were immigrants. Twenty-four had children. Eight were unemployed. Thirteen were house wives. Three non qualified professionals. Eleven retailers and two qualified professionals. Psychopathologic results: Eight patients were totally normal. Golberg'test: twelve patients were normal. Five had mild and twelve severe alterations. Beck'test: thirteen patients were normal. Four suffered from mild depression. Four moderate, four severe and four more severe depression. Stai'test: ten patients didnt anxiety. Eighteen were trait anxiety and sixteen state anxiety. Interview: nineteen patients had sleep problems. Twenty-four were fatigued and fifteen had relapse thinking.

Conclusion The absence of psychopathology in 28% of patients with fibromyalgia, and the mild changes in others, allows us to differentiate between psychogenic rheumatism and fibromyalgia, and obligues us to establish a psychogenic evaluation in these.

AB0169 FIBROMYALGIA. GERIATRICS PROBLEM?

A Laiz, H Corominas, M Peiró, S Fernández, C Díaz, G Vázquez. *Internal Medicine, Hospital de La Santa Creu I Sant Pau, Barcelona, Spain*

10.1136/annrheumdis-2001.586

Background

Objectives To study in a group of patients, of a rheumatology unit in primary care the presence of fibromyalgia and tocompare the results between those with more and less than 65 years.

Methods From a rheumatology unit in primary care, in Barcelona, during one year, we selected the patients with fibromyalgia.

The diagnostic was done according with ACR criteria 1990.

Statistical method used for comparisons was the chi- square test and correction of continuidad.

Results Visits performed were 1555. The mean was 61 ± 15 (Table 1).

Between them ($p < 0.05$).

Abstract AB0169 Table 1

	Less than 65	Equal or more than 65	Total
Without fibromyalgia	777	660	1437
With fibromyalgia	93	25	118
Total	870	685	1555

Conclusion The prevalence of fibromyalgia is similar to others studies.